

# CLASSICALLY INTEGRABLE COSMOLOGICAL MODELS WITH A SCALAR FIELD

Hisao Suzuki, Eiichi Takasugi and Yasuhiro Takayama

*Department of Physics, Osaka University*

*1-16 Toyonaka, Osaka 560, Japan*

## Abstract

New classes of classically integrable models in the cosmological theories with a scalar field are obtained by using freedoms of defining time and fields. In particular, some models with the sum of exponential potentials in the flat spatial metric are shown to be integrable. The model with the Sine-Gordon potential can be solved in terms of analytic continuation of the non-periodic Toda field theory.

After the original inflational scenario was presented, various models have been proposed to obtain a natural inflation such as the chaotic inflation and the modified Brans-Dicke theories. In all of these scenarios, scalar fields play an important role. In other words, in the early universe, the dynamics seems to be governed by the gravity theory coupled with a scalar field.

As for classical solutions, it is known that we can easily get them once we start by assuming the time variation of the scale factor  $a$ : The classical solution of a scalar field and the potential are expressed as functions of  $a$  and its derivatives so that in principle the potential is expressed as a function of a scalar field[1]. Moreover, equations for gauge invariant variables for scalar perturbations can be reduced to a second order homogenous differential equation whose coefficients are solely determined by  $a$  and its derivatives. By solving this equation, several analytic solutions for the inflation models are derived[1].

In this paper, we present a method to get classically integrable models for gravitational theories coupled with a scalar field in the case where the dilatational mode and scalar field do not have spatial dependence. A method to get integrable models was proposed in Ref.[2], but unfortunately the integrable models were very few. Here, we show that with a slight modification of their approach, we can get integrable models for the larger class of models. In particular, it will be shown that models with the exponential potential corresponding to the Brans-Dicke theory are classically integrable. We also find the Sine-Gordon theory coupled to gravity can be solved in terms of Toda theories.

The action of the gravity coupled to a scalar field is given by

$$S = \int d^4x \sqrt{-g} \left[ \frac{1}{16\pi G} f(\Phi) R + \frac{1}{2} (T(\Phi) g^{\mu\nu} \partial_\mu \Phi \partial_\nu \Phi - U(\Phi)) \right], \quad (1)$$

where  $f(\Phi)$ ,  $T(\Phi)$  and  $U(\Phi)$  are functions of  $\Phi$ . The standard choice of the action is  $f = 1 - \xi \Phi^2$ ,  $T = 1$  (the choice  $\xi = \frac{1}{6}$  is called minimal coupling). The modified Brans-Dicke theories correspond to the choice  $f = \Phi$ ,  $T = \omega(\Phi)/\Phi^2$ , where the ordinary Brans-Dicke theory is given by  $\omega = 1$ , and other choice is called the extended Brans-Dicke theories[3]. For  $T(\Phi) = 0$ , the scalar field is just the auxiliary field and by eliminating it with the use of equation of motion, we get the action of the higher derivative gravity[4].

It is known that the general form of the action in Eq.(1) can be transformed to a action in the Einstein frame as followings[1]: We make a conformal transformation

$$g_{\mu\nu} = \frac{1}{f(\Phi)} \tilde{g}^{\mu\nu}, \quad (2)$$

and define a new scalar field as

$$d\chi = [T(\Phi)f^{-1} - 3(\frac{f'}{f})^2]d\Phi. \quad (3)$$

Then, we change the scalar potential as

$$\tilde{U}(\chi) \equiv U(\Phi)f^4(\Phi) \quad (4)$$

and finally we get the action in the Einstein frame:

$$S = \int d^4x \sqrt{-\tilde{g}} [\frac{1}{16\pi G} R(\tilde{g}) + \frac{1}{2}(\tilde{g}_{\mu\nu} \partial_\mu \chi \partial_\nu \chi - \frac{1}{2} \tilde{U}(\chi))]. \quad (5)$$

Except for the singular case  $Tf^{-1} - 3(\frac{f'}{f})^2 = 0$  where we just obtain the action with no kinetic term of the scalar field. Hereafter, we consider the non singular case and thus we treat the action in Eq.(1) with  $T = f = 1$ .

We start from the metric of the universe to be the spatially homogeneous metric as

$$(ds)^2 = \sigma^2(N^2(dt_N)^2 - a^2(t_N)[\frac{(dr)^2}{1 - Kr^2} + r^2(d\theta)^2 + r^2 \sin^2 \theta (d\varphi)^2]), \quad (6)$$

where  $N$  is the gauge function and  $\sigma^2 = 2G/3\pi$ . We assume that  $\phi$  depends only on the time  $t_N$ . Writting

$$\phi = (2\pi^2\sigma^2)^{\frac{1}{2}}\Phi(t_N), \quad V(\phi) = 2\pi^2\sigma^4 U(\Phi), \quad (7)$$

the 1 dimensional action in Eq.(1) with  $f = T = 1$  is expressed as follows:

$$S = \frac{1}{2} \int dt_N [N^{-1}(-a\dot{a}^2 + a^3\dot{\phi}^2) + N(-a^3V(\phi) + Ka)], \quad (8)$$

where we have omitted the volume factor of the space. Note that the variation with respect to  $N$  gives us the hamiltonian constraint

$$(\frac{\dot{a}}{a})^2 - \dot{\phi}^2 - V + \frac{K}{a^2} = 0. \quad (9)$$

An important point in our method is that the equation of motion does not depend on the choice of this gauge function  $N$ . In other words, we can choose this function in such a way that the system becomes easy to be solved. This is the main advancement from the method discussed in Ref.[2].

We are now going to change the action in Eq.(8) to the one with ordinary kinetic terms. All we have to do is to change variables  $a$  and  $\phi$  to  $z$  and  $w$  in such a way that the action is transformed to

$$S = \int dt_N [(2\tilde{N})^{-1}(-\dot{z}^2 + \dot{w}^2) - \tilde{N}W], \quad (10)$$

or by introducing variables  $z_{\pm} = z \pm w$ , we write Eq.(10) as

$$S = \int dt_N [-(2\tilde{N})^{-1}\dot{z}_+\dot{z}_- - \tilde{N}W(z_+, z_-)]. \quad (11)$$

When we set  $\tilde{N} = 1$ , the cosmological problem turns out to be just an ordinary kinematical problem with indefinite metric.

In order to find new variables, we define

$$a = e^{h(z_+, z_-)}, \quad \tau_{\pm} = h \pm \phi. \quad (12)$$

Then the action in Eq.(8) is written as

$$S = \frac{1}{2} \int dt_N \left[ -\frac{e^{\frac{3}{2}(\tau_+ + \tau_-)}}{N} \dot{\tau}_+ \dot{\tau}_- + N \left[ -e^{\frac{3}{2}(\tau_+ + \tau_-)} V\left(\frac{1}{2}(\tau_+ - \tau_-)\right) + K e^{\frac{1}{2}(\tau_+ + \tau_-)} \right] \right]. \quad (13)$$

The change of variables which preserves the form of kinetic term is known as the two dimensional conformal transformation with respect to the fields  $\tau_+$  and  $\tau_-$ . We therefore take the following choice of variables:

$$\tau_+ = \tau_+(z_+), \quad \tau_- = \tau_-(z_-). \quad (14)$$

By choosing gauge function as

$$N = \tilde{N} e^{\frac{3}{2}(\tau_+ + \tau_-)} \partial_+ \tau_+ \partial_- \tau_-, \quad (15)$$

we find that the kinetic terms of Eq.(13) transforms to those of (11), as desired. Here  $\partial_+$  and  $\partial_-$  mean derivatives with respect to  $z_+$  and  $z_-$ , respectively. To simplify Eq.(15), we will use the following functions in stead of  $\tau_{\pm}$ :

$$x_+(z_+) = e^{\frac{3}{2}\tau_+}, \quad x_-(z_-) = e^{\frac{3}{2}\tau_-} \quad (16)$$

Then Eq.(15) is just written as

$$N = \frac{4}{9} \tilde{N} \partial_+ x_+ \partial_- x_- \quad (17)$$

In terms of  $x_+$  and  $x_-$ , the original fields  $a$  and  $\phi$  can be written as

$$a = (x_+ x_-)^{\frac{1}{3}}, \quad \phi = \frac{1}{3} \ln \frac{x_+}{x_-} \quad (18)$$

We can also find that the potential term in Eq.(10) or (11) is given by

$$W(z_+, z_-) = \frac{2}{9} \partial_+ x_+ \partial_- x_- [x_+ x_- V[\frac{1}{3} \ln \frac{x_+}{x_-}] - K(x_+ x_-)^{\frac{1}{3}}] \quad (19)$$

Note that the form of the potential  $W$  depends on the function  $x_+$  and  $x_-$  which are arbitrary functions of  $z_+$  and  $z_-$ , respectively, while the kinetic term takes the form of  $\dot{z}_+ \dot{z}_-$  as seen in Eq.(11). We utilize the freedom of choosing functions of  $x_+$  and  $x_-$  to simplify the potential  $W$  such that differential equations are solved.

Firstly, we consider the exponential potential with  $K = 0$ :

$$[A]: \quad V(\phi) = \lambda e^{-\alpha\phi} \quad (\alpha \neq \pm 6)$$

This case is equivalent to the Brans-Dicke theory with a constant potential and is known to possess a power law inflational solution  $a = a_0(t + c)^\beta, \beta = \frac{12}{\alpha^2}$ , as a special solution. The special case of  $\alpha = \pm 2$  in this case is also the special case of [H] whcih we discuss later. The case [H] is well studied and the analytic solution is obtained[5]. Thus,  $\alpha = \pm 2$  case was known to be soluble. But for other cases, the analytic solutions are not known before.

Here we show that the general solution is obtained in our method. We take the ansatz

$$x_+ = z_+^l \quad x_- = z_-^m \quad (20)$$

When we take  $l$  and  $m$  as

$$l = \frac{i}{2 - \frac{\alpha}{3}}, \quad m = \frac{j}{2 + \frac{\alpha}{3}} \quad (21)$$

with  $i$  and  $j$  are integers, then the potential  $W$  is written as

$$W(z_+, z_-) = \frac{2\lambda}{36 - \alpha^2} i j z_+^{i-1} z_-^{j-1}. \quad (22)$$

These are valid for  $\alpha \neq \pm 6$ . When the choice of the pair  $(i, j)$  is one of (1,1), (1,2), (2,1), (1,3), (2,2) and (3,1), the theory can be solved by the separation of variables. In particular, if we take  $i = j = 1$ , the potential becomes constant. With the gauge choice  $\tilde{N} = 1$ , the equations of motion are written by  $\ddot{z}_+ = \ddot{z}_- = 0$  so that

$$z_+ = a_1 t_N + a_2 \quad z_- = b_1 t_N + b_2. \quad (23)$$

The Hamiltonian constraint  $\frac{\dot{z}_+ \dot{z}_-}{2} - W = 0$  gives a restriction

$$a_1 b_1 = \frac{4\lambda}{36 - \alpha^2}. \quad (24)$$

The scale factor  $a$  and the scalar field  $\phi$  are obtained from Eq.(18) as functions of  $t_N$  through Eq.(23). From Eq.(17) with  $\tilde{N} = 1$ , the time  $t_N$  is expressed as a function of  $t$  as

$$t = \frac{4}{36 - \alpha^2} \int (a_1 t_N + a_2)^{l-1} (b_1 t_N + b_2)^{m-1} dt_N, \quad (25)$$

where  $l$  and  $m$  are given in Eq.(21) with  $i = j = 1$ . In order to compare the power law behavior given before, we have to perform the above integration. Here we consider the large  $t_N$  or  $t$  limit of  $a$  and  $\phi$ . From Eq.(25), we have  $t \propto t_N^{l+m-1}$ . Then,  $a = z_+^{\frac{l}{3}} z_-^{\frac{m}{3}} \propto t_N^{(l+m)/3} \propto t^{\frac{12}{\alpha^2}}$  which reproduces the special solution stated before.

With the ansatz in Eq.(20), some models with potentials consisting of the sum of exponential forms are solvable. We found the following cases with  $K = 0$  are solvable:

$$[B] : V(\phi) = \lambda_1 e^{2\phi} + \lambda_2 e^{-2\phi}$$

We find with the ansatz in Eq.(20)

$$W = \frac{1}{8}(\lambda_1 z_+ + \lambda_2 z_-) \quad \text{with} \quad l = m = \frac{3}{4}. \quad (26)$$

This case is the special case of [H] explained later so that this model was known to be soluble[5].

$$[C] : V(\phi) = \lambda_1 e^{\frac{5}{18}\phi} + \lambda_2 e^{-\frac{6}{5}\phi}$$

We find with the ansatz in Eq.(20)

$$W = \frac{25}{144}(\lambda_1 z_+ + \lambda_2 z_-^2) \quad \text{with} \quad l = \frac{5}{8}, \quad m = \frac{5}{4}. \quad (27)$$

Note that a new soluble model is derived from a model by changing  $\phi$  to  $-\phi$ . In this case,  $W$  is obtained by changing  $z_+$  and  $z_-$  and  $l$  and  $m$ . Thus, the case

$$[C'] : V(\phi) = \lambda_1 e^{-\frac{18}{5}\phi} + \lambda_2 e^{\frac{6}{5}\phi}$$

is also soluble and with the ansatz in Eq.(20) we find

$$W = \frac{25}{144}(\lambda_1 z_- + \lambda_2 z_+^2) \quad \text{with} \quad l = \frac{5}{4}, \quad m = \frac{5}{8}. \quad (28)$$

$$[D] : V(\phi) = \lambda_1 e^{3\phi} + \lambda_2 e^{-3\phi} + \lambda_3$$

We find with the ansatz in Eq.(20)

$$W = \frac{2}{9}(\lambda_1 z_+^2 + \lambda_2 z_-^2 + \lambda_3 z_+ z_-) \quad \text{with} \quad l = m = 1. \quad (29)$$

Solutions for all of above cases can be derived following the same manner demonstrated for the exponential potential case. The special cases of [D], the case with  $\lambda_1, \lambda_2 > 0$  and  $\lambda_3 = -2\sqrt{\lambda_1 \lambda_2}$  and the case with  $\lambda_3 = 0$  are solved by Ritis et. al.[2].

As for the exponential potential in the case [A],  $\alpha = \pm 6$  are excluded. Here we consider these cases:

$$[E] : V(\phi) = \lambda e^{-6\phi}$$

By taking the ansatz

$$x_+ = e^{Az_+}, \quad x_- = z_+^{\frac{1}{4}}, \quad (30)$$

we get

$$W = \frac{1}{18}A\lambda. \quad (31)$$

Thus, this case is solvable. The case with  $\alpha = -6$  can be solved by exchanging  $x_+$  into  $x_-$ , i.e., by taking  $x_+ = z_+^{\frac{1}{4}}$  and  $x_- = e^{Az_-}$ .

Let us examine some other type of ansatz,

$$x_+ = e^{Az_+}, \quad x_- = e^{Bz_-}. \quad (32)$$

There are several solvable cases under this ansatz. One is the potential

$$[\text{F}] : V(\phi) = \lambda_1 e^{\alpha\phi} + \lambda_2 e^{\beta\phi}, \quad \alpha\beta = 36 \quad (\alpha \neq \beta)$$

In this case, with the choice

$$B = \frac{2 + \frac{\alpha}{3}}{2 - \frac{\alpha}{3}} A, \quad (33)$$

the potential reduces to

$$W = \frac{2}{9} AB \left( \lambda_1 \exp[2A(2 + \frac{\alpha}{3})z] + \lambda_2 \exp[\frac{12}{\alpha} A(2 + \frac{\alpha}{3})w] \right). \quad (34)$$

Another nontrivial example which cannot be solved by the separation of variable is the case of a Sine-Gordon potential

$$[\text{G}] : V(\phi) = \lambda \cos 6\sqrt{3}\phi$$

In this case, we take

$$x_+ = (288)^{\frac{1}{2}} e^{-\frac{1}{8}z_+}, \quad x_- = (288)^{\frac{1}{2}} e^{-\frac{1}{8}z_-}. \quad (35)$$

Then, we get the following potential:

$$W = \lambda e^{-\frac{1}{2}z} \cos \frac{\sqrt{3}}{2} w. \quad (36)$$

In this case, we can find the following conserved currents:

$$H = -\frac{1}{2}p_z^2 + \frac{1}{2}p_w^2 + \lambda e^{-\frac{1}{2}z} \cos \frac{\sqrt{3}}{2} w = 0, \quad (37)$$

$$I = \frac{1}{2}p_z^2 p_w + \frac{1}{6}p_w^3 + \frac{\sqrt{3}}{2} \lambda p_z e^{-\frac{z}{2}} \sin \frac{\sqrt{3}}{2} w + \frac{\lambda}{2} p_w e^{-\frac{z}{2}} \cos \frac{\sqrt{3}}{2} w, \quad (38)$$

where  $p_z$  and  $p_w$  are conjugate momentum of  $z$  and  $w$ , respectively. Since we found two conserved currents, the model is soluble. The origin of the higher dimensional conserved current is understood by noticing that the system can be obtained by the analytic



continuation of the nonperiodic Toda theory[6]. Explicitly, the conserved quantities  $H$  and  $I$  are constructed from a Lax pair  $(L, M)$  of the nonperiodic Toda theory: That is  $H \propto \text{Tr} L^3$  and  $I \propto \text{Tr} L^3$  with the identification  $q_1 = -(z + \frac{i}{\sqrt{3}}\omega)/4$ ,  $q_2 = \frac{i}{2\sqrt{3}}\omega$  and  $q_3 = -(q_1 + q_2)$  where  $q_1$ ,  $q_2$  and  $q_3$  are defined in Ref.6.

Finally, we consider the integrable models with  $K \neq 0$ . This case is in general hard to be solved. Except for a trivial example  $V = \Lambda$ , we can find the soluble model with the potential,

$$[\text{H}] : V(\phi) = \lambda_1 e^{2\phi} + \lambda_2 e^{-2\phi} \quad (K \neq 0)$$

Similarly to the case [B], we take the ansatz

$$x_+ = z_+^{\frac{3}{\alpha A \otimes 4}}, \quad x_- = z_-^{\frac{3}{4}}, \quad (39)$$

we obtain

$$W = \frac{1}{8}(\lambda_1 z_+ + \lambda_2 z_- - K) \quad (40)$$

so that this case is soluble. This is a famous model studied extensively[5] and is known to be soluble.

It may be worthwhile to note that our result depends heavily on the fact that we introduced only one scalar field. In this case, we can utilize the  $2d$  conformal transformation with respect to  $a$  and  $\phi$  and we found several solvable models. If we have  $n$  scalar fields and wish to use similar symmetry, we may need to consider  $n+1$  dimensional conformal transformation. Thus, the integrable models are quite restrictive except for  $n=1$  because the 2 dimensional conformal transformation is known to be special. Note also that the basic form of the soluble potential is of the exponential type. These are quite analogous to the integrability of 2 dimensional gravity.

We have found some class of cosmological models can be solved in terms of the appropriate choice of the gauge function and the conformal transformation. But the integrable models are rare and most of the physically interesting examples are not integrable. For example, the potential which corresponds to  $R^2$  gravity [7] can be transformed to the

quartic potential by an appropriate choice of the conformal transformation. Of course, since we do not know the whole classification of integrable models even in the ordinary kinematic problem, we have an opportunity to get other integrable models. We hope larger class of interesting models can be found and analyzed in our method.

Finally, we point out that for some potentials given above, the Wheeler-De Witt equation can be solved analytically. This equation for the case [H] ( and thus for the case [A] with  $\alpha = \pm 2$  and the case [B]) was solved and the analytical solution is given[5]. By using our transformations, we can get the analytical solutions for the cases [A], [D], [E] and [F] which will be given in the forthcoming paper.

## References

- [1] M. Kuwahara, H. Suzuki and E. Takasugi, Phys. Rev. **D50** (1994) 661.
- [2] R. de Ritis et. al., Phys. Rev. **D42** (1990) 1091; Phys. Lett. **A149** (1990) 79.
- [3] D. La and P.J. Steinhardt, Phys. Rev. Lett. **62** (1989) 376;  
P.J. Steinhardt and F.S. Acetta, Phys. Rev. Lett. **64** (1990) 2740;  
J. Garcia-Bellido and M. Quiros, Phys. Lett. **B 243** (1990) 45.
- [4] B. Whitt, Phys. Lett. **B 145** (1984) 176.
- [5] L.J. Garay, J.J. Halliwell and G.A. Marugan, Phys. Rev. **D43** (1991) 2572.
- [6] M.A. Olshanetsky and A.M. Perelomov, Phys. Rep. [bf C71 No.5 (1981) 313.
- [7] D.S. Salopek, J.R. Bond and J.M. Bardeen, Phys. Rev. **D40** (1989) 1753.