

Unitarity Condition on Quantum Fields in Semiclassical Gravity

Sang Pyo Kim *

Department of Physics

Kunsan National University

Kunsan 573-701, Korea

Abstract

The condition for the unitarity of a quantum field is investigated in semiclassical gravity from the Wheeler-DeWitt equation. It is found that the quantum field preserves unitarity asymptotically in the Lorentzian universe, but does not preserve unitarity completely in the Euclidean universe. In particular we obtain a very simple matter field equation in the basis of the generalized invariant of the matter field Hamiltonian whose asymptotic solution is found explicitly.

Published in Physics Letters A **205**, 359 (1995)

Unitarity of quantum field theory in curved space-time has been a problem long debated but still unsolved. In particular the issue has become an impassioned altercation with the discovery of the Hawking radiation [1] from black hole in relation to the information loss problem. Recently there has been a series of active and intensive investigations of quantum

*E-mail : sangkim@knusun1.kunsan.ac.kr

effects of matter field through dilaton gravity and resumption of unitarity and information loss problem (for a good review and references see [2]).

In this letter we approach the unitarity problem and investigate the condition for the unitarity of a quantum field from the point of view of semiclassical gravity based on the Wheeler-DeWitt equation [3]. By developing various methods [4–18] for semiclassical gravity and elaborating further the new asymptotic expansion method [19] for the Wheeler-DeWitt equation, we derive the quantum field theory for a matter field from the Wheeler-DeWitt equation for the gravity coupled to the matter field, which is equivalent to a gravitational field equation and a matrix equation for the matter field through a definition of cosmological time. The full field equation for the matter field is found to preserve unitarity asymptotically in the limit $\frac{\hbar}{M} \rightarrow 0$ in the Lorentzian universe with an oscillatory gravitational wave function but does not preserve unitarity completely for an exponential gravitational wave function in the Euclidean universe. We find the exact quantum state for the asymptotic field equation in terms of the eigenstates of the generalized invariant of the matter field Hamiltonian.

We consider the Wheeler-DeWitt equation for a quantum cosmological model

$$\left[-\frac{\hbar^2}{2M} \nabla^2 - MV(h_a) + \hat{H}\left(\frac{i}{\hbar} \frac{\delta}{\delta \phi}, \phi, h_a\right) \right] \Psi(h_a, \phi) = 0. \quad (1)$$

Here M is the Planck mass squared, $\nabla^2 = G_{ab}(\delta^2)/(\delta h_a \delta h_b)$, V denotes the superpotential of three-curvature with or without the cosmological constant on superspace with the DeWitt metric G_{ab} with the signature $(-, +, \dots, +)$, and \hat{H} represents the matter field Hamiltonian. We find the wave function of the form

$$\Psi(h_a, \phi) = \psi(h_a) \Phi(\phi, h_a), \quad (2)$$

where $\Phi(\phi, h_a)$ is a gravitational field-dependent quantum state of the matter field. We expand the quantum state by some orthonormal basis

$$\Phi(\phi, h_a) = \sum_k c_k(h_a) |\Phi_k(\phi, h_a)\rangle, \quad \langle \Phi_k | \Phi_n \rangle = \delta_{kn}. \quad (3)$$

Substituting Eqs. (2) and (3) into Eq. (1) and acting $\langle \Phi_n |$ on both sides, one obtains the following matrix equation equivalent to the Wheeler-DeWitt equation

$$\begin{aligned}
c_n(h_a) \left(-\frac{\hbar^2}{2M} \nabla^2 - MV(h_a) + H_{nn}(h_a) \right) \psi(h_a) - \frac{\hbar^2}{M} \nabla \psi(h_a) \cdot \nabla c_n(h_a) \\
+ i \frac{\hbar^2}{M} \nabla \psi(h_a) \cdot \sum_k \mathbf{A}_{nk}(h_a) c_k(h_a) + \psi(h_a) \sum_{k \neq n} H_{nk}(h_a) c_k(h_a) \\
- \frac{\hbar^2}{2M} \psi(h_a) \sum_k \Omega_{nk}(h_a) c_k(h_a) = 0,
\end{aligned} \tag{4}$$

where

$$\begin{aligned}
H_{nk}(h_a) &= \langle \Phi_n(h_a) | \hat{H} | \Phi_k(h_a) \rangle, \\
\mathbf{A}_{nk}(h_a) &= i \langle \Phi_n(h_a) | \nabla | \Phi_k(h_a) \rangle, \\
\Omega_{nk}(h_a) &= \nabla^2 \delta_{nk} - 2i \mathbf{A}_{nk} \cdot \nabla + \Omega_{nk}^{(2)},
\end{aligned} \tag{5}$$

where

$$\Omega_{nk}^{(2)}(h_a) = \langle \Phi_n(h_a) | \nabla^2 | \Phi_k(h_a) \rangle. \tag{6}$$

By noting that $i\nabla$ acts on $c_k(h_a)$ as a hermitian operator, it follows that H , \mathbf{A} , $\Omega^{(2)}$, and thereby Ω are hermitian matrices.

We separate the matrix equation equivalent to the Wheeler-DeWitt equation into the gravitational part

$$\left(-\frac{\hbar^2}{2M} \nabla^2 - MV(h_a) + H_{nn}(h_a) \right) \psi(h_a) = 0, \tag{7}$$

and the matter field part

$$\begin{aligned}
- \frac{\hbar^2}{M} \nabla \psi(h_a) \cdot \nabla c_n(h_a) + i \frac{\hbar^2}{M} \nabla \psi(h_a) \cdot \sum_k \mathbf{A}_{nk}(h_a) c_k(h_a) \\
+ \psi(h_a) \sum_{k \neq n} H_{nk}(h_a) c_k(h_a) - \frac{\hbar^2}{2M} \psi(h_a) \sum_k \Omega_{nk}(h_a) c_k(h_a) = 0.
\end{aligned} \tag{8}$$

The gravitational wave function has an effective potential $MV - H_{nn}$. The gravitational wave function can have either a real action in a region of the Lorentzian universe or an imaginary action in a region of the Euclidean universe.

First, we consider a region of the Lorentzian universe, in which the gravitational wave function takes the form

$$\psi(h_a) = f(h_a) \exp\left(\frac{i}{\hbar} S_{nn}(h_a)\right). \quad (9)$$

In the semiclassical limit $\hbar \rightarrow 0$, the gravitational action satisfies the Einstein-Hamilton-Jacobi equation with the quantum back-reaction

$$\frac{1}{2M} (\nabla S_{nn}(h_a))^2 - MV(h_a) + H_{nn}(h_a) = 0. \quad (10)$$

There are possibly an infinite number of the gravitational wave functions depending on the mode number due to the quantum back-reaction of the matter field. Each gravitational wave function, whose peak corresponds to a classical solution with the matter field, describes a history of evolution of the universe and contains the whole information of the universe. So along each wave function in this region we may introduce a cosmological time

$$\frac{\partial}{\partial \tau^{(n)}} := \frac{1}{M} \nabla S_{nn}(h_a) \cdot \nabla. \quad (11)$$

Substituting (9) into (8) and dividing $\psi(h_a)$, we obtain the matter field equation

$$i\hbar \frac{\partial}{\partial \tau^{(n)}} c_n + \Omega_{nn}^{(1)} c_n + \sum_{k \neq n} \left(\Omega_{nk}^{(1)} - H_{nk} \right) c_k + \frac{\hbar^2}{2M} \sum_k \Omega_{nk}^{(3)} c_k = 0, \quad (12)$$

where

$$\Omega_{nk}^{(1)} = i\hbar \langle \Phi_n | \frac{\partial}{\partial \tau^{(n)}} | \Phi_k \rangle = \frac{\hbar}{M} \nabla S_{nn} \cdot \mathbf{A}_{nk}, \quad (13)$$

and

$$\Omega_{nk}^{(3)} = \Omega_{nk} + 2\frac{1}{f} \nabla f \cdot \nabla \delta_{nk} - 2i\frac{1}{f} \nabla f \cdot \mathbf{A}_{nk}. \quad (14)$$

The full field equation (12) for the matter field is not unitary due to the terms $2\frac{1}{f} \nabla f \cdot \nabla \delta_{nk} - 2i\frac{1}{f} \nabla f \cdot \mathbf{A}_{nk}$ which do not obviously act as unitary operators, even though all the other terms $\Omega^{(1)}$, H_{nk} , and Ω_{nk} are hermitian matrices. The unitarity violating terms were discovered using different method in Ref. [13]. In a previous paper [19], we also obtained a small unitarity violating term, $-i\frac{\hbar}{2M} \nabla^2 S_{nn}(h_a)$, which originated from the gravitational wave function in the form $\psi(h_a) = \exp(\frac{i}{\hbar} S_{nn}(h_a))$.

However, it is only in the asymptotic limit $\frac{\hbar}{M} \rightarrow 0$ that for any gravitational wave function satisfying (7) these unitarity violating terms are suppressed and one gets

$$i\hbar \frac{\partial}{\partial \tau^{(n)}} c_n + \Omega_{nn}^{(1)} c_n + \sum_{k \neq n} \left(\Omega_{nk}^{(1)} - H_{nk} \right) c_k = 0. \quad (15)$$

In this case, since $H^\dagger = H$, $\Omega^{(1)\dagger} = \Omega^{(1)}$, and $\Omega^\dagger = \Omega$, it follows that

$$\frac{\partial}{\partial \tau^{(n)}} \left(\mathbf{c}^\dagger \cdot \mathbf{c} \right) = 0, \quad (16)$$

where \mathbf{c} denotes a column vector of c_k . The norm of vector, $|\mathbf{c}|$, is preserved. This implies a unitary operator such that

$$\mathbf{c}(\tau^{(n)}) = \mathbf{U}_c(\tau^{(n)}, \tau_0^{(n)}) \mathbf{c}(\tau_0^{(n)}). \quad (17)$$

And since the eigenstates of the generalized invariant form an orthonormal basis, there is also a unitary operator

$$|\Phi(\tau^{(n)})\rangle = \mathbf{U}_\Phi(\tau^{(n)}, \tau_0^{(n)}) |\Phi(\tau_0^{(n)})\rangle \quad (18)$$

for the evolution of the column vector of $|\Phi_k\rangle$. The physical implication is that the quantum field theory of the matter field is asymptotically unitary in this sense.

In particular, in terms of the eigenstates of the generalized invariant

$$\begin{aligned} \frac{\partial}{\partial \tau^{(n)}} \hat{I} - \frac{i}{\hbar} [\hat{I}, \hat{H}] &= 0, \\ \hat{I} |\Phi_k\rangle &= \lambda_k |\Phi_k\rangle \end{aligned} \quad (19)$$

there is a well-known decoupling theorem [20] such that

$$H_{nk} = \Omega_{nk}^{(1)}, \quad n \neq k. \quad (20)$$

In the basis of the eigenstates of the generalized invariant the matter field equation takes the simpler form

$$i\hbar \frac{\partial}{\partial \tau^{(n)}} c_n + \Omega_{nn}^{(1)} c_n + \frac{\hbar^2}{2M} \sum_k \Omega_{nk}^{(3)} c_k = 0. \quad (21)$$

In the asymptotic limit of $\frac{\hbar}{M} \rightarrow 0$ we obtain the asymptotic quantum state of matter field [18]

$$\Phi(\phi, h_a) = c_n(\tau_0^{(n)}) \exp\left(\frac{i}{\hbar} \int \Omega_{nn}^{(1)}(h_a) d\tau^{(n)}\right) |\Phi_n(\phi, h_a)\rangle. \quad (22)$$

With an additional phase factor $\exp\left(-\frac{i}{\hbar} \int H_{nn} d\tau^{(n)}\right)$ (22) is also one of the exact quantum states of time-dependent Schrödinger equation

$$i\hbar \frac{\partial}{\partial \tau^{(n)}} \Phi(\phi, h_a) = \hat{H}\left(\frac{i}{\hbar} \frac{\delta}{\delta \phi}, \phi, h_a\right) \Phi(\phi, h_a). \quad (23)$$

The exact quantum state (3) can be obtained solving perturbatively (21), which is a linear superposition of eigenstates of the generalized invariant with gravitational field-dependent coefficient functions.

Second, we consider an exponential gravitational wave function of the form in the Euclidean universe

$$\psi(h_a) = f(h_a) \exp\left(-\frac{1}{\hbar} S_{nn}(h_a)\right). \quad (24)$$

The cosmological time is imaginary and given by

$$\begin{aligned} \frac{\partial}{\partial \tau} &= i \frac{1}{M} \nabla S_{nn} \cdot \nabla \\ &: = -i \frac{\partial}{\partial \tau_{im}^{(n)}}. \end{aligned} \quad (25)$$

The matter field equation becomes

$$\hbar \frac{\partial}{\partial \tau_{im}^{(n)}} c_n + \Omega_{nn}^{(1)} c_n + \sum_{k \neq n} \left(\Omega_{nk}^{(1)} - H_{nk} \right) c_k + \frac{\hbar^2}{2M} \sum_k \Omega_{nk}^{(3)} c_k = 0. \quad (26)$$

Even though the basis evolves unitarily, the coefficient functions in (3) with respect to which the exact quantum state is expanded do not. This violates completely the unitarity of exact quantum state.

As an application, we consider a free massive scalar field coupled to the gravity. Through the definition of cosmological time (11) using the gravitational wave function the Hamiltonian for the matter field depends on the time as

$$\hat{H}(h_a) = \frac{1}{m(\tau)} \frac{\pi_\phi^2}{2} + m(\tau) \omega^2(\tau) \frac{\phi^2}{2}. \quad (27)$$

It is found that the one-parameter-dependent generalized invariant

$$\hat{I}(\tau) = g_-(\tau) \frac{\pi_\phi^2}{2} + g_0(\tau) \frac{\pi_\phi \phi + \phi \pi_\phi}{2} + g_+(\tau) \frac{\phi^2}{2}, \quad (28)$$

is given by

$$\begin{aligned} g_-(\tau) &= c_1 \left(\phi_1^2(\tau) + \phi_2^2(\tau) \right), \\ g_0(\tau) &= c_1 m(\tau) \left(\phi_1(\tau) \dot{\phi}_2(\tau) + \phi_2(\tau) \dot{\phi}_1(\tau) \right), \\ g_+(\tau) &= c_1 m^2(\tau) \left(\dot{\phi}_1^2(\tau) + \dot{\phi}_2^2(\tau) \right) \end{aligned} \quad (29)$$

in the Lorentzian universe [21,22], and by

$$\begin{aligned} g_-(\tau) &= c_1 \left(\phi_1^2(\tau) - \phi_2^2(\tau) \right), \\ g_0(\tau) &= i c_1 m(\tau) \left(\phi_1(\tau) \dot{\phi}_2(\tau) - \phi_2(\tau) \dot{\phi}_1(\tau) \right), \\ g_+(\tau) &= -c_1 m^2(\tau) \left(\dot{\phi}_1^2(\tau) - \dot{\phi}_2^2(\tau) \right) \end{aligned} \quad (30)$$

in the Euclidean universe [23], where

$$\ddot{\phi}_{1,2}(\tau) + \frac{\dot{m}(\tau)}{m(\tau)} \dot{\phi}_{1,2}(\tau) + \omega^2(\tau) \phi_{1,2}(\tau) = 0. \quad (31)$$

It should be remarked that the generalized invariant for a harmonic oscillator in the Euclidean universe leads to an upside-down oscillator.

Finally, we apply the new method to the well-known one-dimensional Klein-Gordon equation

$$\left[-\frac{\hbar^2}{c^2} \frac{\partial^2}{\partial t^2} + \hbar^2 \frac{\partial^2}{\partial x^2} - m^2 c^2 \right] \Psi(t, x) = 0. \quad (32)$$

Here the asymptotic parameter is c the speed of light. Since the Hamiltonian $\hat{H} = \frac{\partial^2}{\partial x^2}$ is independent of parameter time t , we may take the generalized invariant $\hat{I} = \hat{H}$. The eigenstates of the generalized invariant are

$$|\Phi_k(\phi)\rangle = \exp(\pm i k x). \quad (33)$$

Not only the gauge potential A_{nk} but also Ω_{nk} do vanish. The expectation value gives rises to $H_{nk} = -\hbar^2 k^2 \delta_{kn}$. The equation that corresponds to Eq. (7) becomes

$$\left[\frac{\hbar^2}{c^2} \frac{\partial^2}{\partial t^2} + \hbar^2 k^2 + m^2 c^2 \right] \psi(t) = 0, \quad (34)$$

whose solution is

$$\psi(t) = \exp(\pm i\omega t), \quad \omega = \sqrt{k^2 c^2 + \frac{m^2 c^4}{\hbar^2}}. \quad (35)$$

It is to be noted that the new method introduced in this paper gives directly the exact result at the asymptotic limit. This argument can also be verified for a gravitational field-independent $\hat{H}(i\frac{\delta}{\hbar\delta\phi}, \phi)$.

In this letter as a semiclassical gravity we recovered an asymptotic unitary theory of the matter field in curved space in the Lorentzian universe and a non-unitary theory in the Euclidean universe, even though the quantum gravity based on the Wheeler-DeWitt equation is not a complete theory. It is expected that the unitarity of quantum field of matter field may be restored for any complete theory of quantum gravity.

Quite recently Kiefer [17] has shown that the quantum field theory derived from the Wheeler-DeWitt equation is non-unitary in general and Kiefer *et al* [13] have suggested its implication in black hole evaporation. The big difference between this paper and theirs is that we used the new method in which one separates the Wheeler-DeWitt equation into the gravitational wave equation with the quantum back-reaction of the matter field and the matrix equation for the matter field, whereas they used the conventional method in which one separates the Wheeler-DeWitt equation into the Einstein-Hamilton-Jacobi equation for the gravitational field without the quantum back-reaction of the matter field and the time-dependent functional Schrödinger equation for the matter field including higher order quantum gravitational correction. As shown in the Klein-Gordon equation in a flat space-time our method gives the relativistic frequency (35) directly at the lowest order whereas the conventional method would expand the energy as $E = mc^2 + \frac{p^2}{2mc^2} - \frac{p^4}{8m^3c^2} + \dots$. The difference between the new method and the conventional method was also discussed in various quantum cosmological models in Ref. [19].

The author would like to thank S. Carlip, S. -W. Kim, D. La, and K. Maeda for useful discussions. This work was supported by the Korea Science and Engineering Foundation under Grant No 951-0207-056-2.

REFERENCES

- [1] S. W. Hawking, Commun. Math. Phys. 43 (1975) 199.
- [2] D. N. Page, in: Proceedings of the 5th Canadian Conference on General Relativity and Relativistic Astrophysics, eds. R. B. Mann and R. G. Mclenaghan (World Scientific, 1994).
- [3] T. Banks, Nucl. Phys. B249 (1985) 332.
- [4] R. Brout, Found. Phys. Phys. 17 (1987) 603.
- [5] C. Kiefer, Class. Quantum Grav. 4 (1987) 1369.
- [6] T. P. Singh and T. Padmanabhan, Ann. Phys. 196 (1989) 296.
- [7] R. Brout and G. Venturi, Phys. Rev. D 39 (1989) 2436.
- [8] T. P. Singh, Class. Quantum Grav. 7 (1990) L149.
- [9] R. Balbinot, A. Barletta, and G. Venturi, Phys. Rev. D 41 (1990) 1848.
- [10] C. Kiefer and T. P. Singh, Phys. Rev. D 44 (1991) 1067.
- [11] J. P. Paz and S. Sinha, Phys. Rev. D 44 (1991) 1038.
- [12] C. Gundlach, Phys. Rev. D 48 (1993) 1700 (1993).
- [13] C. Kiefer, R. Müller, and T. P. Singh, gr-qc/9308024 (1993).
- [14] D. P. Datta, Mod. Phys. Lett. A 8 (1993) 191; 2523(E).
- [15] S. P. Kim and S.-W. Kim, Phys. Rev. D 49 (1994) R1679.
- [16] S. P. Kim, Phys. Lett. A 191 (1994) 365.
- [17] C. Kiefer, in: Canonical gravity - from classical to quantum, eds. J. Ehlers and H. Friedrich (Springer, Berlin, 1994).
- [18] S. P. Kim and S.-W. Kim, Phys. Rev. D 51 (1995) 4254.

- [19] S. P. Kim, Phys. Rev. D 52 (1995) 3382.
- [20] H. R. Lewis, Jr. and W. B. Riesenfeld, J. Math. Phys. 10 (1969) 1458.
- [21] S. Kim, J. Phys. A: Math. Gen. 27 (1994) 1387.
- [22] J. Y. Ji, J. K. Kim and S. P. Kim, Phys. Rev. A 51 (1995) 4268.
- [23] S. P. Kim, in: Proceedings of the 1st Seoul Workshop on Gravity and Cosmology, ed. S.-W. Kim (Journ. Korean Phys. Soc. Suppl., to appear)