

**Isothermal spherical perfect fluid model:
Uniqueness and Conformal mapping**

Naresh Dadhich*

Inter University Centre for Astronomy & Astrophysics

P.O. Box 4, Pune-411007, India

Abstract

We prove the theorem: The necessary and sufficient condition for a spherically symmetric spacetime to represent an isothermal perfect fluid (barotropic equation of state with density falling off as inverse square of the curvature radius) distribution without boundary is that it is conformal to the “minimally” curved (gravitation only manifesting in tidal acceleration and being absent in particle trajectory) spacetime.

PACS numbers : 0420, 9880.

* E-mail : naresh@iucaa.ernet.in

Except for Brinkman's theorem [1] conformally relating two Einstein spaces, only spacetimes conformal to flat spacetime have been considered. The well-known examples of such spacetimes, that are also of great astrophysical and cosmological interest, are the Friedman-Robertson-Walker (FRW) model of the Universe and the Schwarzschild solution describing interior of a star in hydrostatic equilibrium. These spacetimes have distinguishing physical properties, like isotropy and homogeneity for the former and uniform density for the latter, but no direct linkage of them with conformal flatness has been shown. In this note we shall establish a unique association between isothermality of fluid with conformal character of spacetime. Such a clear characterisation of physical behaviour with geometric property is rather very rare.

Here the base spacetime is of course not flat but "minimally" curved. It is a spherically symmetric spacetime from which radial acceleration has been annulled out but curvature is not zero, which manifests in tidal acceleration for transverse motion. Though it is not a solution of the Einstein equation, but it presents an interesting physical situation which is free of gravity at the linear (Newtonian) level. Further it has all but one curvatures zero, which is also an invariant for spherical symmetry and it is at any given radius proportional to curvature of sphere of that radius [2]. This is why we have termed it as "minimally" curved consideration of MCS [3-4]. We have very recently considered metric in the Kerr-Schild (KS) form with a view to find perfect fluid solutions [5]. It is however well-known that perfect fluid is not compatible with the KS form and hence Senovilla et al [6- 7] have generalized it by replacing flat metric by conformally flat. This would mean that metric can be written as conformal to an original KS metric, though the base space will not in general be a solution of the Einstein equation.

A metric in KS form is given by

$$g_{ij} = \eta_{ij} + 2Hl_i l_j \quad (1)$$

where H is a scalar field and l_i is a null vector relative to both g_{ij} and η_{ij} . For spherically symmetric vacuum solution; i.e. the Schwarzschild solution, H satisfies the Laplace equation $\nabla^2 H = 0$. It is rather curious to note that spacetime is not flat unless $H = 0$; i.e. $H = \text{const.} \neq 0$, represents a curved spacetime and in particular MCS. Let us consider the smallest deviation from flat spacetime by taking H constant. Recall that for perfect fluid seed metric has to be different from vacuum. So we take H constant and write

$$\bar{g}_{ij} = e^{2U}(\eta_{ij} + 2Hl_i l_j), \quad H = \text{const.} \quad (2)$$

which for spherical symmetry can be brought to the orthogonal form to read [5]

$$ds^2 = e^{2U}(-dt^2 + k^2 dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2). \quad (3)$$

Here $U = U(r, t)$ and $k^2 = (1 + 2H)^2 / (1 - 2H)$ and the base metric is MCS.

First, the perfect fluid conditions imply $U = U(r)$, and they yield the general solution for unbounded distribution [5],

$$e^U = r^{-n} \quad (4)$$

with

$$8\pi\rho = \frac{n(n-2)}{k^2 r^{2(1-n)}}, \quad \rho \frac{n-2}{n} p \quad (5)$$

where $k^2 = 1 + 2n(n-2)$ and removable constants have been transformed away. This is the general solution which represents isothermal fluid as it admits a barotropic equation of state and density falls off as inverse square of the curvature radius,

$R = r^{1-n}$. It is important to note that the metric (3) admits this unique isothermal distribution [5].

Thus the metric ansatz (3) with the unique solution (4) is the sufficient condition for isothermal perfect fluid model. We shall now turn to the necessary condition. For that we have to prove that all spherically symmetric isothermal fluid solutions can always be cast in the metric ansatz (3). Let us consider the general spherically symmetric metric

$$ds^2 = -e^\nu dt^2 + e^\lambda dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2) \quad (6)$$

where λ and ν are to begin with functions of both r and t . Now $R_{01} = 0$ will imply $\lambda = \lambda(r)$. With this we have for perfect fluid

$$8\pi\rho = \frac{1}{r^2}[1 + e^{-\lambda}(r\lambda' - 1)] \quad (7)$$

$$8\pi p = \frac{1}{r^2}[-1 + e^{-\lambda}(r\nu' + 1)] \quad (8)$$

$$2\nu'' + \nu'^2 - \lambda'\nu' - \frac{2}{r}(\nu' + \lambda') + \frac{4}{r^2}(e^\lambda - 1) = 0 \quad (9)$$

where $\lambda' = \partial\lambda/\partial r$. Now implimenting isothermality; i.e. $\rho = \gamma p \sim r^{-2}$, which will first determine $\nu = \nu(r)$, and then we obtain the general solution [8]

$$e^\lambda = \text{const.} = k_1^2, \quad e^\nu = r^{-2m}. \quad (10)$$

This belongs to the Tolman class of solutions [9]. Thus we have obtained the general solution for unbounded spherically symmetric isothermal fluid and the metric (6) will read

$$ds^2 = r^{-2m}[-dt^2 + k_1^2 r^{2m} dr^2 + r^{2(1+m)}(d\theta^2 + \sin^2\theta d\varphi^2)]. \quad (11)$$

By redefining the radial coordinate as $\bar{r} = r^{1+m}$, the above metric on dropping overhead bar takes the form

$$ds^2 = r^{\frac{-2m}{1+m}}[-dt^2 + k_1^2 dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2)] \quad (12)$$

which is exactly in the required ansatz form (3). We have thus proved the theorem: The necessary and sufficient condition for a spherically symmetric spacetime to represent an isothermal perfect fluid distribution without boundary is that it is conformal to the minimally curved spacetime.

It is quite remarkable that there exists a one to one association between isothermality and conformally MCS. Isothermality of fluid picks up uniquely the conformally MCS. This is really very interesting because it is a rare case of a specific physical property singling out a geometric property. Like the isothermal case, stresses generated by K in MCS also fall off as $T_0^0 = T_1^1 \sim r^{-2}$. These stresses have been identified with the geometric string dust [10] as well as with gravitational monopole [11]. These are very exotic objects which are relevant in the very extraordinary setting of the very early Universe. The simplest way to get to MCS is that it is a spacetime corresponding to a constant potential. Recall that H in (1) satisfies the Laplace equation. However we confess that its interpretation as potential may not be generally agreed upon. This as well as all other related issues have been taken up in detail in refs. [3-4].

Viewing MCS as a small departure (as it differs from flat spacetime only in tidal acceleration for transverse motion) from flat spacetime, it is interesting that for its conformal spacetime it does not permit non-static perfect fluid solution. It admits the general solution having three free parameters, which can represent bounded fluid

spheres as well as the unbounded cosmological model considered above. In contrast conformally flat metric admits non- conformally MCS is highly constrained and has the unique spherically symmetric perfect fluid solution. Even though MCS may be a small deviation from flat spacetime, their conformal spacetimes have very different physical properties.

Isothermal fluid structures have been considered in astrophysics for a long time as an equilibrium approximation to more complicated systems approaching dynamical relaxed state. Recently the spacetime (11) has been argued as an ultimate end state of the Einstein-deSitter universe [8]. It is envisioned that the Einstein-deSitter model asymptotically tends to an expansion free state, and then it condenses into a stable isothermal fluid sphere.

Finally it is perhaps for the first time such a clear linkage between a certain physical property of matter and a specific geometric property of spacetime has been demonstrated. It is also for the first time that spacetime conformal to a non-flat spacetime has been considered.

References

- [1] H W Brinkman Math. Ann. **91**, 269 (1924).
- [2] N Dadhich Ph.D. thesis (Pune University), (1970).
- [3] N Dadhich On the Schwarzschild field and a little beyond, submitted.
- [4] N Dadhich and L K Patel, On the Machian generalization of vacuum spacetimes, to be submitted.
- [5] N Dadhich, A conformal mapping and isothermal perfect fluid, submitted.
- [6] J M M Senovilla and C F Sopuerta Class. Quantum Grav. **11**, 2073 (1994).
- [7] J Martin and J M M Senovilla J. Math. Phys. **27**, 265 (1986).
- [8] W C Saslaw, S D Maharaj and N Dadhich, An isothermal universe, submitted.
- [9] R C Tolman Phys. Rev. **55**, 365 (1934).
- [10] P S Letelier Phys. Rev. D 20, 1294 (1979).
- [11] M Barriola and A Vilenkin Phys. Rev. Lett. **63**, 341 (1989).