

INSTABILITY OF SPACE-TIME DUE TO EUCLIDEAN WORMHOLES

V.A.Rubakov and O.Yu. Shvedov

*Institute for Nuclear Research of the Russian Academy of Sciences,
60-th October Anniversary Prospect 7a, Moscow 117312, Russia*

and

*Sub-department of Quantum Statistics and Field Theory,
Department of Physics, Moscow State University
Vorobievsky gory, Moscow 119899, Russia*

November 7, 2017

gr-qc/9608065

Abstract

The problem of topology change transitions in quantum gravity is discussed. We argue that the contribution of the Giddings-Strominger wormhole to the Euclidean path integral is pure imaginary. This is checked by two techniques: by the functional integral approach and by the analysis of the Wheeler-De Witt equation. We present also a simple quantum mechanical model which shares many features of the system consisting of parent and baby universes. In this simple model, we show that quantum coherence is completely lost and obtain the equation for the effective density matrix of the "parent universe".

This talk concerns the contribution of wormhole configurations, fig.1, into the Euclidean functional integral for the forward "flat space \rightarrow flat space" amplitude in quantum gravity. In many models, there exist classical Euclidean wormhole solutions, which may be considered as saddle points of this integral. The question is whether their contributions are real or imaginary, i.e. whether wormholes are analogous to instantons or to bounces in quantum field theory, and what could be their interpretation in the latter case.

The simplest wormhole solution [1] emerges in the theory of gravity interacting with massless three-form field (axion), or, equivalently [2], of the theory of massless scalar field $\theta(x)$ minimally coupled to gravity. The wormhole solution is $O(4)$ – symmetric,

$$ds^2 = N^2(\tau)d\tau^2 + R^2(\tau)d\Omega_3^2, \quad (1)$$

where

$$\dot{\theta} = \frac{iq}{2\pi^2 R^3 \kappa^{1/2}}, \quad \dot{R}^2 = 1 - \frac{q^2}{24\pi^4 R^4}, \quad N = 1. \quad (2)$$

Here τ is the coordinate running into the wormhole, $d\Omega_3^2$ is metrics of the unit 3-sphere, κ is the gravitational constant, and q is an arbitrary parameter of the solution, which has meaning of the global charge flowing into the wormhole. The values of the field θ are purely imaginary, so that the wormhole contributes to the amplitude not in coordinate, but in momentum representation, since the momentum $p(x)$ conjugate to $\theta(x)$ is real [2, 3, 4].

The analysis of $O(4)$ – symmetric fluctuations has been recently performed in ref.[5]. It has been found that there exists a negative mode about the wormhole solution. It has been argued, therefore, that the wormhole contribution to the forward amplitude is imaginary. It was suggested in ref.[5] that the latter fact means instability of flat space-time with respect to spontaneous emission of baby universes which in turn implies the loss of quantum coherence in the large universe, as originally suggested in refs.[6, 7]. Further support of this point of view comes from the semiclassical analysis of the Wheeler-DeWitt equation for baby universe that branches off [8]. The latter analysis reveals the role of the boundary conditions for the wave function of the baby universe that has to be imposed at singularity, $R = 0$.

There are a few subtleties concerning the diagram of fig.1. The purpose of this talk is to discuss some of these subtleties in turn.

1. For arbitrary positions of the wormhole ends in flat space-time, the values of the field θ evaluated along the flat space and along the wormhole do not match each other. In other words, the field θ has to be multivalued.

The resolution of this subtlety is in fact well-known [3]. Let us consider only those wormholes that carry fixed global charge q . This is accomplished by inserting the δ -function:

$$1 = \int dq \delta\left(\int_S p(\mathbf{x})d\mathbf{x} - q\right).$$

into the Euclidean functional integral, where S is the 3-sphere shown in fig.1, and $p(\mathbf{x})$, as before, is momentum conjugate to θ . Representing this δ -function as $\delta(\xi) = (2\pi)^{-1} \int d\lambda e^{i\lambda\xi}$, one finds that the saddle-point equation for θ is

$$\dot{\theta} = \lambda\delta(\tau - \tau_S) + \text{smooth function};$$

this means that θ may indeed have the discontinuity on the surface S .

2. The wormhole ends can be placed at any two points x and y in flat space-time. At large $|x - y|$ the wormhole action is independent of $|x - y|$. Therefore, the

wormhole contribution is proportional to $(VT)^2$ rather than usual VT , where V and T are normalization spatial volume and time, respectively.

This infrared peculiarity has to do with infinite degeneracy of vacuum of the field θ in infinite flat space: there exists a state with zero energy for any value of the global charge q . To see the infrared nature of the factor $(VT)^2$, consider finite volume case (fig.2). One can show that at large time separation, $|x^0 - y^0| \gg V^{1/3}$, the wormhole action has the following dependence on $|x^0 - y^0|$:

$$S \sim \frac{q^2 |y^0 - x^0|}{2V}$$

Hence in finite volume, the integral over the positions of the wormhole ends ("almost zero modes"), $\int dx dy e^{-S/\kappa}$, is proportional to T . We conclude that in the theory with infrared cut-off, the dependence of the wormhole amplitude on normalization time is exactly as it should be.

3. The integration over the $O(4)$ – symmetric fluctuations about the wormhole solution, as it stands, is ill defined because of the wrong sign of the action for conformal factor in quantum gravity. To cure this problem, the Gibbons-Hawking-Perry rotation [9] has been performed in ref. [5]. Even though GHP prescription works nicely in many cases, additional arguments confirming the conclusion that the wormhole amplitude of fig.1 is imaginary, are of importance.

One of these arguments comes from the observation that at large separation between the wormhole ends the diagram of fig.1 can be cut as shown in fig.3. Since the negative mode found in ref.[5] is $O(4)$ – symmetric, it suffices to evaluate the latter diagram in the minisuperspace approximation, leaving $R(\tau)$, $N(\tau)$, $\theta(\tau)$ and $p(\tau)$ as the only integration variables. Then the diagram of fig.3 is proportional to the product $\Phi_+(R, q)\Phi_-(R, q)$ of the two semiclassical wave functions, each of which is represented by the functional integral:

$$\begin{aligned} \Phi_{\pm}(R, q) = & \int DN(\tau)DR(\tau)D\theta(\tau)Dp(\tau) \\ & \times \exp \left(-\frac{1}{\kappa} S_{gr}(R(\cdot), N(\cdot)) - \frac{1}{\kappa} \int_{-\infty}^{\tau_f} d\tau \left(\frac{p^2 N}{4\pi^2 R^3} - i\sqrt{\kappa} \theta \dot{\pi} \right) \right), \end{aligned} \quad (3)$$

where S_{gr} is the gravitational action for metrics of the form (1), the boundary conditions are

$$p(\tau_f) = q, \quad (4)$$

(we consider the wormholes of global charge q),

$$R(\tau_f) = R, \quad R(-\infty) = +\infty. \quad (5)$$

The functional integral (3) is to be evaluated in the semiclassical approximation; Φ_- and Φ_+ contain the contributions of the classical solutions $R_-(\cdot)$ and $R_+(\cdot)$ (obeying

boundary conditions (4), (5)) that do and do not pass through the turning point ($\dot{R} = 0, \dot{\theta} = 0$), respectively.

Integrating eq.(3) over θ , one obtains the factor $\delta(\dot{p})$. This means that $p = \text{const} = q$, and the integration over p is immediate. The remaining integral is

$$\Phi = \int DRDN e^{-S/\kappa},$$

with

$$S = \int_{-\infty}^{\tau_f} d\tau \left[-6\pi^2 NR \left(1 - \frac{R}{N} \frac{d}{d\tau} \left(\frac{\dot{R}}{N} \right) - \frac{\dot{R}^2}{N^2} \right) + \frac{q^2 N}{4\pi^2 R^3} \right] - \frac{6\pi^2}{N} R^2(\tau_f) \dot{R}(\tau_f).$$

where we have taken into account the boundary term in the gravitational action. To evaluate the contributions of the two saddle-point configurations, one first integrates over R by the Maslov technique [10] and then evaluates the integral over N by using the following definition of measure, which coincides with the definition of ref.[11],

$$\int DNF \left(\int N d\tau \right) \equiv \int dTF(T)$$

The result is

$$\Phi_{\pm}(R, q) = \frac{1}{\sqrt{-\dot{R}_{\pm}(\tau_f)}} e^{-S[R_{\pm}(\cdot)]/\kappa}. \quad (6)$$

Since \dot{R}_+ and \dot{R}_- have opposite signs, the product

$$\Phi_+ \Phi_-$$

is pure imaginary. This is the direct consequence of the fact that the original Giddings-Strominger solution (2) has a turning point. We conclude that the amplitude of fig.1 is indeed purely imaginary.

4. The above argument is close in spirit to one of ref.[8]. Each half of the diagram of fig.3 may be viewed as a contribution to the wave function of the "ground state" of the system "parent universe + baby universes". Indeed, one can argue (cf. [12]) that the latter can be expressed through the Euclidean functional integral over 4-manifolds with the following boundary conditions: final 3-geometry is the argument of the wave function, while initial 3-geometry is the classical ground state (flat space in our case, not a point as in ref.[12]). The semiclassical wave function of the system has therefore the form

$$\Phi(R, q) = \Phi_+(R, q) + \Phi_-(R, q).$$

Then one can construct the conserved current in minisuperspace,

$$j_R = -i(\Phi \partial_R \Phi^* - \Phi^* \partial_R \Phi)$$

that measures the flux of baby universes towards the singularity. From eq.(6) it follows that

$$j_R = e^{-S_{GS}/\kappa}, \quad (7)$$

where S_{GS} is the euclidean action of the wormhole of fig.1. The fact that j_R is non-zero is interpreted as a signal that the flat space is unstable with respect to the emission of baby universes that then evolve in their intrinsic time towards the singularity.

5. The latter argument suggests also that the picture of unstable flat space is valid beyond the minisuperspace approximation. Indeed, in the general case, there still exists a current that is conserved in superspace [13]. Its semiclassical value integrated over the surface of fixed R in superspace is still expected to be given by eq.(7), since there is non-zero flux of baby universes towards $R = 0$ if the "radiation" boundary conditions are imposed. Therefore, one expects that the amplitude of fig.1 will still be pure imaginary when $O(4)$ – asymmetric fluctuations about the wormhole are properly accounted for. This expectation has to be confirmed by direct analysis of general fluctuations about the wormhole solution.

6. The above arguments support the picture of baby universes that are steadily emitted by the parent universe and then evolve into the singularity (or towards $R = \infty$ in other models [14, 15]). This picture is similar to one emerging in (1+1)-dimensional stringy model of parent and baby universes [16, 17]. In these cases one naturally expects that quantum coherence in the parent universe is lost. However, the latter property has never been shown explicitly. Here we present a simple model which shares many features of the above systems; the simplicity of the model enables one to see the loss of quantum coherence explicitly and even obtain the equation for the effective density matrix in the parent "universe".

Let us consider a quantum mechanical system (an analog of the "parent universe") which is able to emit particles ("baby universes"). The radiated particles can move along one-dimensional line in one direction (to the right) only. We choose the Hamiltonian of each particle to be equal to $-i\partial/\partial x$, so that all emitted particles move with the velocity equal to 1. Suppose that the particles can be emitted only at $x = 0$, so that the full Hamiltonian of the system under consideration is

$$H = T + \int dx a^+(x) \left(-i\frac{\partial}{\partial x}\right) a^-(x) + V a^+(0) + V^+ a^-(0). \quad (8)$$

Here $a^\pm(x)$ are creation and annihilation operators of "baby universes", operator T corresponds to free evolution of the "parent universe", while V is the operator in the "parent universe" state space which corresponds to emission of "baby universes".

Our purpose is to find whether there is loss of quantum coherence. The answer to this question depends on the boundary conditions imposed on the radiated particles. If at the initial moment of time there existed "baby universes" at $x = x_0 < 0$, they would move to the right, $x(t) = x_0 + t$, and at time $t = |x_0|$ they would reach the point $x = 0$ and would be able to interact with the "parent universe" according to

eq.(8). We see that the information about the "baby universes" residing at $x < 0$ is not lost.

For example, if $V = V^+$ then there exist so-called " α -states" [18]

$$(a^+(0) + a^-(0))\Psi = \alpha\Psi;$$

$$\int dx a^+(x) \left(-i \frac{\partial}{\partial x}\right) a^-(x) \Psi = 0.$$

If the state of the full system is an α -state at the initial moment of time, then the same property is satisfied at arbitrary time. The effective evolution of the "parent universe" is described then by the Hamiltonian $T + \alpha V$, where the "constant of nature" α is not specified by the theory but depends on the state of "baby universes". We see that Coleman's " α -state" interpretation of the Hamiltonian (8) is certainly possible.

However, one can consider another boundary condition which leads to the loss of coherence for the "parent universe". This condition is that there are no "baby universes" at $x < 0$,

$$a^-(x)\Psi = 0, x < 0. \quad (9)$$

This condition is invariant under time evolution, since "baby universes" are able to move only to the right and are created only at $x = 0$. If the condition (9) is imposed, the information about "baby universes" is lost completely, as opposed to the " α -state" case. Therefore, the initial reduced density matrix ρ_0 of the "parent universe" determines in a unique fashion its effective density matrix ρ at arbitrary time. We show in Appendix that the evolution equation for ρ is a special case of the equation of refs.[19, 20]:

$$\dot{\rho} = -i[T, \rho] - \frac{1}{2}V^+V\rho - \frac{1}{2}\rho V^+V + V\rho V^+. \quad (10)$$

It is straightforward to see [19] that $Tr\rho^2$ decreases with time, so that quantum coherence is indeed lost for the "parent universe" subsystem.

The authors are indebted to T. Banks, K. Lee, Kh. Nirov and P. Tinyakov for helpful discussions. This work is supported in part by the Russian Foundation for Basic Research, project 96-02-17449a.

Appendix

The purpose of this appendix is to obtain the evolution equation for the reduced density matrix of the "parent universe", eq.(10). Consider the interaction representation

$$\Phi(t) = e^{iH_0t}\Psi(t) \quad (11)$$

of the state vector $\Psi(t) = e^{-iHt}\Psi_0$; here

$$H_0 = T + \int dx a^+(x) \left(-i \frac{\partial}{\partial x}\right) a^-(x).$$

Since

$$e^{iH_0 t} a^\pm(\xi) e^{-iH_0 t} = a^\pm(\xi - t),$$

the vector $\Phi(t)$ satisfies the following equation

$$i\dot{\Phi}(t) = H_{int}(t)\Phi(t), \quad (12)$$

where

$$V(t) = e^{iTt} V e^{-iTt},$$

$$H_{int}(t) = V(t)a^+(-t) + V^+(t)a^-(-t).$$

Let us show that under conditions (9) the operator $H_{int}(t)$ can be replaced by the following effective Hamiltonian,

$$H_{eff}(t) = V(t)a^+(-t) - \frac{i}{2}V^+(t)V(t) \quad (13)$$

Note that the second term of eq.(13) is a direct analog of the imaginary part of the forward "flat space \rightarrow flat space" amplitude discussed in this talk.

To check the formula (13), let us show that the following vector

$$Texp(-i \int_0^t H_{eff}(t) dt) \Phi_0$$

satisfies eq.(12) if the condition (9) is imposed on the initial state vector Φ_0 . The left hand side of eq.(12) takes the form

$$H_{eff}(t)\Phi(t), \quad (14)$$

while the right-hand side of eq.(12) is as follows:

$$\sum_{n=0}^{\infty} (V(t)a^+(-t) + V^+(t)a^-(-t)) \int_{t > t_1 > \dots > t_n > 0} dt_1 \dots dt_n (-i)^n \times$$

$$\times (V(t_1)a^+(-t_1) - \frac{i}{2}V^+(t_1)V(t_1)) \dots (V(t_n)a^+(-t_n) - \frac{i}{2}V^+(t_n)V(t_n)) \Phi_0. \quad (15)$$

Because of conditions (9), the terms with $[a^-(-t), a^+(-t_i)]$ give $\delta(t - t_i)$. If $i \geq 2$ then the range of t_1 collapses to a point, so that the contributions of terms containing $a^-(-t)a^+(t_i)$, $i \geq 2$ vanish. The only non-vanishing contribution is given by the term $[a^-(-t), a^+(-t_1)]$. We now have to define the integrals like

$$\int_{t_2 < t_1 \leq t} \delta(t_1 - t) V^+(t) V(t_1) dt_1. \quad (16)$$

To resolve this ambiguity, let us regularize the Hamiltonian (8) by the replacement

$$a^+(0) \rightarrow \int a^+(\xi) f(\xi) d\xi,$$

where f is a sharply peaked real function. After this regularization the quantity

$$[a^-(-t), a^+(-t_1)] = \delta(t - t_1)$$

is replaced by

$$\int d\xi d\xi_1 [a^-(-t + \xi), a^+(-t_1 + \xi_1)] f(\xi) f(\xi_1) = \int d\xi f(\xi) f(t_1 - t + \xi).$$

The integral (16) takes the form $\frac{1}{2}V^+(t)V(t)$, so that the quantity (15) coincides with (14).

Making use of the effective Hamiltonian (13), it is straightforward to obtain eq.(10). Namely, consider the trace with respect to "baby universes" of the matrix $|\Phi(t)\rangle\langle\Phi(t)|$

$$\rho_\Phi(t) = Tr_{baby} |\Phi(t)\rangle\langle\Phi(t)|.$$

Since

$$Tr_{baby} a^+(-t) |\Phi(t)\rangle\langle\Phi(t)| = 0,$$

and

$$Tr_{baby} a^+(-t_1) |\Phi(t_1)\rangle\langle\Phi(t_2)| a^-(-t_2) = \delta(t_1 - t_2) \rho_\Phi(t_1),$$

the difference between matrices ρ_Φ at times $T + t$ and T is

$$\begin{aligned} \rho_\Phi(T + t) - \rho_\Phi(T) = & t \left[-\frac{1}{2} V^+(T) V(T) \rho_\Phi(T) - \frac{1}{2} \rho_\Phi(T) V^+(T) V(T) + \right. \\ & \left. + V(T) \rho_\Phi(T) V^+(T) \right] + O(t^2). \end{aligned}$$

Making use of formula (11) one obtains the equation for the reduced density matrix, eq. (10).

References

- [1] S.B.Giddings and A. Strominger, *Nucl. Phys.* **B306** (1988) 890.
- [2] K. Lee, *Phys. Rev. Lett.* **61** (1988) 263.
- [3] S. Coleman and K. Lee, *Nucl. Phys.* **B329** (1990) 387.
- [4] L.F. Abbott and M.B. Wise, *Nucl. Phys.* **B325** (1989) 687.
- [5] V.A. Rubakov and O.Yu. Shvedov, gr-qc/9604038, to appear in *Phys. Lett. B*.
- [6] S.W. Hawking, *Phys. Lett.* **B195** (1987) 337;
in: *Quantum Gravity. Proc. 4th Int. Seminar, Moscow, 1987*, eds. M.A. Markov, V.A. Berezin and V.P. Frolov (*World Scientific, Singapore*, 1988).

- [7] G.V. Lavrelashvili, V.A. Rubakov and P.G. Tinyakov, *Nucl. Phys.* **B299** (1988) 757;
in: Quantum Gravity. Proc. 4th Int. Seminar, Moscow, 1987, eds. M.A. Markov, V.A. Berezin and V.P. Frolov (*World Scientific, Singapore*, 1988).
- [8] O.Yu. Shvedov, gr-qc/9603039, to appear in Proceedings of the 24th ITEP Winter School of Physics.
- [9] G.W. Gibbons, S.W. Hawking and M.J. Perry, *Nucl. Phys.* **B138** (1978) 141.
- [10] V.P. Maslov, *Teor. Mat. Fiz.* **2** (1970) 30.
- [11] E. Mottola, *J. Math. Phys.* **36** (1995) 2470.
- [12] J.Hartle and S.W.Hawking, *Phys. Rev.* **D28** (1983) 2960.
- [13] B. DeWitt, *Phys. Rev.* **160**, (1967) 1113.
- [14] G.V. Lavrelashvili, V.A. Rubakov and P.G. Tinyakov, *Mod. Phys. Lett.* **A3** (1988) 1231.
- [15] V.A.Rubakov and P.G.Tinyakov, *Phys. Lett.* **B214** (1988) 334.
- [16] V.A. Rubakov, *Nucl. Phys.* **B453** (1995) 395.
- [17] Kh.S. Nirov and V.A. Rubakov, *Mod. Phys. Lett.* **A10** (1995) 3059.
- [18] S.Coleman, *Nucl. Phys.* **B307** (1988) 867.
- [19] T.Banks, L.Susskind and M.Peskin, *Nucl. Phys.* **B244** (1984) 125.
- [20] J.Ellis, J.S.Hagelin, D.V. Nanopoulos and M.Srednicki, *Nucl. Phys.* **B241** (1984) 381.




Fig.1




Fig.2




Fig.3