

accepted for publication in
Phys. Lett. B

MPI-PhT/96-69

Group Theoretical Quantization of Schwarzschild and Taub-NUT

Helia Hollmann¹

Max-Planck-Institut für Physik, Werner-Heisenberg-Institut
Föhringer Ring 6, 80805 Munich, Germany

Abstract

Stationary spherically symmetric gravity is equivalent to a nonlinear coset sigma model on $SL(2, \mathbb{R})/SO(2)$ coupled to a gravitational remnant. Classically there are stationary solutions besides the static Schwarzschild metric labeled by the Schwarzschild mass m and the Taub-NUT charge l . Imposing the $SL(2, \mathbb{R})$ symmetry at the quantum level the Wheeler-DeWitt equation becomes related to the Casimir operator on the coset, which makes the system amenable to exact quantization.

¹e-mail: hollmann@mppmu.mpg.de

The quantization of Einstein's gravity is one of the outstanding and most challenging areas of theoretical physics. Freezing all but a few of the infinitely many degrees of freedom of the gravitational field, solutions for special sectors of Einstein's vacuum theory are within reach [11], [12], [14], [5]. It is believed that at least some relevant features of the quantum theory are preserved. Even restricting oneself to a particularly symmetric case, that is to stationary spherical symmetry — as it is done here — the space of classical solutions contains black holes. They are in some respect the true particles of general relativity: coupling a point particle to its own gravitational field turns it into a black hole [6]. Furthermore classical black holes are stable under local perturbations [4], and they have remarkable uniqueness properties [3]. This is why they are considered to be the fundamental pieces of a quantum theory of gravitation, and the investigation of the spherically symmetric sector of Einstein's theory of gravitation is seen in a different light.

This paper mainly splits into a classical and a quantum part. In the beginning the spherically symmetric sector of Einstein's gravity is introduced and a Kaluza-Klein like reduction is performed [2]. Classically the theory can be solved, and the physical parameters turn out to be the Taub-NUT charge l and the Schwarzschild mass m . In the following the problem is treated as a constrained system. The observables are identified and shown to form the Lie algebra $\mathfrak{sl}(2, \mathbb{R})$. The Hilbert space of the model is built from the unitary irreducible representations of the $\mathrm{SL}(2, \mathbb{R})$ symmetry group. The quantum mechanical model can be solved completely.

One starts with the following parametrization of the four dimensional line element

$$ds^2 = \tau(dt - B_m dx^m)^2 - \tau^{-1} h_{mn} dx^m dx^n.$$

As in Kaluza-Klein theories the existence of a timelike Killing vector field can be used to perform a dimensional reduction [10] of Einstein's gravity. It leads to three dimensional gravity coupled to an abelian vector field and a scalar. After elimination of the field strength F_{mn} of B_m by the Bianchi identity

$$\tau^2 F_{mn} = \epsilon_{mnp} \partial^p \omega$$

one arrives at a nonlinear $\mathrm{SL}(2, \mathbb{R})/\mathrm{SO}(2)$ sigma model coupled to gravity:

$$\mathcal{L} = \sqrt{h} \left(-\frac{R}{2} + \frac{h^{mn}}{8} \mathrm{Tr} \left(\chi^{-1} \partial_m \chi \chi^{-1} \partial_n \chi \right) \right), \quad (1)$$

where

$$\chi = \begin{pmatrix} \tau + \frac{\omega^2}{\tau} & \frac{\omega}{\tau} \\ \frac{\omega}{\tau} & \frac{1}{\tau} \end{pmatrix}.$$

h_{mn} denotes the three dimensional metric, \sqrt{h} the square root of the determinant, R is the scalar curvature associated with h_{mn} , and χ is an element

of the Riemannian space $SL(2, \mathbb{R})/SO(2)$. τ is the norm of the Killing vector and plays the role of a gravitational potential. ω is the so called twist or NUT potential, which can also be considered as a kind of “magnetic potential” in analogy with Maxwell’s theory. To further simplify the problem one imposes an additional symmetry on the remaining three spatial dimensions:

$$h_{mn}dx^m dx^n = N^2 d\rho^2 + f^2 d\Omega^2.$$

Then the Lagrangian (1) simplifies to

$$\mathcal{L} = N \left(\frac{f'^2}{N^2} + 1 - \frac{f^2}{4N^2\tau^2}(\tau'^2 + \omega'^2) \right).$$

The additional $SO(3)$ symmetry reduces the equations of motion to ordinary nonlinear differential equations [15], [7], which can be integrated. The space of classical solutions consists of the Schwarzschild metric and a one parameter family connecting the Taub-NUT solution with the Schwarzschild solution. The geometrical difference between these is that the latter admits an isometry group whose action on four dimensional space-time has three dimensional orbits, whereas they are 2-spheres for the Schwarzschild solution. Another interpretation is given in terms of the difference between staticity and stationarity [3]: In the static case there are two symmetries: a time translation symmetry and a time reflection symmetry. For Taub-NUT the fields being time translation invariant fail to be time reflection invariant: the adjacent orbits of the Killing vector fields twist around each other! Both solutions are asymptotically flat, but only the Schwarzschild solution is asymptotically Minkowski. The space of classical solutions, which consists of the Schwarzschild solution only, is the static truncation $GL(1)$ of the original sigma model $SL(2, \mathbb{R})/SO(2)$.

A suitable choice of coordinates leads to

$$\omega = \frac{l}{\rho} + O\left(\frac{1}{\rho^2}\right) \quad \text{and} \quad \tau = 1 - \frac{2m}{\rho} + O\left(\frac{1}{\rho^2}\right).$$

The coefficients l and m of the asymptotic expansion of ω and τ are called the Taub-NUT charge l and the Schwarzschild mass m . They are the “physical” parameters of the space of classical solutions. Within the sigma model formalism the currents which are in this case also “global charges”, are contained in the matrix

$$J = \frac{f^2}{2N^2} \chi^{-1} \chi' = \chi_0^{-1} \chi_1,$$

where χ_0 and χ_1 are the coefficients of order zero and one of the asymptotic multipole expansion of the $SL(2, \mathbb{R})/SO(2)$ matrix χ . After a suitable reparametrization one of the equations of motion becomes the equation of geodesic motion in the coset space $SL(2, \mathbb{R})/SO(2)$. This yields a geometrical interpretation of the Taub-NUT charge l and the Schwarzschild mass m . The requirement that χ tends to the “vacuum” configuration in the limit $\rho \rightarrow \infty$ fixes the values of ω and τ at the starting point of the geodesic. Apart from this there exists a uniquely determined velocity vector $(\dot{\tau}_\infty, \dot{\omega}_\infty) = (m, l)$ which characterizes the geodesic completely.

The gravitational remnant parametrized by the coordinate f decouples from the sigma model and can be solved separately.

The quantization of the spherically symmetric sector of Einstein's gravity is achieved via the canonical ADM quantization. At the classical level the ADM formalism selects the coordinates for which the Hamiltonian itself becomes a constraint. The four dimensional space-time is split into a three dimensional spacelike hypersurface and a time. A lapse function N and a shift function N_i are introduced to describe how the 3D hypersurface at time $x_0 = \text{constant}$ with a given 3-metric and an extrinsic curvature is related to the hypersurface at time $x_0 + \delta x_0$. The lapse and shift functions enter the Hamiltonian formalism as Lagrange multipliers. In this paper a modified Hamiltonian formalism is to be implemented. The slicing is performed according to ρ , which is a spacelike coordinate. The formalism can be found in [13]. Due to spherical symmetry the lapse function is the only Lagrange multiplier which survives. In other words: it expresses the invariance under ρ -reparametrization and leads to the only primary first class constraint of the theory, the Hamiltonian

$$H = \frac{1}{4}\pi_f^2 - 1 - \frac{\tau^2}{f^2}(\pi_\tau^2 + \pi_\omega^2),$$

where π_ω, π_τ and π_f denote the canonical ρ -momenta. H generates the gauge transformations. Observables are functions on the constraint surface which are gauge invariant, i.e. their Poisson brackets with the first class constraints vanish. By definition they do not evolve in "time" and therefore there is a one to one correspondence between the observables and the initial data. Expansion of the sigma model current

$$\begin{aligned} J &= \begin{pmatrix} -J^0 & J^+ \\ J^- & J^0 \end{pmatrix} \\ &= \begin{pmatrix} -\tau\pi_\tau - \omega\pi_\omega & -\pi_\omega \\ \tau^2\pi_\omega - 2\omega\tau\pi_\tau - \omega^2\pi_\omega & \tau\pi_\tau + \omega\pi_\omega \end{pmatrix} \end{aligned} \quad (2)$$

into a suitable Lie algebra basis yields two observables J^0 and J^+ of the theory, which commute with H :

$$\{H, J^0\} = 0, \quad \{H, J^+\} = 0.$$

Evaluating J^0 and J^+ on the classical solutions gives the Schwarzschild mass m and the Taub-NUT charge l respectively. The third coefficient J^- is not linearly independent. The currents J^0, J^+ and J^- form an $\mathfrak{sl}(2, \mathbb{R})$ algebra which becomes important in the quantization process. The finite transformation related to J^0 is a scale transformation $\omega \mapsto e^\alpha \omega, \tau \mapsto e^\alpha \tau$. To J^+ corresponds a shift in ω .

In the sequel a Dirac quantization of this sector of Einstein's gravity is performed. In the Schrödinger representation the fields are turned into multiplication operators $\Phi \mapsto \hat{\Phi}$ and the momenta become differentiation operators

$\pi_\Phi \mapsto -i\partial_\Phi$, where ∂_Φ is an abbreviation for $\frac{\partial}{\partial_\Phi}$:

$$\begin{aligned}\widehat{J^0} &= -i\tau\partial_\tau - i\omega\partial_\omega \\ \widehat{J^+} &= i\partial_\omega \\ \widehat{J^-} &= i(\tau^2 - \omega^2)\partial_\omega - 2i\omega\tau\partial_\tau \\ \widehat{\text{Tr}J^2} &= -2\tau^2(\partial_\tau^2 + \partial_\omega^2) \\ \widehat{H} &= -\frac{\partial_f^2}{4} + \frac{1}{2f^2}\widehat{\text{Tr}J^2} - 1.\end{aligned}$$

Applied to a wave function ψ the Hamiltonian defines the so called Wheeler-DeWitt equation $\widehat{H}\psi = 0$. The following commutation relations hold:

$$[\widehat{H}, \widehat{J^0}] = 0 \quad [\widehat{H}, \widehat{J^+}] = 0 \quad [\widehat{H}, \widehat{J^-}] = 0,$$

i.e. the current operators are observables. The Casimir operator $\widehat{\text{Tr}J^2}$ commutes with the Hamiltonian and the currents:

$$[\widehat{\text{Tr}J^2}, \widehat{H}] = 0 \quad [\widehat{\text{Tr}J^2}, \widehat{J^0}] = 0 \quad [\widehat{\text{Tr}J^2}, \widehat{J^+}] = 0 \quad [\widehat{\text{Tr}J^2}, \widehat{J^-}] = 0,$$

and $\widehat{J^0}, \widehat{J^+}, \widehat{J^-}$ form an $\mathfrak{sl}(2, \mathbb{R})$ algebra:

$$[\widehat{J^+}, \widehat{J^0}] = -i\widehat{J^+}, \quad [\widehat{J^+}, \widehat{J^-}] = 2i\widehat{J^0}, \quad [\widehat{J^0}, \widehat{J^-}] = -i\widehat{J^-}.$$

The Casimir operator $\widehat{\text{Tr}J^2}$ is an essential part of the Wheeler-DeWitt equation. As it commutes with the current operators, $\widehat{\text{Tr}J^2}$ and any one of the J 's can be “measured simultaneously”, i.e. they can be simultaneously diagonalized. $\widehat{J^0}, \widehat{J^+}$ and $\widehat{J^-}$ do not commute with each other. Therefore even formally a direct “measurement” of the Schwarzschild mass and the Taub-NUT charge is not possible. The interpretation of the current operators as observables of the theory strongly suggests to preserve the $\text{SL}(2, \mathbb{R})$ symmetry at the quantum level. That is, the quantum mechanical Hilbert space is built from the unitary irreducible representations of the group $\text{SL}(2, \mathbb{R})$. Finding a solution of the Wheeler-DeWitt equation becomes therefore basically a group theoretical problem as the equation splits into an f dependent and an f independent part, which is the Casimir operator on the group up to a constant term. The solutions of the eigenvalue equation for $\widehat{\text{Tr}J^2}$ yield a complete set of eigendistributions in the sense that they are a basis of the representation space.

There are two particularly useful possibilities to diagonalize the Casimir operator $\widehat{\text{Tr}J^2}$. On one hand one solves the differential equations for $\widehat{\text{Tr}J^2}$ and the Taub-NUT charge operator $\widehat{J^+}$ simultaneously:

$$-\tau^2(\partial_\tau^2 + \partial_\omega^2)\psi_{\lambda L}(\omega, \tau) = \lambda\psi_{\lambda L}(\omega, \tau), \quad (3a)$$

$$i\partial_\omega\psi_{\lambda L}(\omega, \tau) = L\psi_{\lambda L}(\omega, \tau). \quad (3b)$$

This system of differential equations yields

$$\psi_{\lambda L}(\omega, \tau) = e^{-iL\omega} \sqrt{\tau} [C_1 \text{I}_k(|L|\tau) + C_2 \text{K}_k(|L|\tau)],$$

where $k = \sqrt{1 - 4\lambda}$. I_k and K_k denote the Bessel functions of first and second kinds with imaginary index k . On the other hand one can diagonalize the Schwarzschild mass operator \widehat{J}^0 and the Casimir operator $\widehat{\text{Tr}} J^2$

$$-\tau^2 (\partial_\tau^2 + \partial_\omega^2) \psi_{\lambda M}(\omega, \tau) = \lambda \psi_{\lambda M}(\omega, \tau) \quad (4a)$$

$$-i (\tau \partial_\tau + \omega \partial_\omega) \psi_{\lambda M}(\omega, \tau) = M \psi_{\lambda M}(\omega, \tau), \quad (4b)$$

which yields

$$\psi_{\lambda M}(\omega, v) = \psi_{\lambda M}^1(\omega, v) + \psi_{\lambda M}^2(\omega, v) \quad (5)$$

with

$$\begin{aligned} \psi_{\lambda M}^1(\omega, v) &= C_1 \omega^{iM} \sqrt{v} \left(1 + v^2\right)^{\frac{iM}{2} - \frac{1}{4}} P_{-iM - \frac{1}{2}}^{-\frac{k}{2}} \left(\frac{1}{\sqrt{1 + v^2}} \right), \\ \psi_{\lambda M}^2(\omega, v) &= C_2 \omega^{iM} v^{k + \frac{1}{2}} |v|^{-k} \left(1 + v^2\right)^{\frac{iM}{2} - \frac{1}{4}} P_{-iM - \frac{1}{2}}^{\frac{k}{2}} \left(\frac{1}{\sqrt{1 + v^2}} \right) \end{aligned}$$

and $v = \omega/\tau$ and $\lambda \neq \frac{1}{4} - n^2$, $n \in \mathbb{Z}$. $P_{-iM - \frac{1}{2}}^{\pm \frac{k}{2}}$ are the associated Legendre functions of first kind.

There is a one to one correspondence between the coset $\text{SL}(2, \mathbb{R})/\text{SO}(2)$ and the Poincaré upper half plane $\mathbb{H} := \{\omega + i\tau \in \mathbb{C}, \tau > 0\}$. Taking a function of the dense subspace $C_c^\infty(\mathbb{H})$ of the representation space $L^2(\mathbb{H}, d\mu)$, where the group invariant measure is $d\mu = d\omega d\tau/\tau^2$, the current operators \widehat{J}^+ and \widehat{J}^0 are shown to be symmetric. The current operators \widehat{J}^+ and \widehat{J}^0 turn out to be already essentially self-adjoint. No constraints on the spectra of \widehat{J}^+ and \widehat{J}^0 arise: it consists of the whole real line for each operator. Negative Taub-NUT charges as well as negative masses belong to the spectrum, too. Suitable solutions are those which belong to the continuous spectrum of the operators. The Bessel functions of first kind with imaginary argument grow exponentially at infinity and do not contribute to the spectral decomposition of the Laplacian. Concerning the second system of differential equations, $\psi_{\lambda M}^2$ drops out because of discontinuities on the imaginary axis.

The “time” dependent part of the Wheeler-DeWitt equation is an ordinary differential equation in f :

$$\psi'' + \left(4 - \frac{2\lambda}{f^2}\right) \psi = 0. \quad (7)$$

The prime $'$ denotes differentiation with respect to f . The dynamical state of the system is entirely hidden in the entities described by ω and τ . In contrast to the coordinate ρ which was arbitrarily chosen, the coordinate f has a geometrical meaning, as $1/f$ is the curvature of the 2-spheres, and serves as a geometrically defined “time”. Therefore (7) describes the “time” evolution of the system. Because it is a second order differential equation, one cannot construct a positive semidefinite probability density that is conserved in “time”. But as in the case of the Klein-Gordon equation [8] a conserved density

$$\rho = \psi^* \partial_f \psi - \psi \partial_f \psi^*$$

can be found. One can interpret the f dependent term in the differential equation (7) as a “time dependent” perturbation. It is instructive to transform the unperturbed differential equation into a system of first order equations $\partial_f \Psi = H_0 \Psi$, $\Psi = (\phi_0, \chi_0)^T$. For $2\phi_0 = \partial_f \psi_0 + 2i\psi_0$ and $2\chi_0 = \partial_f \psi_0 - 2i\psi_0$ the operator H_0 is diagonal, and the density ρ_0 reads: $\rho_0 = i(\phi_0^* \phi_0 - \chi_0^* \chi_0)$. It is a difference of two positive semidefinite parts which correspond to two equivalent states. Given λ the solution of (7) approaches the free solution in the limit $f \rightarrow \infty$ leading to a suggestion how to define two independent but equivalent states in this case, too. The differential equation is of Bessel type, and the appropriate linear combinations turn out to be the Bessel functions of third kind $H_\nu^{(1)}$ and $H_\nu^{(2)}$

$$\psi(f) = C_1 \sqrt{f} H_\nu^{(1)}(2f) + C_2 \sqrt{f} H_\nu^{(2)}(2f), \quad \nu = \frac{1}{2} \sqrt{1 + 8\lambda}.$$

Using the Wronskian it can be shown that the density is again a difference of two positive semidefinite parts.

This paper concludes with some remarks. There is a nice relation between the quantization of four dimensional reduced spherically symmetric gravity in its dual representation and a $SL(2, \mathbb{R})$ WZNW model in the point particle version, which is further reduced to a Liouville theory [9]. One ends up with a Hamiltonian system with constraints, which correspond to initial values. Choosing these initial values, the Hamiltonian of the latter system has exactly the form of the remaining part of the Wheeler-DeWitt equation (7), where the initial values take the place of the eigenvalue of the Casimir operator. The Liouville Hamiltonian is interpreted to be the Hamiltonian of a relativistic particle which moves in a potential. Formally, there are three cases to be distinguished, depending on the sign of the constant λ for the initial values. If $\lambda > 0$, the particle is affected by an infinitely high potential barrier, when it travels in the negative f -direction. For $\lambda = 0$ the particle is free, and for $\lambda < 0$ an infinitely deep potential valley attracts the particle towards the negative f -direction. Here $\lambda > 0$ holds, because the Casimir operator can be shown to be the square of the Taub-NUT charge and the Schwarzschild mass operator.

There are also some important differences between these models. In Fülöp’s paper [9] the third coordinate f has an intrinsic group theoretical meaning leading to an interpretation of the potential. $\lambda > 0$ corresponds to the continuous and $\lambda < 0$ to the discrete series of representation theory. Fülöp considers the momenta of the particle to be fundamental. As the physical meaning is contained in the currents, there is no reason for the momenta to become hermitian operators. Moreover it turns out that starting with hermitian momenta the currents are non-hermitian. On the other hand, the conditions that $\widehat{\mathcal{J}}^0, \widehat{\mathcal{J}}^+$ and $\widehat{\mathcal{J}}^-$ are hermitian defines these operators to be Lie derivatives. This also fixes the operator ordering of the Laplacian and therefore of the Hamiltonian, too.

The more technical aspects of this paper are treated in a separate forthcoming publication.

Acknowledgment

The author is indebted to P. Breitenlohner and D. Maison for numerous discussions on the subject.

References

- [1] Bargmann V.: Irreducible Unitary Representations of the Lorentz Group, *Ann. Math.* **48**, 568 (1947).
- [2] Breitenlohner P., Maison D.: Solitons in Kaluza-Klein Theories, unpublished.
- [3] Breitenlohner P., Maison D., Gibbons G.: 4-Dimensional Black Holes from Kaluza-Klein Theories, *Commun. Math. Phys.* **120**, 295 (1988).
- [4] Chandrasekhar S.: *Mathematical Theory of Black Holes*. Oxford (1983).
- [5] Cavaglià M., de Alfaro V., Filippov A. T.: Hamiltonian Formalism of the Schwarzschild Black Hole, *Int. J. Mod. Phys.* **D4**, 661 (1995)
- [6] Damour T.: *Gravitational Radiation*. N. Deruelle and T.Piran eds., North Holland (1986).
- [7] Dobiasch P., Maison D.: Stationary, Spherically Symmetric Solutions of Jordan's Unified Theory of Gravity and Electromagnetism, *Gen. Rel. Grav.* **14**, 231 (1982).
- [8] Feshbach H., Villars F.: Elementary Relativistic Wave Mechanics of Spin 0 and Spin 1/2 Particles, *Rev. Mod. Phys.* **30**, 24 (1958).
- [9] Fülöp T.: Reduced $SL(2, \mathbb{R})$ WZNW Quantum Mechanics, *J. Math. Phys.* **37** (4), 1617 (1996).
- [10] Kaluza T.: Sitzungsber. Preuss. Akad. Wiss. 966 (1921).
- [11] Kastrup H.A., Thiemann T.: Canonical Quantization of Spherically Symmetric Gravity in Ashtekar's Self-Dual Representation, *Nucl. Phys.* **B339**, 211 (1993)
- [12] Kuchar K.: Geometrodynamics of Schwarzschild Black Holes, *Phys. Rev* **D50**, 3961 (1994)
- [13] Martin J.: Hamiltonian Quantization of General Relativity with the change of Signature, *Phys. Rev.* **D49**, 5089 (1994)
- [14] Louis-Martinez D., Gegenberg J., Kunstatter G.: Exact Dirac Quantization of all 2D-Dilaton Gravity Theories, *Phys. Lett.* **B321**, 193 (1994)
- [15] Taub A.H.: Empty Space-Times Admitting a Three Parameter Group of Motions, *Ann. Math.* **53**, 472 (1951)