

Gravitational coupling constant in higher dimensions

Reza Mansouri^{1,2★} and Ali Nayeri^{3†}

¹*Institute for Studies in Theoretical Physics and Mathematics, P.O.Box 5746, 19395 Tehran, Iran.*

²*Department of Physics, Sharif University of Technology, P.O.Box 11365-9161, Tehran, Iran.*

³*Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411007, India*

ABSTRACT

Assuming the equivalence of FRW-cosmological models and their Newtonian counterparts, we propose using the Gauss law in arbitrary dimension a general relation between the Newtonian gravitational constant G and the gravitational coupling constant κ .

Key words: cosmology: theory.

1 INTRODUCTION

It is now a common belief among cosmologists and relativists that although spacetime appears smooth, nearly flat, and four dimensional on large scales, at sufficiently small distances and early times, it is highly curved with all possible topologies and of arbitrary dimensions. The initial idea of Kaluza-Klein has been extensively used in unified theories of fundamental interactions; e.g. Green, Schwarz, & Witten (1987). There the extra dimensions are assumed to be compactified to Planckian size, and therefore do not display themselves in macroscopic processes. The multidimensional cosmologies based on these ideas have been extensively studied in the last years [see e.g. Cho (1990), Kirillov & Melnikov (1995), Wesson (1992)].

Usually in cosmological models based on higher dimensions the problem of the dimensionality of the gravitational coupling constant is not tackled on, being tacitly assumed to be

$$\kappa = 8\pi G, \quad (1)$$

where G is the Newtonian gravitational constant. This is of course a relation being derived in four dimension. Even in some textbooks this has not been differentiated [see e.g. Kolb & Turner (1990)]. The effective gravitational constant in the multidimensional cosmologies is defined through the multiplication of this four-dimensional value with some volume of the internal space which could even be infinite [Rainer & Zhuk (1996)].

There are however cosmological models in higher dimension where this picture do not work and one should be cautious about the value of the coupling constant in dimensions higher than four [see e.g. Chatterjee (1992), Chatterjee & Bhui (1990), Khorrami, *et. al* (1995), Khorrami, Mansouri

& Mohazzab (1996)]. In fact, Chatterjee (1992), Chatterjee & Bhui (1990) calculate a homogeneous model in a higher dimensional gravity using the same 4-dimensional relation as in (1).

We propose to show some deficiencies of this misuse and propose a generalization for the gravitational coupling constant for any arbitrary dimension. Any definition of a coupling constant in higher dimension will influence the time dependence of G and therefore could have direct observational consequences; e.g. Barrow (1978), Degl’Innocenti, *et. al* (1996), Ramero & Melnikov (1995).

In this note we confine ourselves to a mere generalization of the κ , using discrepancies in comparing relativistic and Newtonian cosmologies in higher dimensions. It belongs to the folklore of the theories of gravitation that their weak limit must be the Newtonian theory of Gravitation. Moreover, on account of the relation (1), there is a Newtonian derivation of the FRW cosmologies. We will show that this derivation, using the coupling constant (1) is just valid in $(3+1)$ -dimension, and has to be changed for higher dimensional theories. On account of the Gauss theorem we give a generalization of it, which makes the Newtonian derivation valid in arbitrary dimensions.

2 FRIEDMANN MODELS AND NEWTONIAN COSMOLOGY IN D DIMENSION

Consider the following Hilbert-Einstein action in $D+1$ dimensional space-time with D as the fixed dimension of space:

$$S_g = -\frac{1}{2\kappa} \int (D+1) R \sqrt{-g} d^{D+1}x + \frac{1}{2} \int T \sqrt{-g} d^{D+1}x, \quad (2)$$

where κ is the gravitational constant, again. For simplicity, we consider the $k=0$ FRW model. In an arbitrary fix dimension we obtain the following Friedmann equation, (Khorrami, *et. al* 1995)

★ email: mansouri@netware2.ipm.ac.ir

† email: ali@iucaa.ernet.in

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{2\kappa}{D(D-1)}\rho, \quad (3)$$

which leads to the familiar Friedmann equation in $\mathbf{D}=3$:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{\kappa}{3}\rho = \frac{8\pi G}{3}\rho. \quad (4)$$

Now, it is well known that the Friedmann equation (4) can be derived from a Newtonian point of view. Taking the Newtonian equation of gravitation in \mathbf{D} dimension in the usual form

$$\nabla_d^2 \varphi = 4\pi G\rho, \quad (5)$$

with φ the gravitational constant and ∇_d the \mathbf{D} dimensional ∇ operator, then the corresponding Friedmann equation can easily be obtained to be

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{D(D-2)}\rho. \quad (6)$$

Now, as expected, for $\mathbf{D} = 3$, the familiar relation (4) is obtained, assuming the relation (1). But, how if $\mathbf{D} \neq 3$. Then one will realise that the agreement between (4) and (6) will fail.

3 MODIFICATION

Looking for the roots of the factor 8π in (1) we come across the relation

$$R_{00} = \nabla^2 \varphi. \quad (7)$$

Now, the coefficient in the Poisson equation, i.e. 4π has been obtained, using Gauss law, for three dimensional space. Thus we should first derive the correct coefficient for a \mathbf{D} dimensional space. Applying Gauss's law for a \mathbf{D} dimensional volume, we find the Poisson equation for arbitrary fixed dimension,

$$\nabla^2 \varphi = \frac{2\pi^{D/2}G}{(D/2-1)!}\rho. \quad (8)$$

On the other hand we get for arbitrary D

$$R_{00} = \left(\frac{D-2}{D-1}\right)\kappa\rho. \quad (9)$$

A comparison of (7), (8), and (9) will give us the following modified Einstein gravitational constant,

$$\kappa_D = \frac{2(D-1)\pi^{D/2}G}{(D-2)(D/2-1)!}. \quad (10)$$

The correct form of Friedmann equation in any arbitrary fixed dimension is now derived to be

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{4\pi^{D/2}G}{D(D-2)(D/2-1)!}\rho. \quad (11)$$

As it is easily seen, the above relation is in complete agreement with its Newtonian counterpart in all dimensions.

4 CONCLUSION

The dimensional dependence of the gravitational constant κ may have very different and serious field theoretic and astrophysical consequences hitherto unnoticed. It would be interesting, e.g., to see which changes are to be expected

if the results of this note are combined with Kaluza-Klein paradigm. The consequences on time variation of G within various models is another very interesting cosmological issue which is under investigation.

ACKNOWLEDGEMENT

A.N. wishes to thank Prof. Padmanabhan for many useful discussions. A.N. was financially supported by the Council of Scientific and Industrial Research, India.

REFERENCES

- Barrow, J.D. 1978, *MNRAS*, **184**, 677.
- Chatterjee, S., 1992. *Ann. Phys.*, **218**, 121.
- Chatterjee, S. & Bhui, B., 1990. *Mon. Not. R. astr. Soc.*, **247**, 57.
- Cho, Y.M., 1990. *Phys. Rev. D.*, **41**, 2462.
- Degl'Innocenti, S., Fiorentini, G., Raffelt, G.G., Ricci, B. & Weiss, A., 1996. *Astron. Astrophys.*, **312**, 345.
- Green, M.B., Schwarz, J.H., & Witten, E., 1987. *Superstring Theory*, Cambridge University Press, Cambridge.
- Kirillov, A.A., & Melnikov, V.N., 1995. *Phys. Rev. D.*, **52**, 723.
- Kolb, E.W., & Turner, M.S., 1990. *The early universe*, Addison-Wesley publishing company, New York.
- Khorrami, M., Mansouri, R., Mohazzab, M. & Ejtehadi, 1995. **gr-qc/9507059**.
- Khorrami, M., Mansouri, R. & Mohazzab, M., 1996. to be published in *Helv. Phys. Acta*.
- Rainer, M. & Zhuk, A. 1996. *preprint*, FUB-HEP/96-3 and IPM-96-149.
- Romero, C., & Melnikov, V.N., 1995. *Gravitation and Cosmology*, **1**, 277.
- Wesson, P.S., 1992. *Astrophys. J.*, **394**, 19.