

EDO-EP-7
October 1996

Black Hole Radiation inside Apparent Horizon in Quantum Gravity

AKIO HOSOYA[†]

*Department of Physics, Tokyo Institute of Technology,
Oh-Okayama, Meguro-ku, Tokyo 152, JAPAN*

and

ICHIRO ODA[‡]

Edogawa University, 474 Komaki, Nagareyama City, Chiba 270-01, JAPAN

ABSTRACT

[†] E-mail address: ahosoya@th.phys.titech.ac.jp

[‡] E-mail address: sjk13904@mgw.shijokyo.or.jp

We study a black hole radiation inside the apparent horizon in quantum gravity. First we perform a canonical quantization for spherically symmetric geometry where one of the spatial coordinates is dealt as the time variable since we would like to consider the interior region of a black hole. Next this rather general formalism is applied for a specific model where the ingoing Vaidya metric is used as a simple model of an evaporating black hole. Following Tomimatsu's idea, we will solve analytically the Wheeler-DeWitt equation in the vicinity of the apparent horizon and see that mass-loss rate of a black hole by thermal radiation is equal to the result obtained by Hawking in his semiclassical treatment. The present formalism may have a wide application in quantum gravity inside the horizon of a black hole such as mass inflation and strong cosmic censorship etc.

1. Introduction

The canonical formalism of general relativity was pioneered in the sixties by Dirac [1], and by Arnowitt, Deser and Misner (ADM) [2,3]. Afterward an extension of the canonical formalism to systems with a black hole was undertaken by Hajicek where special attention was paid for the properties of the apparent horizon and the choice of hypersurfaces covering the spacetime between the apparent horizon and the spatial infinity [4,5]. Recently Kuchař has given a detailed analysis of the geometrodynamics of the Kruskal extension of the Schwarzschild black hole [6].

In this article, we would like to study the canonical formalism of a system with a spherically symmetric black hole only in the interior region bounded by the apparent horizon and the singularity. In this region the Killing vector field $\frac{\partial}{\partial t}$ becomes spacelike, while it does timelike in the exterior region. Consequently one must foliate the interior region of a black hole by a family of spacelike hypersurfaces, for example, $r = \text{const.}$ One will see that it is straightforward to extend Hajicek's works [5] performed in the exterior region to the present case.

So far, the region inside the event horizon of a black hole has been received little attention because it is physically of no relevance for external observers outside the horizon. However, this situation has remarkably changed by the recent development of understanding the internal geometry near the Cauchy horizon inside the Reissner-Nordström black hole, what we call, the mass inflation [7,8,9], and the phenomenon of the smearing of black hole singularity in quantum gravity [10,11]. Anyway since both spacetime singularity and the Cauchy horizon exhibit highly pathological behavior in the classical theory of general relativity, and indeed it is expected that quantum effects play a dominant role around them, the studies of physics inside the horizon of a black hole may give us some important clues for constructing a theory of quantum gravity in future. From this viewpoint, in this article we will present the canonical formalism which describes the region inside the horizon of a spherically symmetric black hole. As one of applications of this

canonical formalism, we shall consider the black hole radiation [12]. The analysis follows the idea by Tomimatsu [13] that the Hamiltonian and supermomentum constraints have a simple and tractable form, and consequently it is possible to solve the Wheeler-DeWitt equation near the apparent horizon. The mass-loss rate by the black hole radiation will be shown to be equal to that evaluated by Hawking in the semiclassical approximation. Hawking's discovery of thermal radiation from quantum black holes was a pivotal event in the development of a quantum theory of gravity [14], thus the results obtained in this paper seems to encourage us to investigate various related problems further along the present canonical formalism.

The article is organized as follows. In section 2, we consider a system of fields in a four dimensional spacetime containing gravity, and transform it to the ADM first-order form. In section 3, we apply the canonical formalism constructed in section 2 for a calculation of the mass-loss rate by the black hole radiation. The last section is devoted to conclusion.

2. Canonical formalism

We begin by constructing a canonical formalism of a spherically symmetric system with a black hole. In this paper, since we have a mind to apply the canonical formalism for various physically interesting phenomena occurring inside the horizon of a black hole we will construct a canonical formalism which holds only in the internal region between the apparant horizon and the singularity of a black hole. First of all, following the conventional procedure, one needs to choose arbitrary spherically symmetric spacelike hypersurfaces to form a foliated structure of the spacetime. The important point is that the radial coordinate plays a role of time inside the horizon in the spherically symmetric coordinate system. As a simple choice, let us take a choice of $x^1 = \text{const}$ hypersurfaces to slice the spacetime. Later we will take the simplest choice $x^1 = r$.

The four dimensional action which we consider in this section is of the form

$$S = \int d^4x \sqrt{-^{(4)}g} \left[\frac{1}{16\pi} {}^{(4)}R - \frac{1}{4\pi} {}^{(4)}g^{\mu\nu} (D_\mu \Phi)^\dagger D_\nu \Phi - \frac{1}{16\pi e^2} F_{\mu\nu} F^{\mu\nu} \right], \quad (1)$$

where Φ is a complex scalar field,

$$D_\mu \Phi = \partial_\mu \Phi + iA_\mu \Phi \quad (2)$$

is its covariant derivative, A_μ is the electromagnetic field, $F_{\mu\nu}$ is the corresponding field strength as usual given by

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu, \quad (3)$$

and e is the electric charge of Φ . To clarify the four dimensional meaning we put the suffix (4) in front of the metric tensor and the curvature scalar. We follow the conventions adopted in the MTW textbook [3] and use the natural units $G = \hbar = c = 1$. The Greek letters μ, ν, \dots take 0, 1, 2, and 3, on the other hand, the Latin ones a, b, \dots do 0 and 1. Of course, the inclusion of other matter fields and the cosmological constants in this formalism is straightforward even if we limit ourselves to the action (1).

The most general spherically symmetric assumption for the metric is

$$\begin{aligned} ds^2 &= {}^{(4)}g_{\mu\nu} dx^\mu dx^\nu, \\ &= g_{ab} dx^a dx^b + \phi^2 (d\theta^2 + \sin^2 \theta d\varphi^2), \end{aligned} \quad (4)$$

where the two dimensional metric g_{ab} and the radial function ϕ are the functions of only the two dimensional coordinates x^a . The substitution of (4) into (1) and then integration over the angular coordinates (θ, φ) leads to the following two

dimensional effective action

$$S = \frac{1}{2} \int d^2x \sqrt{-g} \left[1 + g^{ab} \partial_a \phi \partial_b \phi + \frac{1}{2} R \phi^2 \right] - \int d^2x \sqrt{-g} \phi^2 g^{ab} (D_a \Phi)^\dagger D_b \Phi - \frac{1}{4} \int d^2x \sqrt{-g} \phi^2 F_{ab} F^{ab}, \quad (5)$$

where

$$D_a \Phi = \partial_a \Phi + ie A_a \Phi. \quad (6)$$

Next let us rewrite the action (5) into the first-order ADM form. As mentioned before, we shall take the x^1 coordinate as time to cover the internal region of a black hole by spacelike hypersurfaces. The appropriate ADM splitting of (1+1)-dimensional spacetime is given by

$$g_{ab} = \begin{pmatrix} \gamma & \alpha \\ \alpha & \frac{\alpha^2}{\gamma} - \beta^2 \end{pmatrix}, \quad (7)$$

and the normal unit vector n^a which is orthogonal to the hypersurface $x^1 = \text{const}$ reads

$$n^a = \left(\frac{\alpha}{\beta\gamma}, -\frac{1}{\beta} \right). \quad (8)$$

The induced metric on the hypersurfaces, that is, the projection operator over $x^1 = \text{const}$ hypersurfaces, is given by

$$h^{ab} = g^{ab} + n^a n^b. \quad (9)$$

It is easy to check that h^{ab} is indeed the projection operator by substituting (7) and (8) into (9).

The extrinsic curvature K_{ab} , its trace K , and the scalar curvature R are given by [15]

$$K_{ab} = K_{ba} = h_a^c \nabla_c n_b, \quad (10)$$

$$K = g^{ab} K_{ab} = \nabla_a n^a = \frac{1}{\sqrt{-g}} \partial_a (\sqrt{-g} n^a), \quad (11)$$

and

$$R = 2n^a \partial_a K + 2K^2 - 2\nabla_c (n^a \nabla_a n^c). \quad (12)$$

A straightforward calculation yields

$$K = -\frac{\gamma'}{2\beta\gamma} + \frac{\dot{\alpha}}{\beta\gamma} - \frac{\alpha}{2\beta\gamma^2} \dot{\gamma}, \quad (13)$$

and

$$R = 2n^a \partial_a K + 2K^2 - \frac{2}{\beta\sqrt{\gamma}} \partial_0 \left(\frac{\dot{\beta}}{\sqrt{\gamma}} \right), \quad (14)$$

where $\frac{\partial}{\partial x^0} = \partial_0$ and $\frac{\partial}{\partial x^1} = \partial_1$ are also denoted by an overdot and a prime, respectively. As a result, the action (5) can be written as

$$\begin{aligned} S = \int d^2x L = \int d^2x & \left[\frac{1}{2} \beta \sqrt{\gamma} \{ 1 - (n^a \partial_a \phi)^2 + \frac{1}{\gamma} \dot{\phi}^2 - K n^a \partial_a (\phi^2) \right. \\ & \left. + \frac{\dot{\beta}}{\beta\gamma} \partial_0 (\phi^2) \} + \beta \sqrt{\gamma} \phi^2 \{ |n^a D_a \Phi|^2 - \frac{1}{\gamma} |D_0 \Phi|^2 \} + \frac{1}{2} \beta \sqrt{\gamma} \phi^2 E^2 \right] \\ & + \int d^2x \left[\frac{1}{2} \partial_a (\beta \sqrt{\gamma} K n^a \phi^2) - \frac{1}{2} \partial_0 \left(\frac{\dot{\beta}}{\sqrt{\gamma} \phi^2} \right) \right], \end{aligned} \quad (15)$$

where

$$\begin{aligned} E &= \frac{1}{\sqrt{-g}} F_{01}, \\ &= \frac{1}{\beta\sqrt{\gamma}} (\dot{A}_1 - A'_0). \end{aligned} \quad (16)$$

By differentiating the action (15) with respect to the canonical variables $\Phi(\Phi^\dagger)$, ϕ , γ_0 ■

and A_0 we have the corresponding conjugate momenta $p_\Phi(p_{\Phi^\dagger}), p_\phi, p_\gamma$ and p_A

$$p_\Phi = -\sqrt{\gamma}\phi^2(n^a D_a \Phi)^\dagger, \quad (17)$$

$$p_\phi = \sqrt{\gamma}n^a \partial_a \phi + \sqrt{\gamma}K\phi, \quad (18)$$

$$p_\gamma = \frac{1}{4\sqrt{\gamma}}n^a \partial_a (\phi^2), \quad (19)$$

$$p_A = -\phi^2 E. \quad (20)$$

The Hamiltonian H which is defined as

$$H = \int dx^0 (p_\Phi \Phi' + p_{\Phi^\dagger} \Phi'^\dagger + p_\phi \phi' + p_\gamma \gamma' + p_A A'_0 - L) \quad (21)$$

is expressed by a linear combination of constraints as expected

$$H = \int dx^0 (\alpha H_0 + \beta H_1 + A_1 H_2), \quad (22)$$

where

$$H_0 = \frac{1}{\gamma} [p_\Phi D_0 \Phi + p_{\Phi^\dagger} (D_0 \Phi)^\dagger] + \frac{1}{\gamma} p_\phi \dot{\phi} - 2\dot{p}_\gamma - \frac{1}{\gamma} p_\gamma \dot{\gamma}, \quad (23)$$

$$\begin{aligned} H_1 = & \frac{1}{\sqrt{\gamma}\phi^2} p_\Phi p_{\Phi^\dagger} - \frac{\sqrt{\gamma}}{2} - \frac{\dot{\phi}^2}{2\sqrt{\gamma}} + \partial_0 \left(\frac{\partial_0 (\phi^2)}{2\sqrt{\gamma}} \right) \\ & + \frac{\phi^2}{\sqrt{\gamma}} |D_0 \Phi|^2 - \frac{2\sqrt{\gamma}}{\phi} p_\phi p_\gamma + \frac{2\gamma\sqrt{\gamma}}{\phi^2} p_\gamma^2 + \frac{\sqrt{\gamma}}{2\phi^2} p_A^2, \end{aligned} \quad (24)$$

$$H_2 = -ie(p_\Phi \Phi - p_{\Phi^\dagger} \Phi^\dagger) - \dot{p}_A. \quad (25)$$

Note that α, β and A_1 are non-dynamical Lagrange multiplier fields.

The action can be cast into the first-order ADM canonical form by the dual Legendre transformation

$$S = \int dx^1 \left[\int dx^0 (p_\Phi \Phi' + p_{\Phi^\dagger} \Phi'^\dagger + p_\phi \phi' + p_\gamma \gamma' + p_A A'_0) - H \right]. \quad (26)$$

As Regge and Teitelboim pointed out [16], in order to have the correct Hamiltonian which produces the Einstein equations through the Hamilton equations, one has to supplement the surface terms to the Hamiltonian (22). In the present formalism, since we take the variation of all the fields to be zero at both the singularity and the apparent horizon we do not have to add any surface terms to the Hamiltonian.

3. Black hole radiation

We now turn our attention to an application of the canonical formalism constructed in the previous section for understanding the Hawking radiation [12] from the viewpoint in the internal region of a black hole in quantum gravity. A similar analysis was performed in the external region of a black hole by Tomimatsu [13].

To consider the simplest model of the Hawking radiation, let us turn off the electromagnetic field and treat the neutral scalar field by which the constraint H_2 generating the $U(1)$ gauge transformations identically vanishes. Moreover, we shall use the ingoing Vaidya metric to express the black hole radiation. The treatment of the case of the outgoing Vaidya metric can be made in a perfectly similar way. Here it is useful to introduce the advanced time coordinate v defined as [3]

$$v = t + r^*, \quad (27)$$

with the tortoise coordinate

$$r^* = \int^r \frac{dr}{1 - \frac{2M}{r}}, \quad (28)$$

where $M = M(x^a)$ is the mass function. Here let us define the two dimensional

coordinate x^a by

$$x^a = (x^0, x^1) = (v - r, r). \quad (29)$$

Note that we have chosen $x^1 = r$ as mentioned above. And we fix the gauge freedoms corresponding to the two dimensional reparametrization invariances by the gauge conditions

$$\begin{aligned} g_{ab} &= \begin{pmatrix} \gamma & \alpha \\ \alpha & \frac{\alpha^2}{\gamma} - \beta^2 \end{pmatrix}, \\ &= \begin{pmatrix} -(1 - \frac{2M}{r}) & \frac{2M}{r} \\ \frac{2M}{r} & 1 + \frac{2M}{r} \end{pmatrix}. \end{aligned} \quad (30)$$

From these equations the two dimensional line element takes a form of the Vaidya metric

$$\begin{aligned} ds^2 &= g_{ab} dx^a dx^b, \\ &= -(1 - \frac{2M}{r}) dv^2 + 2dvdr. \end{aligned} \quad (31)$$

This is, of course, rewritten in terms of the (t, r) coordinates by using Eq.s (27), (28) into the Schwarzschild form

$$ds^2 = -(1 - \frac{2M}{r}) dt^2 + \frac{1}{(1 - \frac{2M}{r})} dr^2. \quad (32)$$

For a dynamical black hole, it is useful to consider the local definition of horizon, i.e., the apparent horizon, rather than the global one, the event horizon. The apparent horizon is then defined as

$$x^1 = r = 2M(x^0, x^1). \quad (33)$$

Following the idea in the reference [13], let us attempt to solve the Hamiltonian and supermomentum constraints only in the vicinity of the apparent horizon. Near

the apparent horizon, by intuition we would be able to make an approximation

$$\Phi \approx \Phi(v), M \approx M(v), \phi \approx r. \quad (34)$$

From now on we shall use \approx to indicate the equalities which hold approximately near the apparent horizon. Indeed one can prove the above assumptions (34) to be consistent with the field equations as follows.

The field equations stemming from the action (5) are given by

$$-\frac{2}{\phi}\nabla_a\nabla_b\phi + \frac{2}{\phi}g_{ab}\nabla_c\nabla^c\phi + \frac{1}{\phi^2}g_{ab}\partial_c\phi\partial^c\phi - \frac{1}{\phi^2}g_{ab} = 2(\partial_a\Phi\partial_b\Phi - \frac{1}{2}g_{ab}\partial_c\Phi\partial^c\Phi), \quad (35)$$

$$\frac{1}{\sqrt{-g}}\partial_a(\sqrt{-g}g^{ab}\partial_b\phi) - \frac{1}{2}R\phi = -\phi\partial_a\Phi\partial^a\Phi, \quad (36)$$

$$\partial_a(\sqrt{-g}\phi^2g^{ab}\partial_b\Phi) = 0. \quad (37)$$

As before the Vaidya metric is given by

$$\begin{aligned} g_{ab} &= \begin{pmatrix} g_{vv} & g_{vr} \\ g_{rv} & g_{rr} \end{pmatrix}, \\ &= \begin{pmatrix} -(1 - \frac{2M}{r}) & +1 \\ +1 & 0 \end{pmatrix}. \end{aligned} \quad (38)$$

To prove Eq.(34), in the vicinity of the apparent horizon we make an ansatz that the fields have a form

$$M \approx M(v), \phi \approx r, \quad (39)$$

but the scalar field is still given by $\Phi \approx \Phi(v, r)$. Under this ansatz, the field

equations (35), (36), and (37) become

$$\partial_r \Phi \approx 0, \quad (40)$$

$$\partial_v \Phi \approx \frac{\sqrt{\partial_v M}}{r} \approx \frac{\sqrt{\partial_v M}}{2M}, \quad (41)$$

$$\partial_r \partial_v \Phi \approx \partial_v \partial_r \Phi \approx -\frac{\sqrt{\partial_v M}}{r^2} \approx -\frac{\sqrt{\partial_v M}}{4M^2}. \quad (42)$$

Then one can easily find the solution which satisfies at the same time Eqs. (40), (41), and (42)

$$\Phi(v, r) = \left(1 - \frac{2M}{r}\right)^2 \frac{1}{4\sqrt{\partial_v M}} + \int^v dv \frac{\sqrt{\partial_v M}}{2M(v)}. \quad (43)$$

Consequently,

$$\Phi(v, r) \approx \int^v dv \frac{\sqrt{\partial_v M}}{2M(v)}, \quad (44)$$

which means that one can set $\Phi(v, r) \approx \Phi(v)$ like other fields in the vicinity of the apparent horizon. Thus our assumptions (34) are consistent with fields equations.

Near the apparent horizon (33), Eq.(30) yields

$$\alpha \approx +1, \gamma = \frac{1}{\beta^2} \approx 0, \quad (45)$$

and the canonical conjugate momenta (17), (18) and (19) are given approximately

as

$$\begin{aligned}
p_\Phi &\approx -\phi^2 \partial_v \Phi, \\
p_\phi &\approx -\frac{1}{\gamma} \partial_v M, \\
p_\gamma &\approx -\frac{1}{2} \phi \approx -M.
\end{aligned} \tag{46}$$

Moreover, the two constraints are proportional to each other

$$\begin{aligned}
-\gamma H_0 &\approx \sqrt{\gamma} H_1, \\
&\approx \frac{1}{\phi^2} p_\Phi^2 + \gamma p_\phi.
\end{aligned} \tag{47}$$

Here we have one important remark. Just at the apparent horizon γ becomes zero so that the various equalities approximately hold when we restrict our attention to the interior region near but not at the apparent horizon. Under this sort of regularization γ takes a small but finite value.

An imposition of the constraint (47) as an operator equation on the state produces the Wheeler-DeWitt equation

$$i \frac{\partial \Psi}{\partial \phi} = -\frac{1}{\gamma \phi^2} \frac{\partial^2}{\partial \Phi^2} \Psi. \tag{48}$$

This Wheeler-DeWitt equation can be interpreted as the Schrödinger equation with the superspace Hamiltonian $H_S = \frac{1}{\gamma \phi^2} p_\Phi^2$.

Now it is easy to find a special solution of the Wheeler-DeWitt equation (48) by the method of separation of variables. The result is

$$\Psi = (B e^{\sqrt{A}\Phi} + C e^{-\sqrt{A}\Phi}) e^{-i \frac{A}{\gamma \phi}}, \tag{49}$$

where A , B , and C are integration constants. If one defines an expectation value

$\langle \mathcal{O} \rangle$ of an operator \mathcal{O} as

$$\langle \mathcal{O} \rangle = \frac{1}{\int d\Phi |\Psi|^2} \int d\Phi \Psi^* \mathcal{O} \Psi, \quad (50)$$

one can calculate $\langle \partial_v M \rangle$ by using either (46) or (47)

$$\langle \partial_v M \rangle = -\frac{A}{\phi^2}. \quad (51)$$

This equation shows the black hole radiation when one chooses the constant A to be a positive constant k^2 . Then the mass-loss rate becomes

$$\langle \partial_v M \rangle = -\frac{k^2}{4 \langle M \rangle^2}. \quad (52)$$

This result exactly corresponds to that calculated by Hawking in the semiclassical approach [12] and by Tomimatsu in the exterior region of the apparent horizon in quantum gravity [13]. Thus a black hole completely evaporates within a finite time. However, it is worth pointing out the difference between the Hawking formulation and the present one. In the Hawking's semiclassical approach the gravitational field is fixed as a classical background and only the matter field is treated to be quantum-mechanical. By contrast, our present formulation is purely quantum-mechanical.

At this stage, we must pay our attention to the boundary condition for the physical state. Our physical state which satisfies the Wheeler-Dewitt equation

$$\Psi = (B e^{|k|\Phi} + C e^{-|k|\Phi}) e^{-i \frac{k^2}{\gamma \phi}}, \quad (53)$$

certainly does neither satisfy the Dirichlet boundary condition $\Psi \rightarrow 0$ for $|\Phi| \rightarrow \infty$ nor its norm $\int d\Phi |\Psi|^2$ is finite. Of course, these requirements might be too heavy

since we do not have any physical principle to choose the appropriate boundary condition and the present formalism is too restrictive to select the correct definition of the inner product. Nevertheless, it seems interesting to impose these requirements on our physical state (53). Although we have derived the expression (53) which is valid only in the vicinity of the apparent horizon, we would expect that the qualitative features remain valid in the whole interior region of a black hole except terms coming from some higher-order quantum corrections. Near the spacetime singularity the matter field Φ would strongly fluctuate owing to huge quantum effects, thus the Dirichlet boundary condition seems to be most suitable to suppress this unwieldy behavior of the state. It seems that one of candidates satisfying the Dirichlet boundary condition would be of the form

$$\Psi = (Be^{|k|\Phi} + Ce^{-|k|\Phi})e^{-K\Phi^2}e^{-i\frac{k^2}{\gamma\Phi}}, \quad (54)$$

where K is a certain positive constant. It is interesting that this state (54) also leads to the finite norm under the naively defined measure. It would be wonderful if we can derive such a welcoming physical state by improving the present model in the future.

4. Conclusion

In this article, we have constructed a canonical formalism which would be suitable for a description of a black hole inside the apparent horizon. The aim in mind to use this formalism is to give a basis for analysing various problems associated with an internal structure of a black hole in quantum gravity. As a concrete application based on this canonical formalism, we have studied the Hawking radiation from quantum mechanical viewpoint and shown that the physical state satisfying the Wheeler-DeWitt equation gives us the same mass-loss rate as Hawking's semi-classical calculation. Thus the canonical formalism constructed in this paper seems to give us to a framework in understanding the quantum mechanical properties of a black hole.

In future works, we would like to investigate further other recently developed problems such as the mass inflation, the smearing of the black hole singularities, and the strong cosmic censorship by the present formalism. These studies are currently under active investigation, hence the results will appear in the future publications [17].

References

- [1] P.A.M.Dirac, Lectures on Quantum Mechanics (Yeshiva University, 1964).
- [2] P.Arnowitt, S.Deser, and C.W.Misner, in Gravitation: An Introduction to Current Research, edited by L.Witten (Wiley, New York, 1962).
- [3] C.W.Misner, K.S.Thorne, and J.A.Wheeler, Gravitation (Freeman, 1973).
- [4] P.Thomi, B.Isaak, and P.Hajicek, Phys. Rev. **D30** (1984) 1168.
- [5] P.Hajicek, Phys. Rev. **D30** (1984) 1178.
- [6] K.V.Kuchar, Phys. Rev. **D50** (1994) 3961.
- [7] E.Poisson and W.Israel, Phys. Rev. **D41** (1990) 1796.
- [8] A.Ori, Phys. Rev. Lett. **67** (1991) 789; **68** (1992) 2117.
- [9] P.R.Brady and C.M.Chambers, Phys. Rev. **D51** (1995) 4177; P.R.Brady and J.D.Smith, Phys. Rev. Lett. **75** (1995) 1256.
- [10] A.Hosoya, Class. Quant. Grav. **12** (1995) 2967.
- [11] A.Hosoya and I.Oda, “Black Hole Singularity and Generalized Quantum Affine Parameter”, TIT/HEP-334/COSMO-73, EDO-EP-5, gr-qc/9605069.
- [12] S.W.Hawking, Comm. Math. Phys. **43** (1975) 199.
- [13] A.Tomimatsu, Phys. Lett. **B289** (1992) 283.
- [14] N.D.Birrell and P.C.W.Davies, Quantum Fields in Curved Space (Cambridge University Press, Cambridge, 1982).

- [15] R.M.Wald, General Relativity (The University of Chicago Press, 1984).
- [16] T.Regge and C.Teitelboim, Ann. Phys. (N.Y.) **88** (1974) 286.
- [17] A.Hosoya and I.Oda, “Mass Inflation in Quantum Gravity”, to appear