

Can Temperature Be Quantized?

Li-Xin Li

Department of Physics, Beijing Normal University, Beijing 100875, China

Abstract: It is shown that the temperature may be quantized in some spacetime due to the periodically topological structure of the Euclidean section. The quanta of the temperature is the Hawking-Unruh temperature.

PACS numbers: 04.62.+v, 05.30.-d

In the Euclidean quantum theory physical functions in the Lorentzian section of a complexified spacetime are assumed to be analytically continued from some functions in the Euclidean section [1, 2]. The non-trivial topological structure of the Euclidean section may cause some interesting effects in the Lorentzian section. For example, people have found that in a Lorentzian section whose corresponding Euclidean section is periodic in the Euclidean time $\tau = it$ there is the Hawking-Unruh effect (such as in the black hole spacetime and the de Sitter spacetime). In such a spacetime (Here we call it the Lorentzian Hawking-Unruh type spacetime (L-HU-spacetime), and call the corresponding Euclidean section the Euclidean Hawking-Unruh type spacetime (E-HU-spacetime)), observers whose worldlines are the integral curves of $(\partial/\partial t)^a$, feel that they are in a thermal bath with the temperature $T_0 = 1/\beta_0 = \kappa/2\pi$, where β_0 is the period of the Euclidean time and κ is the surface gravity of the event horizon [3 - 5]. In this letter I show that in a L-HU-spacetime the temperature may be quantized with the quanta T_0 which is the lowest possible temperature for the thermal equilibrium.

It is well known that the Euclidean thermal Green function $G_T(\tau, \vec{x}; \tau', \vec{x}')$ is a periodic (for bosons) or anti-periodic (for fermions) function of τ (and τ') with the period $\beta = 1/T$ where T is the temperature [6, 7]. This holds for systems with vanishing chemical potential in a stationary spacetime or a conformally stationary spacetime [7]. It is easy to show that β is the *fundamental period* of such functions. For example, for the scalar field, $G_T(\tau, \vec{x}; \tau', \vec{x}') = i\text{tr}[e^{-\beta H}\mathcal{T}(\phi(\tau, \vec{x})\phi(\tau', \vec{x}'))]/\text{tr}(e^{-\beta H})$ where \mathcal{T} denotes the Wick time-ordering. $G_T(\tau, \vec{x}; \tau', \vec{x}') = G_T(\tau + \alpha, \vec{x}; \tau', \vec{x}')$ ($0 < \alpha \leq \beta$) leads to $\sum_{mn}(e^{(\alpha-\beta)E_m-\alpha E_n} - e^{-\beta E_n})|\langle m|\phi(\tau, \vec{x})|n\rangle|^2 = 0$ ($H|m\rangle = E_m|m\rangle$ and $\langle m|n\rangle = \delta_{mn}$) in the limit $\tau' \rightarrow \tau$ and $\vec{x}' \rightarrow \vec{x}$, which immediately results in $\alpha = \beta$.

In an E-HU-section with the period β_0 in the Euclidean time τ , every thermal Green function for bosons should be periodic with respect to the translation $\tau \rightarrow \tau + \beta_0$ [4, 7]. While it is also a periodic function of τ with the *fundamental period* β , we must have $\beta_0 = n\beta$ ($n = 1, 2, \dots$) because every doubly-periodic function reduces to a singly-periodic function when the ratio of the two periods is real [8]. Such a conclusion also holds for fermions since an anti-periodic function with the period β is also a period function with the period 2β and every spinor thermal Green function for fermions should be anti-periodic with respect to the translation $\tau \rightarrow \tau + \beta_0$ [7]. Therefore the allowed temperature in the Lorentzian section is

$$T_n = nT_0, \quad n = 1, 2, \dots$$

with $T_0 = 1/\beta_0 = \kappa/2\pi$, which means that in the L-HU-spacetime, the temperature should be quantized with quanta T_0 , and T_0 is the lowest possible temperature for the thermal equilibrium.

References

- [1] Hawking S W 1979 *Recent Developments in Gravitation* ed M Levy and S Deser (New York: Plenum Press)
- [2] Hawking S W 1979 *General Relativity: An Einstein Centenary Survey* ed S W Hawking and W Israel (Cambridge: Cambridge University Press)
- [3] Birrel N D and Davies P C W 1982 *Quantum Fields in Curved Space* (Cambridge: Cambridge University Press)
- [4] Wald R M 1984 *General Relativity* (Chicago: The University of Chicago Press)
- [5] Hawking S and Penrose R 1996 *The Nature of Space and Time* (Princeton: Princeton University Press)
- [6] Fetter A L and Walecka J D 1971 *Quantum Theory of Many-Particle Systems* (San Francisco: McGraw-Hill)
- [7] Gibbons G W and Perry M J 1978 Proc. R. Soc. Lond. A **358** 467
- [8] Whittaker E T and Watson G N 1978 *A Course of Modern Analysis: An Introduction to the General Theory of Infinite Processes and of Analytic Functions; with An Account of the Principal Transcendental Functions* (Cambridge: At the University Press)