

Submitted to Comm. Math. Phys. — January 15, 1997

REISSNER-NORDSTRÖM-LIKE SOLUTIONS OF THE $SU(2)$ EINSTEIN-YANG/MILLS EQUATIONS

by J. A. Smoller and A.G. Wasserman

In this paper we study a new type of solution of the spherically symmetric Einstein-Yang/Mills (EYM) equations with $SU(2)$ gauge group. These solutions are well-behaved in the far-field, and have a Reissner-Nordström type essential singularity at the origin $r = 0$. These solutions display some novel features which are not present in particle-like, or black-hole solutions.

In order to describe these solutions and their properties, we recall that for the spherically symmetric EYM equations, the Einstein metric is of the form

$$ds^2 = -AC^2 dt^2 + A^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \quad (1.1)$$

and the $SU(2)$ Yang-Mills curvature 2-form is

$$F = w'\tau_1 dr \wedge d\theta + w'\tau_2 dr \wedge (\sin d\phi) - (1 - w^2)\tau_3 d\theta \wedge (\sin \theta d\phi) \quad (1.2).$$

Here A, C , and w are functions of r , and τ_1, τ_2, τ_3 form a basis for the Lie algebra $su(2)$. These equations have been studied in many papers; see e.g. [3-19].

Smooth solutions of the EYM equations, defined for all $r \geq 0$, are called (Bartnik-McKinnon, BM) particle-like solutions; such solutions satisfy $1 > A(r) > 0$ for all $r > 0$, and $A(0) = 1$. The EYM equations also admit black-hole solutions; i.e. solutions defined for all $r \geq \rho > 0$, where $A(r) = 0$. Here again, $1 > A(r) > 0$ for all $r > \rho$. The classical

Reissner-Nordström (RN) solutions of the Einstein equations with zero electric charge, $A(r) = 1 - \frac{c}{r} + \frac{1}{r}$, ($c = \text{const.}$), $(AC^2)(r) = (1 - \frac{c}{r} + \frac{1}{r^2})$, are also solutions to the EYM equations, with $w(r) \equiv 0$. We note that for this solution, $A(r) > 1$ for r near 0. We prove the existence of solutions which have this feature, ($A(r) > 1$ for r near 0), of the classical RN solution, and we study their properties; we call these Reissner-Nordström-Like (RNL) solutions.¹

For these RNL solutions, we show that if $A(r_1) = 1$ for some $r_1 > 0$, then the solution is defined for all r , $0 < r \leq r_1$, and $\lim_{r \searrow 0} A(r) = \infty = \lim_{r \searrow 0} (AC^2)(r)$. However, the function $B(r) \equiv r^2 A(r)$ is analytic, on $0 \leq r \leq r_1$, as is the function $w(r)$; moreover $\lim_{r \searrow 0} w'(r) = 0$.

If we consider solutions that are defined in the far field; i.e., for $r \gg 1$, then it was shown in [10] that $\lim_{r \searrow \infty} (A(r), w^2(r), w'(r)) = (1, 1, 0)$. Thus the projection of the solution in the $w - w'$ plane for a particle like solution starts at the “rest point” $(\pm 1, 0)$, and goes to a “rest point” $(\pm 1, 0)$. Black-hole solutions start at certain curves in the $w - w'$ plane, ([9]), and end at a rest point $(\pm 1, 0)$. In both of these cases, there are an infinite number of solutions, distinguished by their nodal class ([8,9]). For RNL solutions, there is a parameter $\sigma > 0$ defined by $A(\sigma) = 1$. We prove that for fixed σ , there are an infinite number of RNL solutions distinguished by their integral nodal class, which must start at $r = 0$ on the line $w' = 0$, and end at a rest point $(\pm 1, 0)$. The RNL solutions corresponding to the special case $w(0) = 0$ are tangent to the line $w = 0$; these give rise to half-integral nodal classes.

¹ We base the name RNL on the behavior of such solutions near $r = 0$. For such solutions which are connecting orbits and have finite (ADM) mass, the results in [2, p. 393] show that $A(r) = 1 - \frac{c}{r} + \frac{0(1)}{r^2}$, as $r \rightarrow \infty$ and thus behave differently at $r = \infty$ from the RN solutions. (The RN solutions that we consider have zero electric charge and unit magnetic charge; c.f. [18]. We thank P. Bizon for pointing this out to us.)

The proof of the existence of locally defined RNL solutions relies on a local existence theorem at $r = 0$, where we show that there is a 3-parameter family of analytic solutions starting at $r = 0, w'(0) = 0$. The proof of the existence of these local analytic solutions is non-trivial because the associated vector field is not even continuous at $r = 0$; (see the last part of §3). Some of these solution have been found numerically in [20]. It is interesting to note that when the first parameter $w(0) = \pm 1$, and the second parameter $b_1 = 0$, we recover the BM solutions. If $b_1 > 0$, we get RNL solutions and if $b_1 < 0$ we get Schwarzschild-like solutions.

We also prove that for fixed $\sigma > \frac{1}{2}$, the (ADM) masses of a sequence of our RNL solutions for σ fixed, and increasing nodal class, tends to $\frac{1}{\sigma}$. Furthermore, the globally defined RNL solutions which we obtain all have naked singularities at $r = 0$; there may well be other RNL solutions for which the singularity at $r = 0$ is inside an event horizon. We prove that the singularity at $r = 0$ for these RNL solutions is always non-removable.

We show how the results which we have obtained, enable us to classify all spherically symmetric EYM solutions, with $SU(2)$ gauge group, which are smooth and satisfy $A > 0$ in the far field.

Please direct requests for hard copies of complete paper to awass@math.lsa.umich.edu or smoller@math.lsa.umich.edu

REFERENCES

1. Adler, R., Bazin, M. Schiffer, M., Introduction to General Relativity, 2nd ed., McGraw Hill, New York, (1975).
2. Arnowitt, R., Deser, S., Misner, C.W., The dynamics of general relativity, in: Gravitation, ed. by L. Witten, Wiley, N.Y. 227-265, (1962).
3. Bartnik, R., and McKinnon, J., Particle-like solutions of the Einstein-Yang-Mills equations, Phys. Rev. Lett., 61, 141-144, (1988).
4. Bizon, P., Colored black holes, Phys. Rev. Lett. 64, 2844-2847, (1990).

5. Breitenlohner, P., Forgacs, P., and Maison, D., Static spherically symmetric solutions of the Einstein-Yang-Mills equations, *Comm. Math. Phys.*, 163, 141-172, (1994).
6. Kunzle, H.P., and Masood-ul-Alam, A.K.M., Spherically symmetric static $SU(2)$ Einstein-Yang-Mills fields, *J. Math. Phys.* 31, 928-935 (1990).
7. Smoller, J., Wasserman, A., Yau, S.-T., McLeod, J., Smooth static solutions of the Einstein/Yang-Mills equations, *Comm. Math. Phys.*, 143, 115-147 (1991).
8. Smoller, J., and Wasserman, A., Existence of infinitely-many smooth static, global solutions of the Einstein/Yang-Mills equations, *Comm. Math. Phys.*, 151, 303-325, (1993).
9. Smoller, J., Wasserman, A., and Yau, S.-T., Existence of black hole solutions for the Einstein-Yang/Mills equations, *Comm. Math. Phys.*, 154, 377-401, (1993).
10. Smoller, J., and Wasserman, A., Regular solutions of the Einstein-Yang/ Mills equations, *J. Math. Phys.*, 36, 4301-4323, (1995).
11. Smoller, J. and Wasserman, A., Limiting masses of solutions of Einstein- Yang/Mills equations, *Physica D*, 93, 123-136, (1996).
12. Smoller, J. and Wasserman, A., Uniqueness of extreme Reissner-Nordstr?m solution in $SU(2)$ Einstein-Yang-Mills theory for spherically symmetric spacetime, *Phys. Rev. D*, 15 Nov. 1995, 52, 5812-5815, (1995).
13. Straumann, N., and Zhou, Z., Instability of a colored black hole solution, *Phys. Lett. B.* 243, 33-35, (1990).
14. Smoller, J., and Wasserman, A., Uniqueness of zero surface gravity $SU(2)$ Einstein-Yang/Mills black-holes, *J. Math. Phys.* 37, 1461-1484, (1996).
15. Smoller, J., and Wasserman, A., An investigation of the limiting behavior of particle-like solutions to the Einstein-Yang/Mills equations and a new black hole solution, *Comm. Math. Phys.*, 161, 365-389, (1994).

16. Ershov, A.A., and Galtsov, D.V., Non abelian baldness of colored black holes, Phys. Lett. A, 150, 747, (1989), 160-164.
17. Lavrelashvili, G., and Maison, D., Regular and black-hole solutions of Einstein/Yang-Mills dilaton theory, Phys. Lett. B, 295, (1992), 67.
18. Bizon, P. and Popp, O., No-hair theorem for spherical monopoles and dyons in SU(2) Einstein-Yang-Mills theory, Class. Quantum Gravity, 9, (1992), 193.
19. Volkov, M.S., and Gal'tsov, D.V., Sphalerons in Einstein/Yang-Mills theory. Phys. Lett. B, 273, (1991), 273.
20. Donets, E.E., Gal'tsov, D.V., and Zotov, M.Yu., Internal structure of Einstein-Yang-Mills black holes, gr-qc/9612067.