

DARBOUX PARAMETER FOR EMPTY FRW QUANTUM UNIVERSES AND QUANTUM COSMOLOGICAL SINGULARITIES

HARET C. ROSU

Instituto de Física, Universidad de Guanajuato, Apdo Postal E-143, León, Gto, Mexico

Received (May 27, 1997)

Revised (January 17, 1998)

I present the factorization(s) of the Wheeler-DeWitt equation for vacuum FRW minisuperspace universes of arbitrary Hartle-Hawking factor ordering, including the so-called strictly isospectral supersymmetric method. By the latter means, one can introduce an infinite class of singular FRW minisuperspace wavefunctions characterized by a Darboux parameter that mathematically speaking is a Riccati integration constant, while physically determines the position of these strictly isospectral singularities on the Misner time axis.

In cosmology one can encounter arbitrary parameters that are difficult to determine by experiment and theory. One famous example is Einstein's cosmological constant, which does set the time scale in the vacuum dominated universe.¹ Another, very recent example is Immirzi's parameter in quantum general relativity.² In this work, using the factorization(s) of the Wheeler-DeWitt (WDW) minisuperspace equation for the vacuum Friedmann-Robertson-Walker (FRW) universes, an infinite class of singular FRW wavefunctions are introduced containing an arbitrary parameter of Darboux type³ determining the position of the singularities during their Misner time evolution.

Recalling that factorizations of second order linear one-dimensional differential operators are common tools in Witten's supersymmetric quantum mechanics⁴ and imply particular solutions of Riccati equations known as superpotentials, I explore here the physical consequences of the most general factoring of the FRW WDW equation for arbitrary Hartle-Hawking⁵ factor ordering Q by means of the general solution of Riccati equation, a procedure which has been first used in physics by Mielnik for the quantum harmonic oscillator,⁶ and which is known as the strictly isospectral supersymmetric method and/or the double Darboux method.⁷ Stated differently, I shall exploit the non-uniqueness of the factorization of second-order linear differential operators on the simple example of the FRW WDW differential equation with Hartle-Hawking factor ordering, that I write down in the form

$$D^2\Psi \equiv \left(\frac{d^2}{d\Omega^2} + 2\frac{Q}{2}\frac{d}{d\Omega} - e^{-4\Omega} \right) \Psi = 0 . \quad (1)$$

The independent variable Ω is Misner's time parameter related to the volume of space V at a given cosmological epoch through $\Omega = -\ln(V^{1/3})$.⁸ Using the change

of function $\Psi = \Phi \exp(-\frac{Q}{2}\Omega)$, one gets the following Schrödinger equation of zero eigenvalue on the Misner axis

$$\Phi'' - \left(\frac{Q^2}{4} + e^{-4\Omega}\right) \Phi = 0 . \quad (2)$$

The general solution is the modified Bessel function $Z_{Q/4}(i\frac{1}{2}e^{-2\Omega}) = C_1 I_{Q/4}(\frac{1}{2}e^{-2\Omega}) + C_2 K_{Q/4}(\frac{1}{2}e^{-2\Omega})$ (the Cs are superposition constants). Usually, the strictly isospectral supersymmetric technique requires nodeless, normalizable Schrödinger solutions. Although the modified Bessel functions $Z_{Q/4}$ are not normalizable, one can still use the strictly isospectral technique as first shown by Pappademos *et al* for a few simple scattering examples in ordinary quantum mechanics.⁹ Using the modified Bessel function $Z_{Q/4}$, the strictly isospectral supersymmetric construction says that the class of strictly isospectral potentials for the FRW cosmological models are given by

$$\begin{aligned} S_{iso}(\Omega; \Omega_D) &= S(\Omega) - 2[\ln(\mathcal{J}_Z(\Omega) + \Omega_D)]'' \\ &= S(\Omega) - \frac{4Z_{Q/4}Z'_{Q/4}}{\mathcal{J}_Z + \Omega_D} + \frac{2Z_{Q/4}^4}{(\mathcal{J}_Z + \Omega_D)^2} , \end{aligned} \quad (3)$$

where $S(\Omega) = \frac{Q^2}{4} + e^{-4\Omega}$, Ω_D is the Darboux parameter introduced by this method and

$$\mathcal{J}_Z(\Omega) \equiv \int_{\infty}^{\Omega} Z_{Q/4}^2(i e^{-2y}/2) dy . \quad (4)$$

To get (3) one factorizes the one-dimensional FRW Schrödinger equation with the operators $A = \frac{d}{d\Omega} + \mathcal{W}(\Omega)$ and $A^\dagger = -\frac{d}{d\Omega} + \mathcal{W}(\Omega)$, where the superpotential function is given by $\mathcal{W} = -Z'_{Q/4}/Z_{Q/4}$. The Schrödinger potential and Witten's superpotential enter an initial 'bosonic' Riccati equation $S = \mathcal{W}^2 - \mathcal{W}'$. On the other hand, one can build a 'fermionic' Riccati equation $S^+ = \mathcal{W}^2 + \mathcal{W}'$, corresponding to a 'fermionic' Schrödinger equation for which the operators A and A^+ are applied in reversed order. Thus, the 'fermionic' FRW Schrödinger potential is found to be $S^+ = \frac{Q^2}{4} + e^{-4\Omega} - 2(\frac{Z'_{Q/4}}{Z_{Q/4}})'$. This potential does not have $Z_{Q/4}$ as an eigenfunction. However, it is possible to reintroduce the $Z_{Q/4}$ solution into the spectrum, by using the general superpotential solution of the fermionic Riccati equation. The general Riccati solution reads

$$\mathcal{W}_{gen} = \mathcal{W}(\Omega) + \frac{d}{d\Omega} \ln[\mathcal{J}_Z(\Omega) + \Omega_D] , \quad (5)$$

where $\Omega_D \in (-\infty, +\infty)$ occurs as an integration constant.¹⁰ The way to obtain (5) is well known since the work of Mielnik and will not be repeated here. From it one can easily get (3). The Darboux parameter family of FRW wavefunctions differs from the initial one, being the Ω_D -dependent quotient¹¹

$$Z_{gen}(\Omega; \Omega_D) = \frac{Z_{Q/4}(i\frac{1}{2}e^{-2\Omega})}{\mathcal{J}_Z + \Omega_D} . \quad (6)$$

The last equation has interesting implications for the celebrated issue of cosmological singularities. Indeed, the strictly isospectral supersymmetric method may be considered as a way of generating parametric families of new cosmological wavefunctions with the important feature that there is always a point Ω_s on the Misner axis defined by $\Omega_D = -\mathcal{J}_Z(\Omega_s)$ at which they become singular (as well as the strictly isospectral potentials). In other words, an infinite set of such strictly isospectral supersymmetric singularities occur, whereas in the classical case one is usually bound only to the idea of the original singularity, which is by definition at $\Omega \rightarrow \infty$, “an infinite Ω time in the past” as Misner put it.⁸ Therefore, we agree with Wheeler’s dictum¹² “No, quantum mechanics does not provide an escape from gravitational collapse”. Even more, the solutions with singularities are never a set of measure zero for the quantum FRW universes. There exist well-defined mathematical procedures to produce an infinite amount of solutions with singularities, at least for the FRW case. Finally, for some plots displaying the strictly isospectral supersymmetric FRW singularities in the particular case $Q = 0$ the reader is directed to one of my works with Socorro.¹³

Acknowledgements

This work was partially supported by the CONACyT Project 4868-E9406. The author thanks the referee for suggestions and Dr. B.K. Berger for a useful electronic message.

References

1. For a review see, S. Weinberg, *Rev. Mod. Phys.* **61** (1989) 1.
2. G. Immirzi, *Nucl. Phys. Proc. Suppl.* **57** (1997) 65; C. Rovelli and T. Thiemann, *Phys. Rev.* **D57** (1998) 1009.
3. G. Darboux, *C.R. Acad. Sci.* **94** (1882) 1456.
4. E. Witten, *Nucl. Phys.* **B185** (1981) 513. For a recent review, see F. Cooper, A. Khare, and U. Sukhatme, *Phys. Rep.* **251** (1995) 267.
5. J. Hartle and S.W. Hawking, *Phys. Rev.* **D28** (1983) 2960.
6. B. Mielnik, *J. Math. Phys.* **25** (1984) 3387.
7. See, e.g., H.C. Rosu, *Phys. Rev.* **A54** (1996) 2571; *Phys. Rev.* **E56** (1997) 2269.
8. C.W. Misner, *Phys. Rev. Lett.* **22** (1969) 1071; *Phys. Rev.* **186** (1969) 1319, 1328.
9. J. Pappademos, U. Sukhatme, and A. Pagnamenta, *Phys. Rev.* **A48** (1993) 3525.
10. See, L.J. Boya *et al*, quant-ph/9711059 (Nuovo Cimento **B113** (1998) xxx).
11. See, e.g., H.C. Rosu and J. Socorro, *Phys. Lett.* **A223** (1996) 28; J. Socorro and H.C. Rosu (gr-qc/9612032), in *Proc. of the Second Mexican School of Gravitation and Mathematical Physics*, ed. A. Garcia *et al* (Science Network Publishing, Konstanz, 1997).
12. J.A. Wheeler, *Gen. Rel. Grav.* **8** (1977) 713.
13. H.C. Rosu and J. Socorro, gr-qc/9606030 (to be published).