

Perturbations in the Kerr-Newman Dilatonic Black Hole Background I. Maxwell Waves

R. Casadio, B. Harms and Y. Leblanc

*Department of Physics and Astronomy, The University of Alabama
Box 870324, Tuscaloosa, AL 35487-0324*

P.H. Cox

*Physics Department, Texas A&M University-Kingsville
Kingsville, TX 78363*

Abstract

In this paper we analyze the perturbations of the Kerr-Newman dilatonic black hole background. For this purpose we perform a double expansion in both the background electric charge and the wave parameters of the relevant quantities in the Newman-Penrose formalism. We then display the gravitational, dilatonic and electromagnetic equations, which reproduce the static solution (at zero order in the wave parameter) and the corresponding wave equations in the Kerr background (at first order in the wave parameter and zero order in the electric charge). At higher orders in the electric charge one encounters corrections to the propagation of waves induced by the presence of a non-vanishing dilaton. An explicit computation is carried out for the electromagnetic waves up to the asymptotic form of the Maxwell field perturbations produced by the interaction with dilatonic waves. A simple physical model is proposed which could make these perturbations relevant in the detection of radiation coming from the region of space near a black hole.

4.60.+n, 11.17.+y, 97.60.Lf

I. INTRODUCTION

The theory of extended objects such as strings and membranes is a very attractive candidate for the quantum theory of gravity. The drawback of the theory of extended objects is that the theory is anomaly free only in some space whose dimension is much larger than one in which we can make measurements. This drawback is partially mitigated by the fact that a four-dimensional, low energy effective action can be written in terms of the fields associated with the massless excitations of extended objects. In a series of papers [1]- [8] we have explored the viewpoint that quantum black holes are massive excitations of extended objects and hence are elementary particles. In Ref. [9], starting from the low energy effective action describing the Einstein-Maxwell theory interacting with a dilaton in four dimensions ($G = 1$),

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} \left[R - \frac{1}{2} (\nabla\phi)^2 - e^{-a\phi} F^2 \right] , \quad (1.1)$$

we obtained the static solutions of the field equations for a Kerr-Newman dilaton black hole rotating with arbitrary angular momentum by expanding the fields in terms of the charge-to-mass ratio of the black hole. In the present paper we use these solutions, which appear as source terms in the wave equations, to determine the effect of a scalar component of gravity (the dilaton) on the propagation of electromagnetic waves in the vicinity of a charged, rotating black hole.

Our goal is to obtain expressions which can be compared to experimentally measurable quantities in order to test the idea that quantum black holes are extended objects. In all likelihood the scalar component of gravity will have a coupling to electromagnetic waves which is as weak as that of the tensor component. Therefore the best place to look for the effect of the dilaton on these waves is in the neighborhood of a cosmological black hole, which we consider to be composed of quantum black holes. The expressions we obtain in this paper show how the electromagnetic radiation reaching a distant observer from such black holes is affected by the presence of a scalar component of gravity.

We begin in Section II with a summary of the static background solutions for a Kerr-Newman dilatonic black hole. Perturbative expressions for the metric tensor components and the electromagnetic fields are given through order Q^2 . Expressions for the gravitational field, the dilaton field and the electromagnetic field in terms of the Newman-Penrose formalism complete this section.

In Section III we obtain the wave equations for the various field modes. Our method is to double expand each field in powers of the electric charge and the wave parameters. Substituting these expansions into Maxwell's equations, the dilaton equation and Einstein's equations then give (inhomogeneous) wave equations for the coefficients of the expansion to any desired accuracy. Part of the currents in the wave equations arise due to the presence of the dilaton. We give explicit expressions for the Maxwell wave equations for the coefficients linear in the wave parameter for the three lowest orders of the charge parameter.

The forms of the solutions of the wave equations at large r are examined in Section IV. We derive explicit expressions for the corrections to the Maxwell scalars for both the ingoing and outgoing waves. An infrared problem which arises in the amplitudes of these corrections

is also discussed. In the same Section we analyze the corrections to the ingoing and outgoing waves near the horizon.

In Section V we discuss the production of outgoing electromagnetic waves by the dilatonic waves which occur in the case of dynamic black holes. We give the explicit contributions to the electric and magnetic fields arising from the solutions we found in the previous Section and also use these contributions to write the expression for the correction to the energy flux due to the presence of a dilaton field.

Section VI is a discussion of our results and of the further investigations these results allow us to pursue.

II. THE STATIC BACKGROUND SOLUTION

The action describing the Einstein-Maxwell theory interacting with a dilaton in four dimensions is given in Eq. (1.1), where R is the scalar curvature, ϕ is the dilaton field, a is the dilaton parameter and F is the Maxwell field. In tensor components, $i, j = 0, \dots, 3$, the corresponding field equations are

$$\begin{cases} \nabla^i (e^{-a\phi} F_{ij}) = 0 \\ \nabla_{[k} F_{ij]} = 0, \end{cases} \quad (\text{Maxwell}) \quad (2.1)$$

$$\nabla^2 \phi = -a e^{-a\phi} F^2 \quad (\text{dilaton}), \quad (2.2)$$

$$R_{ij} = \frac{1}{2} \nabla_i \phi \nabla_j \phi + 2 T_{ij}^{EM} \quad (\text{Einstein}), \quad (2.3)$$

where ∇ is the covariant derivative, R_{ij} is the Ricci tensor, and the electromagnetic energy-momentum tensor is given as

$$T_{ij}^{EM} = e^{-a\phi} \left[F_{ik} F_j^k - \frac{1}{4} g_{ij} F^2 \right]. \quad (2.4)$$

In Ref. [9] we found a perturbative static solution of the field equations above representing a charged, rotating black hole of the Kerr-Newman type with a non-vanishing background dilaton field in which the parameter of expansion is the charge-to-mass ratio, Q/M [10].

A. Perturbative Q/M solution

The metric in our solution is of the general form which describes a stationary axisymmetric space-time [11]

$$ds^2 = -\frac{\Delta \sin^2 \theta}{\Psi} (dt)^2 + \Psi (d\varphi - \omega dt)^2 + \rho^2 \left[\frac{(dr)^2}{\Delta} + (d\theta)^2 \right], \quad (2.5)$$

in which $x^0 = t$, $x^1 = r$, $x^2 = \theta$, $x^3 = \varphi$.

The explicit expressions of the functions $\Psi = \Psi(r, \theta)$, $\omega = \omega(r, \theta)$ and $\Delta = \Delta(r)$ as displayed in Ref. [9] can be simplified upon substituting for the (bare) parameters M , Q and $J \equiv \alpha M$ the quantities

$$\begin{aligned}
M_{phys} &\equiv M \left[1 + \frac{a^2 Q^2}{6 M^2} \right] \\
Q_{phys} &\equiv Q \left[1 + \frac{a^2 Q^2}{3 M^2} \right] \\
J_{phys} &\equiv \alpha M_{phys} ,
\end{aligned} \tag{2.6}$$

which are respectively the ADM mass, charge and angular momentum of the hole which determine the geodesic motion in the asymptotically flat region. We also introduce the following shift in the radial coordinate

$$r \rightarrow r - \frac{a^2 Q_{phys}^2}{6 M_{phys}} , \tag{2.7}$$

and we finally get that the metric in Eq. (2.5) coincides (at order Q_{phys}^2/M_{phys}^2) with the pure Kerr-Newman solution, that is

$$\begin{aligned}
\Delta &= r^2 - 2 M_{phys} r + \alpha_{phys}^2 + Q_{phys}^2 \\
\rho^2 &= r^2 + \alpha_{phys}^2 \cos^2 \theta \\
\Psi &= -\frac{\Delta - \alpha_{phys}^2 \sin^2 \theta}{\rho^2} \\
\omega &= -\alpha_{phys} \sin^2 \theta [1 + \Psi^{-1}] ,
\end{aligned} \tag{2.8}$$

where we have also written $\alpha_{phys} = \alpha$. This also implies that the singularity structure is not affected by the presence of the dilaton at that order. In the general case however (if one includes the higher order corrections in Q^2/M^2) this is not so, because the spacetime is very likely no longer a Petrov type-D spacetime. In the present case one still has two horizons for $\Delta = 0$, that is

$$r_{\pm} = M_{phys} \pm \sqrt{M_{phys}^2 - \alpha_{phys}^2 - Q_{phys}^2} . \tag{2.9}$$

However the presence of a non-zero dilaton field

$$\phi = -a \frac{r}{\rho^2} \frac{Q_{phys}^2}{M_{phys}} , \tag{2.10}$$

affects the electric and magnetic field potentials A and B (see Ref. [9] for the definitions),

$$\begin{aligned}
A &= Q_{phys} \frac{r}{\rho^2} \left[1 - \left(\frac{1}{2r} + \frac{r}{\rho^2} \right) \frac{a^2 Q_{phys}^2}{3 M_{phys}} \right] \\
B &= -Q_{phys} \alpha_{phys} \frac{\cos \theta}{\rho^2} \left[1 - \left(\frac{1}{2M} - \frac{r}{\rho^2} \right) \frac{a^2 Q_{phys}^2}{3 M_{phys}} \right] ,
\end{aligned} \tag{2.11}$$

where the terms proportional to Q_{phys}^2 inside the brackets correspond to the corrections with respect to the Kerr-Newman potentials [11]. These corrections also affect the electric and magnetic fields in the limit $r \rightarrow \infty$,

$$\begin{aligned}
\mathcal{E}_{\hat{r}} &\approx \frac{Q_{phys}}{r^2} \\
\mathcal{E}_{\hat{\theta}} &\approx -\frac{2\alpha_{phys}^2 Q_{phys}}{r^4} \sin\theta \cos\theta \\
\mathcal{B}_{\hat{r}} &\approx \frac{2\alpha_{phys} Q_{phys}}{r^3} \cos\theta \left[1 - \frac{a^2 Q_{phys}^2}{6 M_{phys}^2} \right] \\
\mathcal{B}_{\hat{\theta}} &\approx \frac{\alpha_{phys} Q_{phys}}{r^3} \sin\theta \left[1 - \frac{a^2 Q_{phys}^2}{6 M_{phys}^2} \right] ,
\end{aligned} \tag{2.12}$$

where $\mathcal{E}_{\hat{a}}$ and $\mathcal{B}_{\hat{a}}$ ($\hat{a} = \hat{r}, \hat{\theta}$) are respectively the electric and the magnetic field components with respect to the usual spherical coordinate tetrad basis [12]. One would thus experience a relative shift in the intensity of the field \mathcal{B} with respect to \mathcal{E} . This is also the source for the anomalous gyromagnetic ratio

$$g = 2 \left[1 - \frac{a^2 Q_{phys}^2}{6 M_{phys}^2} \right] , \tag{2.13}$$

which should be compared with $g = 2$ for the Kerr-Newman black hole [13].

For the sake of simplicity, from now on we omit the subscript *phys* in all Q , M and α .

B. Newman-Penrose form of the solution

In the next Section we will compute the linear perturbation field equations for the gravitational field, Eq. (2.3), the scalar dilaton field, Eq. (2.2), and the Maxwell field, Eq. (2.1), in this background. We will work in the Newman-Penrose formalism [14,11], which is a special type of tetrad calculus in which one introduces four null vectors e_a^i , $a = 1, \dots, 4$,

$$\begin{aligned}
l^i &\equiv e_1^i \\
n^i &\equiv e_2^i \\
m^i &\equiv e_3^i \\
m^{*i} &\equiv e_4^i ,
\end{aligned} \tag{2.14}$$

at every point of space-time, where m^{*i} is the complex conjugate of m^i . They must satisfy the normalization conditions

$$l^i n_i = 1 \quad \text{and} \quad m^i m_i^* = -1 . \tag{2.15}$$

Further l^i is affinely parameterized while the other vectors are not.

Since our static metric coincides with the Kerr-Newman one, and the latter differs from the Kerr metric only in the definition of Δ [11], all the quantities formally computed in the Kerr case apply. In detail, one has the following expressions for the four vectors

$$\begin{aligned}
l^i &= \frac{1}{\Delta} (r^2 + \alpha^2, +\Delta, 0, \alpha) \\
n^i &= \frac{1}{2\rho^2} (r^2 + \alpha^2, -\Delta, 0, \alpha) \\
m^i &= \frac{1}{\sqrt{2}\bar{\rho}} (i\alpha \sin\theta, 0, 1, i \csc\theta) ,
\end{aligned} \tag{2.16}$$

where

$$\bar{\rho} \equiv r + i \alpha \cos \theta \quad \text{and} \quad \bar{\rho}^* \equiv r - i \alpha \cos \theta . \quad (2.17)$$

Also, the directional derivatives along the four null vectors are given special symbols,

$$\begin{aligned} \hat{D} &\equiv l^i \partial_i \\ \hat{\Delta} &\equiv n^i \partial_i \\ \hat{\delta} &\equiv m^i \partial_i . \end{aligned} \quad (2.18)$$

In the tetrad formalism covariant derivatives (with respect to a coordinate basis) are replaced by intrinsic derivatives (see Ref. [11]) represented by the symbol $|$. For scalar quantities one has $f|_c = f_{,c}$, for covariant vectors

$$V_{a|c} \equiv V_{a,c} - \eta^{bd} \gamma_{bac} V_d , \quad (2.19)$$

and so forth, where the indices $a, b, \dots = 1, \dots, 4$ are tetrad indices, $\eta^{bd} = \text{diag}(-1, +1, +1, +1)$ is the Minkowski tensor and γ_{bac} are the rotation or *spin coefficients*. For the Kerr-Newman metric they are represented by the following Greek letters and have the expressions [11]

$$\begin{aligned} \kappa = \sigma = \lambda = \nu = \epsilon = 0 \\ \tilde{\rho} = -\frac{1}{\bar{\rho}^*} , \quad \beta = \frac{\cot \theta}{2\sqrt{2}\bar{\rho}} , \quad \pi = \frac{i\alpha \sin \theta}{\sqrt{2}(\bar{\rho}^*)^2} , \quad \tau = -\frac{i\alpha \sin \theta}{\sqrt{2}\bar{\rho}^2} \\ \mu = -\frac{\Delta}{2\rho^2 \bar{\rho}^*} , \quad \gamma = \mu + \frac{r-M}{2\rho^2} , \quad \tilde{\alpha} = \pi - \beta^* . \end{aligned} \quad (2.20)$$

Further, among the Weyl scalars Ψ_k , $k = 0, \dots, 4$, there is only one non vanishing quantity, that is

$$\Psi_2 \equiv C_{ijkl} l^i m^j n^k m^{*l} = -\frac{M}{(\bar{\rho}^*)^3} + \frac{Q^2}{\rho^2 (\bar{\rho}^*)^2} . \quad (2.21)$$

We can thus easily claim that our solution is still of Petrov type D up to $\mathcal{O}(Q^2)$ since the Kerr-Newman one is of this type.

The Maxwell field can be fully described by the following three complex quantities

$$\begin{aligned} \phi_0 &= F_{ij} l^i m^j \\ \phi_1 &= \frac{1}{2} F_{ij} (l^i n^j + m^i m^{*j}) \\ \phi_2 &= F_{ij} m^{*i} n^j , \end{aligned} \quad (2.22)$$

which contain all the information about the six components of the electric and magnetic fields. For the background solution, they are shown to satisfy the *phantom gauge* [11] at order Q^2 ,

$$\phi_0 = \phi_2 = \mathcal{O}(Q^3) , \quad (2.23)$$

with the third scalar given by

$$\phi_1 = -i \frac{Q}{2(\bar{\rho}^*)^2} + \mathcal{O}(Q^3) . \quad (2.24)$$

III. FORMAL DEVELOPMENT OF THE WAVE EQUATIONS

We want now to study the linear perturbations of the static solution given in the previous Section and obtain the wave equations for the various field modes. The full set of equations one obtains in the Newman-Penrose formalism for a metric of the form given in Eq. (2.5) is extraordinarily large, and we refer to the book by Chandrasekhar [11] for the details. In Ref. [11] it is shown that, while in the Kerr background the Maxwell and gravitational wave equations decouple, in the Kerr-Newman background (even without the dilaton field) they do not.

Since our solution at order Q^2 coincides with the Kerr-Newman one as far as the metric is concerned, we cannot thus expect the dilaton, Maxwell and gravitational wave equations to disentangle. However, at order Q^0 it is just the Kerr solution, and we can take advantage of the related simplifications. For this purpose, we perform a double expansion of every relevant quantity,

$$\begin{aligned}\phi(t, r, \theta, \varphi) &= \sum_{p,n} g^p Q^n \phi^{(p,n)} \\ \phi_i(t, r, \theta, \varphi) &= \sum_{p,n} g^p Q^n \phi_i^{(p,n)} , \quad i = 0, 1, 2 \\ G(t, r, \theta, \varphi) &= \sum_{p,n} g^p Q^n G^{(p,n)} ,\end{aligned}\tag{3.1}$$

where g is the same wave parameter for the dilaton, Maxwell and gravitational field quantities, the latter (including tetrad components) being collectively represented with the symbol G . Although we have formally included every order in the wave parameter in the series above, we will only study the linear ($p = 1$) case.

For $p = 0$ the static solutions are

$$\begin{aligned}\phi^{(0,n)} &= \phi^{(0,n)}(r, \theta) \\ \phi_i^{(0,n)} &= \phi_i^{(0,n)}(r, \theta) , \quad i = 0, 1, 2 \\ G^{(0,n)} &= G^{(0,n)}(r, \theta) .\end{aligned}\tag{3.2}$$

At order $(0,0)$ we recover the Kerr solution with neither Maxwell nor dilaton fields. The metric tensor elements for this solution are obtained from those defined in Eq. (2.8) by setting $Q = 0$. At order $(0,1)$ the Maxwell fields appear, corresponding to the metric of the Kerr-Newman solution which is obtained at order $(0,2)$ together with the first non-vanishing contribution to the dilaton background. Finally our solution as given in [9] is fully displayed at order $(0,3)$.

Correspondingly, one expects to obtain at order $(1,0)$ the already known, decoupled and separable wave equations in the Kerr background. At order $(1,1)$ new contributions relating to the presence of the dilaton waves are expected. At order $(1,2)$ the effect induced by the dilaton background should appear. In the following we will show that this is actually the case.

We shall also assume that the perturbations to which the various quantities are subject have the following time and azimuthal dependence

$$\begin{aligned}
\phi^{(1,n)}(t, r, \theta, \varphi) &= k_d e^{i\bar{\omega}t + im\varphi} \phi^{(1,n)}(r, \theta) \\
\phi_i^{(1,n)}(t, r, \theta, \varphi) &= k_{EM} e^{i\bar{\omega}t + im\varphi} \phi_i^{(1,n)}(r, \theta) , \quad i = 0, 1, 2 \\
G^{(1,n)}(t, r, \theta, \varphi) &= k_G e^{i\bar{\omega}t + im\varphi} G^{(1,n)}(r, \theta) ,
\end{aligned} \tag{3.3}$$

where k_d , k_{EM} and k_G are (possibly equal) parameters. We note here that each function of r and θ on the R.H.S.s above implicitly carries an extra integer index, m , and the continuous dependence on the frequency $\bar{\omega}$ (both can be positive or negative).

The directional derivatives when acting on the wave modes displayed above can be written

$$\begin{aligned}
\hat{D} &= \mathcal{D}_0 \\
\hat{\Delta} &= -\frac{\Delta}{2\rho^2} \mathcal{D}_0^\dagger \\
\hat{\delta} &= \frac{1}{\sqrt{2}\bar{\rho}} \mathcal{L}_0^\dagger \\
\hat{\delta}^* &= \frac{1}{\sqrt{2}\bar{\rho}^*} \mathcal{L}_0 ,
\end{aligned} \tag{3.4}$$

where

$$\begin{aligned}
\mathcal{D}_n &= \partial_r + i \frac{K}{\Delta} + 2n \frac{r-M}{\Delta} \\
\mathcal{D}_n^\dagger &= \partial_r - i \frac{K}{\Delta} + 2n \frac{r-M}{\Delta} \\
\mathcal{L}_n &= \partial_\theta + \tilde{Q} + n \cot \theta \\
\mathcal{L}_n^\dagger &= \partial_\theta - \tilde{Q} + n \cot \theta ,
\end{aligned} \tag{3.5}$$

with n an integer such that $n \geq 0$ and

$$\begin{aligned}
K &\equiv (r^2 + \alpha^2) \bar{\omega} + \alpha m \\
\tilde{Q} &\equiv \alpha \bar{\omega} \sin \theta + m \csc \theta .
\end{aligned} \tag{3.6}$$

A. Gravitational equations

Following Ref. [15], we consider the following three non vacuum equations,

$$\begin{aligned}
(\hat{\delta}^* - 4\tilde{\alpha} + \pi) \Psi_0 - (\hat{D} - 4\tilde{\rho} - 2\epsilon) \Psi_1 - 3\kappa \Psi_2 &= (\hat{\delta} + \pi^* - 2\tilde{\alpha}^* - 2\beta) R_{11} \\
&\quad - (\hat{D} - 2\epsilon - 2\tilde{\rho}^*) R_{12} + 2\sigma R_{21} - 2\kappa R_{22} - \kappa^* R_{13} \\
(\hat{\Delta} - 4\gamma + \mu) \Psi_0 - (\hat{\delta} - 4\tau - 2\beta) \Psi_1 - 3\sigma \Psi_2 &= (\hat{\delta} + 2\pi^* - 2\beta) R_{12} \\
&\quad - (\hat{D} - 2\epsilon + 2\epsilon^* - \tilde{\rho}^*) R_{13} - \lambda^* R_{11} - 2\sigma R_{22} - 2\kappa R_{23} \\
(\hat{D} - \tilde{\rho} - \tilde{\rho}^* - 3\epsilon + \epsilon^*) \sigma - (\hat{\delta} - \tau + \pi^* - \tilde{\alpha}^* - 3\beta) \kappa - \Psi_0 &= 0 .
\end{aligned} \tag{3.7}$$

The Ricci tensor terms are given by the Einstein field equations, Eq. (2.3),

$$R_{ab} \equiv R_{ij} e_a^i e_b^j = \frac{1}{2} \phi_{|a} \phi_{|b} + 2 T_{ab}^{EM} . \quad (3.8)$$

Now we expand the equations above according to the expression in Eq. (3.1).

For $p = 0$ we recover the static solution described in the previous Section. In particular one has

$$\begin{aligned} R_{ab}^{(0,0)} &= R_{ab}^{(0,1)} = 0 \\ \Psi_k^{(0,0)} &= \Psi_k^{(0,1)} = \Psi_2^{(0,1)} = 0 , \quad k = 0, 1, 3, 4 \\ \kappa^{(0,0)} &= \sigma^{(0,0)} = \nu^{(0,0)} = \lambda^{(0,0)} = \epsilon^{(0,0)} = 0 \\ \kappa^{(0,1)} &= \sigma^{(0,1)} = \nu^{(0,1)} = \lambda^{(0,1)} = \epsilon^{(0,1)} = 0 , \end{aligned} \quad (3.9)$$

which will be used to simplify the expressions at order $p = 1$.

We can express the (wave) perturbation

$$(e_a^i)^{(1)} \equiv \sum_n Q^n (e_a^i)^{(1,n)} , \quad (3.10)$$

in the tetrad vectors as a linear combination of the unperturbed basis vectors $(e_a^i)^{(0,n)}$ [11],

$$(e_a^i)^{(1,n)} = \sum_{q+r=n} (A_a^b)^{(q)} (e_a^i)^{(0,r)} . \quad (3.11)$$

The perturbations in the basis vectors are then fully described by the matrix

$$A_a^b = \sum_n Q^n (A_a^b)^{(n)} , \quad (3.12)$$

whose elements A_1^1 , A_2^2 , A_2^1 and A_1^2 are real, while all the others are complex (complex conjugate to each other when changing any index 3 with 4 or viceversa). It then follows that A_a^b has sixteen independent components, and it is also useful to define

$$\begin{aligned} F_2^1 &\equiv \frac{\Delta}{2\rho^2} A_2^1 , & F_1^2 &\equiv \frac{2\rho^2}{\Delta} A_1^2 , & F_1^3 &\equiv \frac{1}{\bar{\rho}^*} A_1^3 , & F_1^4 &\equiv \frac{1}{\bar{\rho}} A_1^4 \\ F_3^1 &\equiv \frac{\Delta}{2\rho^2 \bar{\rho}} A_3^1 , & F_3^2 &\equiv \frac{1}{\bar{\rho}} A_3^2 , & F_2^3 &\equiv \frac{\Delta}{2\rho^2 \bar{\rho}^*} A_2^3 , & F_2^4 &\equiv \frac{\Delta}{2\rho^2 \bar{\rho}} A_2^4 \\ F_4^1 &\equiv \frac{\Delta}{2\rho^2 \bar{\rho}^*} A_4^1 , & F_4^2 &\equiv \frac{1}{\bar{\rho}^*} A_4^2 , & F_4^3 &\equiv \frac{1}{(\bar{\rho}^*)^2} A_4^3 , & F_3^4 &\equiv \frac{1}{(\bar{\rho})^2} A_3^4 , \end{aligned} \quad (3.13)$$

and

$$\begin{aligned} F &= F_3^1 + F_4^1 , & B_1 &= F_1^3 + F_2^3 + F_1^4 + F_2^4 \\ G &= F_3^2 + F_4^2 , & B_2 &= F_1^3 + F_2^3 - F_1^4 - F_2^4 \\ H &= F_3^1 + F_2^3 , & C_1 &= F_3^1 + F_3^2 - F_4^1 - F_4^2 \\ J &= F_1^4 - F_2^4 , & C_2 &= F_3^1 - F_3^2 - F_4^1 + F_4^2 \\ U &= A_1^1 + A_2^2 , & V &= A_3^3 + A_4^4 . \end{aligned} \quad (3.14)$$

We are now ready to study the perturbations with $p = 1$.

1. (1, 0) order

It is easy to see that

$$R_{ab}^{(1,0)} = 0 , \quad (3.15)$$

so that both the electromagnetic field and the dilaton field decouple from the gravitational field equations. Further, on also using Eq. (3.9), the system of gravitational equations reduces to

$$\begin{aligned} (\hat{\delta}^* - 4\tilde{\alpha} + \pi)^{(0,0)} \Psi_0^{(1,0)} - (\hat{D} - 4\tilde{\rho})^{(0,0)} \Psi_1^{(1,0)} - 3\kappa^{(1,0)} \Psi_2^{(0,0)} &= 0 \\ (\hat{\Delta} - 4\gamma + \mu)^{(0,0)} \Psi_0^{(1,0)} - (\hat{\delta} - 4\tau - 2\beta)^{(0,0)} \Psi_1^{(1,0)} - 3\sigma^{(1,0)} \Psi_2^{(0,0)} &= 0 \\ (\hat{D} - \tilde{\rho} - \tilde{\rho}^*)^{(0,0)} \sigma^{(1,0)} - (\hat{\delta} - \tau + \pi^* - \tilde{\alpha}^* - 3\beta)^{(0,0)} \kappa^{(1,0)} - \Psi_0^{(1,0)} &= 0 , \end{aligned} \quad (3.16)$$

which are the equations for gravitational waves in the Kerr background [15].

2. (1, $n \geq 1$) order

From the expression of the Ricci tensor, Eq. (3.8), one obtains that the background dilaton field $\phi^{(0,2)}$ enters into the equations for the gravitational waves only at order Q^n with $n \geq 2$.

In fact, at order (1, 1) one has

$$\begin{aligned} R_{ab}^{(1,1)} &= 2T_{ab}^{EM(1,1)} \\ &= 2F_{ik}^{(1,0)} F_j^{k(0,1)} + 2F_{ik}^{(0,1)} F_j^{k(1,0)} \\ &\quad - \frac{1}{2} g_{ij}^{(0,0)} \left(F_{kl}^{(1,0)} F^{kl(0,1)} + F_{kl}^{(0,1)} F^{kl(1,0)} \right) . \end{aligned} \quad (3.17)$$

Thus the dilaton field does not affect the equations at this order in the Q expansion, neither directly nor through its effect on the Maxwell waves ($F_{ij}^{(1,0)}$ are free electromagnetic waves in Kerr background, see Section III C below). However, both Maxwell background and wave fields already enter into the gravitational equations, so that for $n \geq 1$ it is no longer possible to decouple gravitational waves from Maxwell excitations.

At order (1, 2) one obtains terms containing products of the form $\phi_{|a}^{(0,2)} \phi_{|b}^{(1,0)}$ which couple the dilaton waves to their own background. Further, one expects corrections coming from effects induced on the electromagnetic waves. For this reason and because of the complex entanglement among all kinds of waves, we do not address this issue in the present paper.

We just notice here that it is only at order (1, 4) that purely background terms like $\phi_{|a}^{(0,2)} \phi_{|b}^{(0,2)}$ appear. However, our knowledge of the static solution does not allow us to write explicitly the equations in Eq. (3.7) at this order, since we do not know the gravitational terms $G^{(0,4)}$.

B. Dilaton equation

The equation for the dilaton field, Eq. (2.2), in tetrad components becomes

$$\eta^{ab} \phi_{a|b} = -a e^{-a\phi} F^2, \quad (3.18)$$

where $\phi_a \equiv e_a^i \phi_{,i}$ are four (complex) scalars.

In the Newman-Penrose tetrad, Eq. (3.18) becomes

$$\begin{aligned} & \left[\hat{D} \hat{\Delta} + \hat{\Delta} \hat{D} - \hat{\delta} \hat{\delta}^* - \hat{\delta}^* \hat{\delta} + (\epsilon + \epsilon^* - \tilde{\rho} - \tilde{\rho}^*) \hat{\Delta} + (\mu + \mu^* - \gamma - \gamma^*) \hat{D} \right. \\ & \left. + (\tau - \pi^* + \tilde{\alpha}^* - \beta) \hat{\delta}^* + (\tau^* - \pi + \tilde{\alpha} - \beta^*) \hat{\delta} \right] \phi = -a e^{-a\phi} F^2, \end{aligned} \quad (3.19)$$

which can now be expanded in Q and the wave parameter.

At order $(0, n)$, $n = 0, 1, 2$, one recovers the static solutions

$$\begin{aligned} \phi^{(0,0)} &= \phi^{(0,1)} = 0 \\ \phi^{(0,2)} &= -\frac{a}{M} \frac{r}{\rho^2}, \end{aligned} \quad (3.20)$$

and we now study the linear perturbations $p = 1$.

1. (1, 0) order

The R.H.S. of Eq. (3.19) vanishes and one finds the Klein-Gordon equation for a free scalar field in Kerr space-time,

$$\begin{aligned} & \left[\hat{D} \hat{\Delta} + \hat{\Delta} \hat{D} - \hat{\delta} \hat{\delta}^* - \hat{\delta}^* \hat{\delta} - (\tilde{\rho} + \tilde{\rho}^*) \hat{\Delta} + (\mu + \mu^* - \gamma - \gamma^*) \hat{D} \right. \\ & \left. + (\tau - \pi^* + \tilde{\alpha}^* - \beta) \hat{\delta}^* + (\tau^* - \pi + \tilde{\alpha} - \beta^*) \hat{\delta} \right]^{(0,0)} \phi^{(1,0)} = 0. \end{aligned} \quad (3.21)$$

This can be further simplified to the final form

$$\left[\Delta_0 \mathcal{D}_1 \mathcal{D}_0^\dagger + \mathcal{L}_0^\dagger \mathcal{L}_1 + 2i\bar{\omega} \bar{\rho} \right] \Phi = 0, \quad (3.22)$$

with $\Phi \equiv \phi^{(1,0)}$, and the operators inside the square brackets are to be evaluated at order $(0, 0)$.

The equation above is separable. One can write

$$\Phi(r, \theta) = R_0(r) S_0(\theta), \quad (3.23)$$

and then refer to the general case for (half) integer spin s functions $R_s(r) S_s(\theta)$ described in Section III C.

2. $(1, n \geq 1)$ order

As for the gravitational waves, the dilaton background $\phi^{(0,2)}$ enters into Eq. (3.19) only at order Q^2 and higher.

At order $(1, 1)$ the R.H.S. of Eq. (3.19) is different from zero and couples the scalar field waves to the Maxwell field,

$$\begin{aligned} \left[\Delta_0 \mathcal{D}_1 \mathcal{D}_0^\dagger + \mathcal{L}_0^\dagger \mathcal{L}_1 + 2i\bar{\omega} \bar{\rho} \right] \phi^{(1,1)} &= -a \frac{k_{EM}}{k_d} \left(F_{ab}^{(1,0)} F^{ab(0,1)} + F_{ab}^{(0,1)} F^{ab(1,0)} \right) \\ &= -4a \frac{k_{EM}}{k_d} \left(\phi_1^{(1,0)} \phi_1^{(0,1)} - \phi_1^{*(1,0)} \phi_1^{*(0,1)} \right), \end{aligned} \quad (3.24)$$

where use has been made of the phantom gauge, Eq. (2.23), and $\phi_1^{(1,0)}$ is the Maxwell wave in the Kerr background, which we will review in the next Subsection. Here we only observe that Eq. (3.24) does not contain a contribution from the static dilaton background, neither explicitly nor implicitly in the metric and Maxwell fields.

At order $(1, 2)$ the L.H.S. of Eq. (3.19) contains terms of the form $G^{(1,0)} \phi^{(0,2)}$ which couple the gravitational waves to the dilaton background as claimed.

C. Maxwell equations

The second equation for the Maxwell field in Eq. (2.1) is automatically satisfied once we introduce the three complex scalars ϕ_i , and we are then left with only the first equation which can be written in a general tetrad frame as

$$F_{ab|c} = a \phi_{|c} F_{ab}. \quad (3.25)$$

Thus one clearly sees that the dilaton generates a source term for the electromagnetic field.

Using the symbols for directional derivatives and spin coefficients introduced in the previous Section, Eq. (3.25) can be written as the following set of four complex equations

$$\begin{aligned} (\hat{D} - 2\bar{\rho}) \phi_1 - (\hat{\delta}^* + \pi - 2\bar{\alpha}) \phi_0 + \kappa \phi_2 &= J_1 \\ (\hat{\delta} - 2\tau) \phi_1 - (\hat{\Delta} + \mu - 2\gamma) \phi_0 + \sigma \phi_2 &= J_3 \\ (\hat{D} - \bar{\rho} + 2\epsilon) \phi_2 - (\hat{\delta}^* + 2\pi) \phi_1 + \lambda \phi_0 &= J_4 \\ (\hat{\delta} - \tau + 2\beta) \phi_2 - (\hat{\Delta} + 2\mu) \phi_1 + \nu \phi_0 &= J_2, \end{aligned} \quad (3.26)$$

where the source terms are given by

$$\begin{aligned} J_1 &= \frac{a}{2} \left[(\phi_1 + \phi_1^*) \hat{D} - \phi_0 \hat{\delta}^* - \phi_0^* \hat{\delta} \right] \phi \\ J_2 &= \frac{a}{2} \left[(\phi_1 + \phi_1^*) \hat{\Delta} - \phi_2 \hat{\delta} - \phi_2^* \hat{\delta}^* \right] \phi \\ J_3 &= \frac{a}{2} \left[(\phi_1 - \phi_1^*) \hat{\delta} - \phi_0 \hat{\Delta} + \phi_2^* \hat{D} \right] \phi \\ J_4 &= \frac{a}{2} \left[(\phi_1 - \phi_1^*) \hat{\delta}^* + \phi_0^* \hat{\Delta} - \phi_2 \hat{D} \right] \phi. \end{aligned} \quad (3.27)$$

We can now expand according to Eq. (3.1).

1. (1, 0) order

We observe that the background dilaton field and the Maxwell background field are zero at order Q^0 . This makes the source terms vanish and, using the vanishing of some of the spin coefficients according to Eq. (3.9), one obtains

$$\begin{aligned}
(\hat{D} - 2\tilde{\rho})^{(0,0)} \phi_1^{(1,0)} - (\hat{\delta}^* + \pi - 2\tilde{\alpha})^{(0,0)} \phi_0^{(1,0)} &= 0 \\
(\hat{\delta} - 2\tau)^{(0,0)} \phi_1^{(1,0)} - (\hat{\Delta} + \mu - 2\gamma)^{(0,0)} \phi_0^{(1,0)} &= 0 \\
(\hat{D} - \tilde{\rho})^{(0,0)} \phi_2^{(1,0)} - (\hat{\delta}^* + 2\pi)^{(0,0)} \phi_1^{(1,0)} &= 0 \\
(\hat{\delta} - \tau + 2\beta)^{(0,0)} \phi_2^{(1,0)} - (\hat{\Delta} + 2\mu)^{(0,0)} \phi_1^{(1,0)} &= 0 .
\end{aligned} \tag{3.28}$$

These are the equations for electromagnetic waves $\phi_i^{(1,0)}$ moving in the Kerr background. Their properties and solutions have already been broadly studied in the literature and we now briefly summarize them for the purpose of introducing some useful notation [11].

We replace the directional derivatives and spin coefficients with their explicit forms as given respectively in Eq. (3.5) and Eq. (2.20) and obtain

$$\begin{aligned}
\left(\mathcal{L}_1 - \frac{i\alpha \sin \theta}{\bar{\rho}^*}\right) \Phi_0 - \left(\mathcal{D}_0 + \frac{1}{\bar{\rho}^*}\right) \Phi_1 &= 0 \\
\left(\mathcal{L}_0 + \frac{i\alpha \sin \theta}{\bar{\rho}^*}\right) \Phi_1 - \left(\mathcal{D}_0 - \frac{1}{\bar{\rho}^*}\right) \Phi_2 &= 0 \\
\left(\mathcal{L}_1^\dagger - \frac{i\alpha \sin \theta}{\bar{\rho}^*}\right) \Phi_2 + \Delta_0 \left(\mathcal{D}_0^\dagger + \frac{1}{\bar{\rho}^*}\right) \Phi_1 &= 0 \\
\left(\mathcal{L}_0^\dagger + \frac{i\alpha \sin \theta}{\bar{\rho}^*}\right) \Phi_1 + \Delta_0 \left(\mathcal{D}_1^\dagger - \frac{1}{\bar{\rho}^*}\right) \Phi_0 &= 0 ,
\end{aligned} \tag{3.29}$$

where $\Delta_0 \equiv \Delta^{(0,0)}$, and we have also introduced

$$\begin{aligned}
\Phi_0 &= \phi_0^{(1,0)} \\
\Phi_1 &= \sqrt{2} \bar{\rho}^* \phi_1^{(1,0)} \\
\Phi_2 &= 2 (\bar{\rho}^*)^2 \phi_2^{(1,0)} .
\end{aligned} \tag{3.30}$$

On using the following commutation properties

$$\begin{aligned}
\left[\left(\mathcal{D}_0 + \frac{1}{\bar{\rho}^*}\right), \left(\mathcal{L}_1^\dagger + \frac{i\alpha \sin \theta}{\bar{\rho}^*}\right)\right] &= 0 \\
\left[\Delta_0 \left(\mathcal{D}_0^\dagger + \frac{1}{\bar{\rho}^*}\right), \left(\mathcal{L}_0 + \frac{i\alpha \sin \theta}{\bar{\rho}^*}\right)\right] &= 0 ,
\end{aligned} \tag{3.31}$$

one can obtain separate equations for Φ_0 and Φ_2 ,

$$\begin{aligned}
\left[\Delta_0 \mathcal{D}_0 \mathcal{D}_0^\dagger + \mathcal{L}_0^\dagger \mathcal{L}_1 - 2i\bar{\omega} \bar{\rho}\right] \Delta_0 \Phi_0 &= 0 \\
\left[\Delta_0 \mathcal{D}_0^\dagger \mathcal{D}_0 + \mathcal{L}_0 \mathcal{L}_1^\dagger + 2i\bar{\omega} \bar{\rho}\right] \Phi_2 &= 0 .
\end{aligned} \tag{3.32}$$

The latter are separable equations and we can factorize their solutions

$$\begin{aligned}\Delta_0 \Phi_0 &= P_{+1}(r) S_{+1}(\theta) , \quad P_{+1} = \Delta_0 R_{+1} \\ \Phi_2 &= P_{-1}(r) S_{-1}(\theta) , \quad P_{-1} = R_{-1} .\end{aligned}\tag{3.33}$$

The radial equations for the functions $P_{s=\pm 1}$ are particular cases of the eigenvalue equations,

$$\begin{aligned}\left[\Delta_0 \mathcal{D}_{1-|s|} \mathcal{D}_0^\dagger - 2i(2|s| - 1)\bar{\omega}r \right] P_{|s|} &= E P_{|s|} \\ \left[\Delta_0 \mathcal{D}_{1-|s|}^\dagger \mathcal{D}_0 + 2i(2|s| - 1)\bar{\omega}r \right] P_{-|s|} &= E P_{-|s|} ,\end{aligned}\tag{3.34}$$

where $P_{|s|} \equiv \Delta_0^s R_s$ and $P_{-|s|} \equiv R_{-s}$, with s integer or half integer. The angular equations for $S_{\pm 1}$ are particular cases of the following equations

$$\begin{aligned}\left[\mathcal{L}_0^\dagger \mathcal{L}_{|s|} + 2\bar{\omega}\alpha \cos \theta \right] S_{|s|} &= -E S_{|s|} \\ \left[\mathcal{L}_0 \mathcal{L}_{|s|}^\dagger - 2\bar{\omega}\alpha \cos \theta \right] S_{-|s|} &= -E S_{-|s|} ,\end{aligned}\tag{3.35}$$

where E is the same (separation) constant which appears in Eq. (3.34).

For any (half) integers $|s| \leq l$ and $|m| \leq 2l + 1$ one has

$$S_{sl}^m(\theta, \bar{\omega}) e^{im\varphi} = Y_{sl}^m(\theta, \varphi; \bar{\omega}) ,\tag{3.36}$$

where $Y(\bar{\omega})$ are *spin-weighted spheroidal harmonics* [16] which form a complete, orthonormal set of functions for every value of s . They reduce to the *spin-weighted spherical harmonics*,

$$Y_{sl}^m(\theta, \varphi) = S_{sl}^m(\theta) e^{im\varphi} ,\tag{3.37}$$

in the limit $\bar{\omega} = 0$ and to the usual spherical harmonics when one has also $s = 0$.

Since the spin-weighted spherical harmonics of a fixed spin s are complete and orthonormal as well, one can perform the following expansion

$$S_{sl}^m(\theta; \bar{\omega}) = \sum_{l'} A_{ss',ll'}^m(\bar{\omega}) S_{s'l'}^m(\theta) ,\tag{3.38}$$

where the $A_{ss',ll'}^m$ are related to Clebsch-Gordan coefficients.

We now introduce the following expansions for the Maxwell wave modes $\Phi_i = \Phi_i^m(r, \theta; \bar{\omega})$ of energy $\bar{\omega}$ and angular momentum m along the axis of symmetry,

$$\begin{aligned}\Phi_i(r, \theta) &= \sum_{l \geq (|m|-1)/2} C_{sl}^m(\bar{\omega}) R_{sl}^m(r, \bar{\omega}) S_{sl}^m(\theta, \bar{\omega}) \\ &= \sum_{l', l \geq (|m|-1)/2} B_{ss',ll'}^m(\bar{\omega}) R_{sl}^m(r, \bar{\omega}) S_{s'l'}^m(\theta) ,\end{aligned}\tag{3.39}$$

where $s = +1$ for $i = 0$ and $s = -1$ for $i = 2$, the functions $R(r, \bar{\omega})$ are solutions of the radial equations, Eq. (3.34), $C_{sl}^m(\bar{\omega})$ are normalization constants and

$$B_{ss',ll'}^m(\bar{\omega}) \equiv C_{sl}^m(\bar{\omega}) A_{ss',ll'}^m(\bar{\omega}) ,\tag{3.40}$$

are related to Clebsch-Gordan coefficients and project the angular solutions on the complete basis of a generic spin s' . By analogy, the dilaton wave Φ can be expanded as

$$\begin{aligned}\Phi(r, \theta) &= \sum_{l \geq (|m|-1)/2} C_{0l}^m(\bar{\omega}) R_{0l}^m(r, \bar{\omega}) S_{0l}^m(\theta, \bar{\omega}) \\ &= \sum_{l', l \geq (|m|-1)/2} B_{00, l'}^m(\bar{\omega}) R_{0l}^m(r, \bar{\omega}) S_{l'}^m(\theta) ,\end{aligned}\quad (3.41)$$

where $S_l^m e^{im\varphi}$ are spherical harmonics.

2. (1, 1) order

Since for the background solution one has $G^{(0,1)} = \phi_0^{(0,1)} = \phi_1^{(0,1)} = 0$, the first equation in Eq. (3.26) can be written,

$$\begin{aligned}k_{EM} \left[(\hat{D} - 2\tilde{\rho})^{(0,0)} \phi_1^{(1,1)} - (\hat{\delta}^* + \pi - 2\tilde{\alpha})^{(0,0)} \phi_0^{(1,1)} \right] + k_G (\hat{D} - 2\tilde{\rho})^{(1,0)} \phi_1^{(0,1)} \\ = \frac{a}{2} k_d (\phi_1 + \phi_1^*)^{(0,1)} \hat{D}^{(0,0)} \phi^{(1,0)} ,\end{aligned}\quad (3.42)$$

which contains contributions from both the dilaton waves $\phi^{(1,0)}$ and the gravitational waves $G^{(1,0)}$.

However, it is possible to show that the gravitational waves actually cancel out in the Kerr background [11]. In fact, one has

$$\begin{aligned}(\hat{D} - 2\tilde{\rho})^{(1,0)} &= -(F_2^1)^{(0)} \left(\partial_r + \frac{2}{\bar{\rho}^*} \right) + \frac{\sqrt{2}\rho^2}{\Delta_0} F^{(0)} \left(\partial_\theta + \frac{2i\alpha \sin \theta}{\bar{\rho}^*} \right) \\ (\hat{\delta} - 2\tau)^{(1,0)} &= -\bar{\rho}^* H^{(0)} \left(\partial_r + \frac{2}{\bar{\rho}^*} \right) + \frac{\bar{\rho}^*}{\sqrt{2}} (F_4^3)^{(0)} \left(\partial_\theta + \frac{2i\alpha \sin \theta}{\bar{\rho}^*} \right) \\ (\hat{\delta}^* + 2\pi)^{(1,0)} &= -\bar{\rho} J^{(0)} \left(\partial_r + \frac{2}{\bar{\rho}^*} \right) + \frac{\bar{\rho}}{\sqrt{2}} (F_3^4)^{(0)} \left(\partial_\theta + \frac{2i\alpha \sin \theta}{\bar{\rho}^*} \right) \\ (\hat{\Delta} + 2\mu)^{(1,0)} &= -\frac{\Delta_0}{2\rho^2} (F_1^2)^{(0)} \left(\partial_r + \frac{2}{\bar{\rho}^*} \right) + \frac{1}{\sqrt{2}} G^{(0)} \left(\partial_\theta + \frac{2i\alpha \sin \theta}{\bar{\rho}^*} \right) ,\end{aligned}\quad (3.43)$$

where the functions F_b^a , F , H , G are defined in Eqs. (3.13), (3.14). Further, from Eq. (2.24), one finds

$$\left(\partial_r + \frac{2}{\bar{\rho}^*} \right) \phi_1^{(0,1)} = \left(\partial_\theta + \frac{2i\alpha \sin \theta}{\bar{\rho}^*} \right) \phi_1^{(0,1)} = 0 ,\quad (3.44)$$

so that all the gravitational wave terms vanish.

One is then left with the following four equations

$$\begin{aligned}(\hat{D} - 2\tilde{\rho}) \phi_1^{(1,1)} - (\hat{\delta}^* + \pi - 2\tilde{\alpha}) \phi_0^{(1,1)} &= a \frac{k_d}{k_{EM}} \phi_1^R \hat{D} \Phi \\ (\hat{\delta} - 2\tau) \phi_1^{(1,1)} - (\hat{\Delta} + \mu - 2\gamma) \phi_0^{(1,1)} &= a \frac{k_d}{k_{EM}} \phi_1^I \hat{\delta} \Phi\end{aligned}$$

$$\begin{aligned}
(\hat{D} - \tilde{\rho}) \phi_2^{(1,1)} - (\hat{\delta}^* + 2\pi) \phi_1^{(1,1)} &= a \frac{k_d}{k_{EM}} \phi_1^I \hat{\delta}^* \Phi \\
(\hat{\delta} - \tau + 2\beta) \phi_2^{(1,1)} - (\hat{\Delta} + 2\mu) \phi_1^{(1,1)} &= a \frac{k_d}{k_{EM}} \phi_1^R \hat{\Delta} \Phi ,
\end{aligned} \tag{3.45}$$

where we have defined $\phi_1^{R/I} \equiv (\phi_1 \pm \phi_1^*)^{(0,1)}/2$, $\Phi = \phi^{(1,0)}$ and all gravitational quantities are understood to be of order $(0,0)$.

The L.H.S. of Eq. (3.45) above is the same as the L.H.S. of Eq. (3.28), so we introduce

$$\begin{aligned}
W_0 &= \phi_0^{(1,1)} \\
W_1 &= \sqrt{2} \bar{\rho}^* \phi_1^{(1,1)} \\
W_2 &= 2 (\bar{\rho}^*)^2 \phi_2^{(1,1)} \\
\Phi_1^{I/R} &= \sqrt{2} \bar{\rho}^* \phi_1^{I/R} ,
\end{aligned} \tag{3.46}$$

and follow the same steps previously done at order $(1,0)$. We then obtain separate equations for W_0 and W_2 ,

$$\begin{aligned}
\left[\left(\mathcal{L}_0^\dagger + \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) \left(\mathcal{L}_1 - \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) + \Delta_0 \left(\mathcal{D}_1 + \frac{1}{\bar{\rho}^*} \right) \left(\mathcal{D}_1^\dagger - \frac{1}{\bar{\rho}^*} \right) \right] W_0 &= T_+ \\
\left[\left(\mathcal{L}_0 + \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) \left(\mathcal{L}_1^\dagger - \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) + \Delta_0 \left(\mathcal{D}_0^\dagger + \frac{1}{\bar{\rho}^*} \right) \left(\mathcal{D}_0 - \frac{1}{\bar{\rho}^*} \right) \right] W_2 &= \Delta_0 T_- ,
\end{aligned} \tag{3.47}$$

which can be further simplified to the final form

$$\begin{aligned}
[\Delta_0 \mathcal{D}_0 \mathcal{D}_0^\dagger + \mathcal{L}_0^\dagger \mathcal{L}_1 - 2i\bar{\omega} \bar{\rho}] \Delta_0 W_0 &= \Delta_0 T_+ \\
[\Delta_0 \mathcal{D}_0^\dagger \mathcal{D}_0 + \mathcal{L}_0 \mathcal{L}_1^\dagger + 2i\bar{\omega} \bar{\rho}] W_2 &= \Delta_0 T_- .
\end{aligned} \tag{3.48}$$

We observe that the L.H.S.s of the equations above are again separable, since they coincide with the expressions one gets at order $(1,0)$.

The currents on the R.H.S.s are given by

$$\begin{aligned}
T_+ &= -a \frac{k_d}{k_{EM}} \left[\left(\mathcal{D}_0 + \frac{1}{\bar{\rho}^*} \right) \Phi_1^I \mathcal{L}_0^\dagger - \left(\mathcal{L}_0^\dagger + \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) \Phi_1^R \mathcal{D}_0 \right] \Phi \\
&= \frac{\sqrt{2}}{2} a \frac{k_d}{k_{EM}} \left[\left(\partial_r + i \frac{K}{\Delta_0} + \frac{1}{\bar{\rho}^*} \right) i \bar{\rho}^* \partial_r \left(\frac{r}{\rho^2} \right) (\partial_\theta - \tilde{Q}) \right. \\
&\quad \left. - \left(\partial_\theta - \tilde{Q} + \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) \frac{\bar{\rho}^*}{\alpha \sin \theta} \partial_\theta \left(\frac{r}{\rho^2} \right) \left(\partial_r + i \frac{K}{\Delta_0} \right) \right] \Phi \\
T_- &= a \frac{k_d}{k_{EM}} \left[\left(\mathcal{L}_0 + \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) \Phi_1^R \mathcal{D}_0^\dagger - \left(\mathcal{D}_0^\dagger + \frac{1}{\bar{\rho}^*} \right) \Phi_1^I \mathcal{L}_0 \right] \Phi \\
&= \frac{\sqrt{2}}{2} a \frac{k_d}{k_{EM}} \left[\left(\partial_r - i \frac{K}{\Delta_0} + \frac{1}{\bar{\rho}^*} \right) i \bar{\rho}^* \partial_r \left(\frac{r}{\rho^2} \right) (\partial_\theta + \tilde{Q}) \right. \\
&\quad \left. - \left(\partial_\theta + \tilde{Q} + \frac{i\alpha \sin \theta}{\bar{\rho}^*} \right) \frac{\bar{\rho}^*}{\alpha \sin \theta} \partial_\theta \left(\frac{r}{\rho^2} \right) \left(\partial_r - i \frac{K}{\Delta_0} \right) \right] \Phi ,
\end{aligned} \tag{3.49}$$

and they cannot be written as the sums of functions of r only and θ only.

However one can expand the currents on the angular basis functions we have introduced at order $(1, 0)$,

$$\begin{aligned} J_{\pm,l}^m(r, \bar{\omega}) &= \Delta_0(r) \int_{-1}^{+1} d(\cos \theta) T_{\pm}(r, \theta) S_{\pm 1,l}^m(\theta, \bar{\omega}) \\ &= \Delta_0(r) \sum_{l'} \int_{-1}^{+1} d(\cos \theta) T_{\pm}(r, \theta) B_{0,l'}^m(\bar{\omega}) S_{l'}^m(\theta) , \end{aligned} \quad (3.50)$$

and define

$$\begin{aligned} W_{+,l}^m(r, \bar{\omega}) &= \Delta_0(r) \int_{-1}^{+1} d(\cos \theta) W_0(r, \theta) S_{+1,l}^m(\theta, \bar{\omega}) \\ W_{-,l}^m(r, \bar{\omega}) &= \int_{-1}^{+1} d(\cos \theta) W_2(r, \theta) S_{-1,l}^m(\theta, \bar{\omega}) . \end{aligned} \quad (3.51)$$

One then obtains the following radial equations for the functions W_{\pm} ,

$$\begin{aligned} \left[\Delta_0 \mathcal{D}_0 \mathcal{D}_0^\dagger - 2i\bar{\omega}r - E_{+1,l}^m \right] W_{+,l}^m &= J_{+,l}^m \\ \left[\Delta_0 \mathcal{D}_0^\dagger \mathcal{D}_0 + 2i\bar{\omega}r - E_{-1,l}^m \right] W_{-,l}^m &= J_{-,l}^m . \end{aligned} \quad (3.52)$$

Given the explicit form of the currents, analytic solutions of these equations do not appear to be feasible.

In this Section we only observe that, without the dilaton ($a = 0$), one would not expect any dependence on the charge Q in the form of the Maxwell waves at order $(1, 1)$, that is we can assume

$$\phi_i^{(1,1)}(a = 0) = 0 . \quad (3.53)$$

But $\phi_i^{(1,1)}(a = 0)$ are also the solutions of the homogeneous equations derived by setting T_{\pm} to zero in Eq. (3.48). It thus follows that the particular solutions W_0 and W_2 of the inhomogeneous Eq. (3.48) precisely represent a purely dilatonic effect.

3. $(1, 2)$ order

So far, the dilaton background field has not appeared in our Maxwell equations. However, one expects $\phi^{(0,2)}$ to enter in the expressions of the currents in the R.H.S.s of Eq. (3.26) at order $(1, 2)$ and in fact it does. Unfortunately, the L.H.S.s of the same equations now contain both gravitational and electromagnetic waves, which are known to be coupled in the Kerr-Newman background [11].

In order to proceed, one can make the following working *ansatz*: we assume

$$k_{EM} \gg k_d, k_G , \quad (3.54)$$

and neglect both dilaton and gravitational waves. This hypothesis is equivalent to assuming (space-time) boundary conditions such that the gravitational and dilaton contents of the wave field are negligibly small when compared to the electromagnetic sources.

We also observe that the currents T_{\pm} at order $(1, 1)$ are proportional to k_d/k_{EM} and become negligible when Eq. (3.54) is satisfied. Thus the corresponding particular solutions $\phi_i^{(1,1)}$ for the inhomogeneous Maxwell equations vanish as well. On the other hand, when Eq. (3.54) does not apply ($k_d \sim k_{EM}$), one can trust the solutions we found at order $(1, 1)$ and neglect the present order of approximation (Q^2). This simple argument shows that the two cases, order $(1, 1)$ with $k_d \sim k_{EM}$ and order $(1, 2)$ with Eq. (3.54), are sufficient to cover most of the electromagnetic physics one expects to be affected by the existence of a dilaton field.

The four Maxwell equations with Eq. (3.54) then read

$$\begin{aligned} (\hat{D} - 2\tilde{\rho})^{(0,0)} \phi_1^{(1,2)} - (\hat{\delta}^* + \pi - 2\tilde{\alpha})^{(0,0)} \phi_0^{(1,2)} &= J_1^{(1,2)} \\ (\hat{\delta} - 2\tau)^{(0,0)} \phi_1^{(1,2)} - (\hat{\Delta} + \mu - 2\gamma)^{(0,0)} \phi_0^{(1,2)} &= J_3^{(1,2)} \\ (\hat{D} - \tilde{\rho})^{(0,0)} \phi_2^{(1,2)} - (\hat{\delta}^* + 2\pi)^{(0,0)} \phi_1^{(1,2)} &= J_4^{(1,2)} \\ (\hat{\delta} - \tau + 2\beta)^{(0,0)} \phi_2^{(1,2)} - (\hat{\Delta} + 2\mu)^{(0,0)} \phi_1^{(1,2)} &= J_2^{(1,2)}. \end{aligned} \quad (3.55)$$

The currents on the R.H.S.s are given by

$$\begin{aligned} J_1^{(1,2)} &= -i \left(\frac{K}{\Delta} \right)^{(0,2)} \phi_1^{(1,0)} + \frac{a}{2} [(\phi_1 + \phi_1^*)^{(1,0)} \partial_r - (\phi_0 + \phi_0^*)^{(1,0)} \partial_\theta] \phi^{(0,2)} \\ J_2^{(1,2)} &= \left(i \frac{K}{\Delta} + 2\mu \right)^{(0,2)} \phi_1^{(1,0)} + \frac{a}{2} [(\phi_1 + \phi_1^*)^{(1,0)} \partial_\theta + (\phi_0^* - \phi_2)^{(1,0)} \partial_r] \phi^{(0,2)} \\ J_3^{(1,2)} &= \left(i \frac{K}{\Delta} + 3\mu \right)^{(0,2)} \phi_0^{(1,0)} + \frac{a}{2} [(\phi_1 - \phi_1^*)^{(1,0)} \partial_r - (\phi_2 + \phi_2^*)^{(1,0)} \partial_\theta] \phi^{(0,2)} \\ J_4^{(1,2)} &= -i \left(\frac{K}{\Delta} \right)^{(0,2)} \phi_2^{(1,0)} + \frac{a}{2} [(\phi_1 - \phi_1^*)^{(1,0)} \partial_\theta - (\phi_0 - \phi_2^*)^{(1,0)} \partial_r] \phi^{(0,2)}, \end{aligned} \quad (3.56)$$

from which one concludes that the perturbations $\phi_i^{(1,2)}$ couple to the dilaton background through free Maxwell waves ($\phi_i^{(1,0)}$).

However, the full consistency of the argument above requires that the same condition in Eq. (3.54) be applied *simultaneously* to all wave equations at order $(1, 2)$. Thus one obtains constraints from both the gravitational and dilaton wave equations at order $(1, 2)$. We leave this analysis to a future investigation.

IV. ASYMPTOTIC EXPANSIONS

The equations we obtained in Section III C for the Maxwell waves at order $(1, 1)$ look almost intractable, due to the complexity of the source terms. For this reason we consider the equations in the limits $r \rightarrow \infty$ and $r \sim r_+$, and study the asymptotic regimes of ingoing and outgoing modes.

A. Ingoing modes at large radius

We first assume that the ingoing modes of the Maxwell field W_0^{in} and W_2^{in} can be approximated for large r as

$$\begin{aligned}\Delta_0 W_0^{in} &\simeq A_+^{in} Z_+^{in} \frac{e^{i\bar{\omega}r}}{r^{n_+}} \\ W_2^{in} &\simeq A_-^{in} Z_-^{in} \frac{e^{i\bar{\omega}r}}{r^{n_-}} ,\end{aligned}\tag{4.1}$$

where $A_{\pm}^{in} = a_{\pm}^{in} + i b_{\pm}^{in}$ are (complex) coefficients, $Z_{\pm}^{in} = Z_{\pm}^{in}(\theta; m, \bar{\omega})$ are angular functions and n_{\pm} are integers, all to be determined later. The L.H.S.s of Maxwell equations in Eq. (3.48) become

$$\begin{aligned}[\Delta_0 \mathcal{D}_0 \mathcal{D}_0^{\dagger} + \mathcal{L}_0^{\dagger} \mathcal{L}_1 - 2i\bar{\omega}\bar{\rho}] \Delta_0 W_0^{in} &= A_+^{in} \frac{e^{i\bar{\omega}r}}{r^{n_+-1}} \left\{ 2\bar{\omega} [2\bar{\omega}M - i(1+n_+)] Z_+^{in} + \mathcal{O}\left(\frac{1}{r}\right) \right\} \\ [\Delta_0 \mathcal{D}_0^{\dagger} \mathcal{D}_0 + \mathcal{L}_0 \mathcal{L}_1^{\dagger} + 2i\bar{\omega}\bar{\rho}] W_2^{in} &= A_-^{in} \frac{e^{i\bar{\omega}r}}{r^{n_+-1}} \left\{ 2\bar{\omega} [2\bar{\omega}M + i(1-n_-)] Z_-^{in} + \mathcal{O}\left(\frac{1}{r}\right) \right\} .\end{aligned}\tag{4.2}$$

Then we use for the dilaton field the asymptotic (ingoing) form [15],

$$\Phi^{in} = \phi^{(1,0)}(r, \theta) = F^{in} S \frac{e^{i\bar{\omega}r}}{r} ,\tag{4.3}$$

where F^{in} is a real coefficient, $S = S_l^m(\theta; \bar{\omega})$ is a spheroidal wave function (of spin zero), and we find

$$\begin{aligned}\Delta_0 T_+ &\simeq \frac{\sqrt{2}}{2} a \frac{k_d}{k_{EM}} F^{in} \bar{\omega} (\partial_{\theta} - \tilde{Q}) S e^{i\bar{\omega}r} \\ \Delta_0 T_- &\simeq \frac{\sqrt{2}}{4} a \frac{k_d}{k_{EM}} F^{in} (i - 2\bar{\omega}M) (\partial_{\theta} + \tilde{Q}) S \frac{e^{i\bar{\omega}r}}{r} .\end{aligned}\tag{4.4}$$

On equating the R.H.S.s in Eq. (4.2) and the expressions for the currents, Eq. (4.4), one gets $n_+ = 1$ and $n_- = 2$. Further, the coefficients A_{\pm}^{in} and the function S_{\pm}^{in} must satisfy the following equations

$$\begin{aligned}A_+^{in} \left\{ 4\bar{\omega} (\omega M - i) Z_+^{in} + \mathcal{O}\left(\frac{1}{r}\right) \right\} &= \frac{\sqrt{2}}{2} a \frac{k_d}{k_{EM}} F^{in} \left\{ \bar{\omega} (\partial_{\theta} - \tilde{Q}) S + \mathcal{O}\left(\frac{1}{r}\right) \right\} \\ A_-^{in} \left\{ 2\bar{\omega} (2\omega M - i) Z_-^{in} + \mathcal{O}\left(\frac{1}{r}\right) \right\} &= \frac{\sqrt{2}}{4} a \frac{k_d}{k_{EM}} F^{in} \left\{ (i - 2\bar{\omega}M) (\partial_{\theta} + \tilde{Q}) S + \mathcal{O}\left(\frac{1}{r}\right) \right\} .\end{aligned}\tag{4.5}$$

Thus one obtains the solutions

$$\begin{cases} n_+ = 1 \\ Z_+^{in} = (\partial_{\theta} - \tilde{Q}) S \\ a_+^{in} = \frac{\sqrt{2}}{8} a \frac{k_d}{k_{EM}} F^{in} \frac{\bar{\omega}M}{1 + \bar{\omega}^2 M^2} \\ b_+^{in} = \frac{\sqrt{2}}{8} a \frac{k_d}{k_{EM}} F^{in} \frac{1}{1 + \bar{\omega}^2 M^2} , \end{cases}\tag{4.6}$$

and

$$\begin{cases} n_- = 2 \\ Z_-^{in} = (\partial_\theta + \tilde{Q}) S \\ a_-^{in} = -\frac{\sqrt{2}}{8} a \frac{k_d}{k_{EM}} F^{in} \frac{1}{\bar{\omega}} \\ b_-^{in} = 0 . \end{cases} \quad (4.7)$$

From the form of a_-^{in} one clearly sees that this asymptotic solution for the ingoing modes introduces some singular behaviour in the infrared region $\bar{\omega} \rightarrow 0$. Actually, in the limit $\bar{\omega} \rightarrow 0$ the full Eqs. (3.48) are well behaved with respect to $\bar{\omega}$ and thus we expect it to disappear after summation of all order in the large r expansion.

However, for the purpose of comparing the particular (dilaton dependent) solutions $\phi_i^{(1,1)}$ to the free waves $\phi_i^{(1,0)}$ in the black hole background, it is sufficient to consider those wavelengths which are short enough to probe the presence of the black hole itself. This provides us with a natural wavelength cut-off $\bar{\lambda} \sim r_+$ and makes the expressions in Eqs. (4.6) and (4.7) sufficiently well-behaved for $\bar{\omega} > 1/\bar{\lambda}$.

B. Outgoing modes at large radius

The analysis of outgoing modes can be greatly simplified once one notices that Eq. (3.48) can be written

$$\mathcal{G}_\pm(\bar{\omega}, m) f_\pm = \Delta_0 \mathcal{T}_\pm(\bar{\omega}, m) \Phi , \quad (4.8)$$

where $f_+ = \Delta_0 W_0$ and $f_- = W_2$. The differential operators \mathcal{G}_\pm and \mathcal{T}_\pm share the following symmetry,

$$\begin{aligned} \mathcal{G}_+(-\bar{\omega}, -m) &= \mathcal{G}_-(\bar{\omega}, m) \\ \mathcal{T}_+(-\bar{\omega}, -m) &= \mathcal{T}_-(\bar{\omega}, m) . \end{aligned} \quad (4.9)$$

On the other side, the transformation $\bar{\omega} \rightarrow -\bar{\omega}$ (together with the relabelling $F^{in} \rightarrow F^{out}$) maps *ingoing* modes to *outgoing* modes, and vice versa. Thus, it is sufficient to study one set of modes and then apply the mapping above to obtain the solutions for the other set.

For the dilaton field we have the asymptotic (outgoing) form [15],

$$\Phi^{out} = F^{out} S \frac{e^{-i\bar{\omega}r}}{r} , \quad (4.10)$$

where S is a spheroidal wave function and F^{out} an arbitrary coefficient. Since we already know the ingoing solutions from the previous Subsection, we can take full advantage of the symmetry shown above and write

$$\begin{aligned} \Delta_0 W_0^{out} &\simeq A_+^{out} Z_+^{out} \frac{e^{-i\bar{\omega}r}}{r^2} \\ W_2^{out} &\simeq A_-^{out} Z_-^{out} \frac{e^{-i\bar{\omega}r}}{r} . \end{aligned} \quad (4.11)$$

Since $\tilde{Q}(-\bar{\omega}, -m) = -\tilde{Q}(\bar{\omega}, m)$, the angular parts are now given by

$$\begin{aligned} Z_+^{out} &= (\partial_\theta - \tilde{Q}) S \\ Z_-^{out} &= (\partial_\theta + \tilde{Q}) S , \end{aligned} \quad (4.12)$$

and the coefficients $A_\pm^{out} = a_\pm^{out} + i b_\pm^{out}$ are given by

$$\begin{cases} a_\pm^{out} = -a_\mp^{in} \\ b_\pm^{out} = b_\mp^{in} . \end{cases} \quad (4.13)$$

These coefficients again display the same infrared behaviour as A_\pm^{in} in the leading order of the large r expansion. The discussion on a natural cut-off, as explained in the previous Subsection, however is equally applicable for the present outgoing modes.

C. Ingoing and outgoing modes close to the horizon

From the classical analysis of wave propagation close to the horizon, one usually obtains special boundary conditions on the surface $r = r_+$. Namely, only ingoing modes (as seen by any local observer) are allowed [15,11] and there are no non-special outgoing modes available for the dilaton waves on the horizon. Thus one could infer that the currents T_\pm^{out} are identically zero and $W_0^{out} = W_2^{out} \equiv 0$ on the horizon.

However, one can think of a dynamical situation in which some of the parameters of the black hole are changed and generate gravitational and dilatonic waves whose intensity is higher near (and outside) the horizon. The dilaton waves can result in a source for electromagnetic perturbations which is sufficiently strong at least in a region $r \sim r_+$. The outgoing modes then propagate freely once they exit this region (see next Section and Eq. (5.3)). The ingoing modes instead can be detected away from the horizon if they satisfy the superradiant condition, $\bar{\omega} < m\omega_+$, where $\omega_+ = \alpha/(2Mr_+)$ is the superradiant frequency.

A useful radial parameterization of the external region ($r_+ < r < +\infty$) is given by the *turtle* coordinate [15,11],

$$r_* \simeq \frac{2Mr_+}{r_+ - r_-} \ln(r - r_+) , \quad r \sim r_+ , \quad (4.14)$$

We thus find it convenient to introduce the new radial variable

$$x \equiv r - r_+ , \quad (4.15)$$

and then to perform an expansion of Eq. (3.48) at leading order in $1/x$.

We have for the dilaton field the following ingoing form [15],

$$\Phi^{in} = F^{in} S e^{i\tilde{k}r_*} \simeq F^{in} S x^{i\Gamma} , \quad (4.16)$$

where S is again a spheroidal wave function, and

$$\begin{aligned} \tilde{k} &= \bar{\omega} - m\omega_+ \\ \Gamma &= \frac{2Mr_+\bar{\omega} - m\alpha}{r_+ - r_-} . \end{aligned} \quad (4.17)$$

We assume that the Maxwell waves are given by

$$\begin{aligned}\Delta_0 W_0^{in} &= Z_+^{in} x^{i\Gamma+n_+} \\ W_2^{in} &= Z_-^{in} x^{i\Gamma+n_-} ,\end{aligned}\tag{4.18}$$

and we determine the integers n_\pm and the complex functions Z_\pm^{in} by equating the leading orders in $1/x$ of both sides of Eq. (3.48).

One easily finds $n_\pm = 1$, but the forms of S_\pm^{in} turn out to be inconveniently long and we write the solutions symbolically as

$$\begin{aligned}W_0^{in} &= a \frac{k_d}{k_{EM}} \bar{\omega} A_+^{in} e^{i\tilde{k}r_*} \\ W_2^{in} &= a \frac{k_d}{k_{EM}} A_-^{in} \Delta_0 e^{i\tilde{k}r_*} ,\end{aligned}\tag{4.19}$$

where $A_\pm^{in} = A_\pm^{in}(\theta; M, m, \bar{\omega})$ are complex rational functions such that $\lim_{\bar{\omega} \rightarrow 0} A_\pm^{in} \neq 0$. Thus it turns out that W_0 is special on the horizon [15], in that a well-behaved observer on the horizon would see it vanish identically, while W_2 is not.

The outgoing solutions are now easily obtained. We assume for the dilaton field the outgoing form,

$$\Phi^{out} = F^{in} S e^{-i\tilde{k}r_*} ,\tag{4.20}$$

and employ the symmetry between ingoing and outgoing modes described in the previous Subsection. The latter are then given by

$$\begin{aligned}W_0^{out} &= a \frac{k_d}{k_{EM}} A_+^{in}(-\bar{\omega}, -m) e^{-i\tilde{k}r_*} \\ W_2^{out} &= a \frac{k_d}{k_{EM}} \bar{\omega} A_-^{in}(-\bar{\omega}, -m) \Delta_0 e^{-i\tilde{k}r_*} ,\end{aligned}\tag{4.21}$$

and again W_0 is special while W_2 is not.

V. WAVE DETECTION

The outgoing waves at $r \rightarrow \infty$ are the ones that might possibly be detected, thus they carry most of the information one can try to retrieve from an experimental observation. It is thus only with these modes that we deal here.

First we observe that the solutions we get in Section IV are inherently related to the existence of dilaton waves (through the coefficients F^{out}), which in turn can be interpreted as fluctuations in the gravitational-electromagnetic coupling constant (see the form of the action in Eq. (1.1)). These fluctuations are likely to be negligible in the very asymptotic region ($r \rightarrow \infty$). Therefore, one does not expect them to be any appreciable source of electromagnetic waves in the region where a detector can be possibly located.

This is consistent with our analysis at large r as given in section IV. In fact the free outgoing waves at order $(1, 0)$ for $r \rightarrow \infty$ are [15]

$$\begin{aligned}\phi_0^{(1,0)} &\sim \frac{e^{-i\bar{\omega}r}}{r^3} \\ \phi_2^{(1,0)} &\sim \frac{e^{-i\bar{\omega}r}}{r},\end{aligned}\tag{5.1}$$

and the dominant contribution comes from $\phi_2^{(1,0)}$. Our outgoing solutions fall off more rapidly,

$$r\phi_0^{(1,1)} \sim \phi_2^{(1,1)} \sim e^{-i\bar{\omega}r}/r^3.\tag{5.2}$$

This implies that, when the interaction between dilaton waves and the background Maxwell field of the black hole takes place at large r , it produces electromagnetic waves which are negligible when compared to free waves.

However, dilatonic waves might be a strong source in a region $\mathcal{R} \subseteq (r_+, r_d)$ just outside the horizon (see Figure), where they could be generated by some violent dynamical process involving changes in the mass, charge and angular momentum of the black hole. For instance, one can think of the injection of new matter from the accreting disk or a companion star, therefore r_d would be of the order of the typical size of such a structure. Also the background electromagnetic field of the black hole is stronger near r_+ and the local static fields might be even stronger if one takes into account possible ionization and polarization of the surrounding matter. This would further enhance the production in a manner dependent on the charge distribution inside \mathcal{R} .

Any Maxwell waves which cross the region \mathcal{R} interact with all the other fields before they reach $r \rightarrow \infty$. However, since we only consider corrections at order $(1, 1)$ for which the waves $\phi_i^{(1,1)}$ decouple from the other fields (except the background metric) outside \mathcal{R} , one has that, once produced at any point inside \mathcal{R} , $\phi_i^{(1,1)}$ simply propagate freely (outside \mathcal{R}) to $r \rightarrow \infty$.

The way we formalize this model is as follows. Inside \mathcal{R} , we regard $\phi_i^{(1,1)}$ to be given by particular outgoing solutions of the inhomogeneous equations in Eq. (3.48) with boundary conditions for the dilaton waves such that the sources T_{\pm} are appreciably strong. Then, in the region $r > r_d$, we use these solutions as initial conditions at $r = r_d$ for the amplitudes of the fields $\phi_i^{(1,1)}$ which are now required to solve the same Eq. (3.48) but with $T_{\pm} \simeq 0$. When $T_{\pm} = 0$, the equations in Eq. (3.48) reduce to the homogeneous equations given in Eq. (3.28) at order $(1, 0)$ and whose solutions are exactly the free waves with asymptotic behaviour displayed above, Eq. (5.1). Thus one has

$$\begin{cases} \phi_0^{(1,1)}(r > r_d) \simeq |\phi_0^{(1,1)}(r_d)| \frac{r_d^3}{r^3} e^{-i\bar{\omega}r} \\ \phi_2^{(1,1)}(r > r_d) \simeq |\phi_2^{(1,1)}(r_d)| \frac{r_d}{r} e^{-i\bar{\omega}r}, \end{cases}\tag{5.3}$$

where $\phi_i^{(1,1)}(r_d)$ are the values taken at $r = r_d$ by the solutions of the inhomogeneous equations inside \mathcal{R} and the matching conditions at $r = r_d$ have been applied as stated above. It also follows that, for $r \rightarrow \infty$, $\phi_0^{(1,1)} \sim \phi_0^{(1,0)}$ is again negligible with respect to $\phi_2^{(1,0)}$, but $\phi_2^{(1,1)} \sim \phi_2^{(1,0)}$ is not.

If r_d is relatively big (compared to r_+) one can still assume the dilaton inside \mathcal{R} to be of the form given in Eq. (4.10) and obtains the corresponding electromagnetic perturbations in Eq. (4.11). However since \mathcal{R} is likely to extend down to r_+ , we found it interesting to study Maxwell equations in a region very close to the horizon (see Subsection IV C) In that case, the relevant form of the dilatonic waves is given in Eq. (4.20), and $\phi_2^{(1,1)}(r_d)$ is obtained from Eq. (4.21). Of course a complete treatment would contain contributions from both the approximations.

For this model one can easily compute the energy flux at infinity. In order to do this, we first observe that

$$\begin{aligned}\phi_0 &= \frac{\sqrt{2}}{2} \left[\mathcal{E}_\theta \frac{r^2 + \alpha^2}{\Delta \bar{\rho}} + \mathcal{B}_\phi \frac{1}{\bar{\rho}} + i \left(\mathcal{E}_\phi \frac{\bar{\rho}^*}{\Delta \sin \theta} - \mathcal{B}_\theta \frac{1}{\bar{\rho} \sin \theta} \right) \right] \\ \phi_2 &= \frac{\sqrt{2}}{2} \left[\mathcal{B}_\phi \frac{\Delta}{\bar{\rho}^* \rho^2} - \mathcal{E}_\theta \frac{r^2 + \alpha^2}{\bar{\rho}^* \rho^2} + i \left(\mathcal{E}_\phi \frac{1}{\bar{\rho}^* \sin \theta} + \mathcal{B}_\theta \frac{\Delta}{\bar{\rho}^* \rho^2 \sin \theta} \right) \right],\end{aligned}\quad (5.4)$$

which, in the limit $r \rightarrow \infty$ and for $\phi_0^{(1,0)}$, $\phi_2^{(1,0)}$ given in Eq. (5.1) above, imply that the coordinate components \mathcal{E}_i , \mathcal{B}_i are of order r^0 and the local components of the electromagnetic waves are given by

$$\begin{aligned}\mathcal{E}_{\hat{\theta}}^{(1,0)} &= -\mathcal{B}_{\hat{\phi}}^{(1,0)} = -\frac{\sqrt{2}}{2} (\phi_2 + \phi_2^*)^{(1,0)} \\ \mathcal{E}_{\hat{\phi}}^{(1,0)} &= \mathcal{B}_{\hat{\theta}}^{(1,0)} = \frac{\sqrt{2}}{2i} (\phi_2 - \phi_2^*)^{(1,0)}.\end{aligned}\quad (5.5)$$

The (outgoing) flux per unit solid angle at infinity is thus given by

$$\frac{d^2 E}{dt d\Omega} = \lim_{r \rightarrow \infty} \frac{r^2}{2\pi} |\phi_2^{(1,0)}|^2 \quad (5.6)$$

On using the solution in Eq. (5.3) one obtains new contributions to the electric and magnetic fields,

$$\begin{aligned}\mathcal{E}_{\hat{\theta}}^{(1,1)} &= -\mathcal{B}_{\hat{\phi}}^{(1,1)} \simeq -\frac{\sqrt{2}}{2} \frac{a_-^{out}}{r_d^3} |Z_-^{out}| \\ \mathcal{E}_{\hat{\phi}}^{(1,1)} &= \mathcal{B}_{\hat{\theta}}^{(1,1)} \simeq \frac{\sqrt{2}}{2} \frac{b_-^{out}}{r_d^3} |Z_-^{out}|, \quad (r = r_d),\end{aligned}\quad (5.7)$$

where we assumed the asymptotic solution in Eq. (4.11). These imply a correction to the energy flux,

$$\frac{d^2 E}{dt d\Omega} \simeq \lim_{r \rightarrow \infty} \frac{r^2}{2\pi} \left| \phi_2^{(1,0)} + e^{i\gamma} Q \phi_2^{(1,1)} \right|^2, \quad (5.8)$$

where the relative phase γ between the (1,0) term and the (1,1) term has been written explicitly. Since the phase of the solution in Eq. (5.3) is determined only with respect to the phase of the source (dilatonic wave) and the latter is unrelated to $\phi_2^{(1,0)}$, the phase γ is

essentially arbitrary. We can also assume that dilatonic waves exist with all possible relative phases and average over $\gamma \in (0, 2\pi)$. One then obtains

$$\begin{aligned} \frac{d^2 E}{dt d\Omega} &\simeq \lim_{r \rightarrow \infty} \frac{r^2}{2\pi} \int_0^{2\pi} \frac{d\gamma}{2\pi} \left| \phi_2^{(1,0)} + e^{i\gamma} Q \phi_2^{(1,1)} \right|^2 \\ &\simeq \lim_{r \rightarrow \infty} \frac{r^2}{2\pi} |\phi_2^{(1,0)}|^2 + \frac{a^2 Q^2}{128 \pi^2 r_d^4} \frac{k_d}{k_{EM}} \frac{(F^{out})^2}{1 + \bar{\omega}^2 M^2} |Z_-^{out}|^2, \end{aligned} \quad (5.9)$$

or an analogous expression when the solution for $\phi_2^{(1,1)}$ inside \mathcal{R} is of the form in Eq. (4.21). The main feature of the expression above is that the correction that we get contains information about the charge of the black hole and the angular structure of the dilatonic perturbations close to the horizon, the latter being encoded in the angular function Z_- . Of course, the contribution we show here is for just one frequency $\bar{\omega} > 1/\bar{\lambda}$ ($\bar{\lambda}$ is the cut-off discussed in Section IV). The general case is a superposition of all modes.

VI. CONCLUSIONS

The double perturbative expansion (Eq.(3.1)) provides a means of calculating the electromagnetic, dilaton and gravitational fields to any desired accuracy. Of course, beyond the lowest orders the field equations must be solved numerically due to the increasing complexity of the equations with increasing order of the expansion. Even at the lowest order we were able to find explicit expressions for the corrections to the electromagnetic waves only in the asymptotic limits $r \rightarrow \infty$ and $r \sim r_+$. We also obtained an explicit form for the correction to the energy flux due to the presence of a scalar gravitational field (the dilaton). Measurement of this flux could in principle be used to test the idea that black holes have a scalar component if the flux from a source could be measured for two cases – one case being when the waves from the source are far away from a black hole and the other being when the waves pass near to the horizon of a black hole. An occultation observation would be a suitable event to test the idea that black hole gravitational fields have a scalar component.

In this paper we have calculated the lowest order corrections to the electromagnetic waves propagating on a black hole background possessing a scalar gravity component. A future project will be to determine the corresponding corrections for the gravitational waves propagating on the same background. The dilaton corrections to these waves first appear at order (1, 2).

ACKNOWLEDGMENTS

This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-96ER40967.

REFERENCES

- [1] B. Harms and Y. Leblanc, Phys. Rev. D **46**, 2334(1992); D **47**, 2438 (1993).
- [2] B. Harms and Y. Leblanc, Ann. Phys. **244**, 262(1995); **244**, 272 (1995).
- [3] P.H. Cox, B. Harms and Y. Leblanc, Europhys. Letts. **26**, 321 (1994).
- [4] B. Harms and Y. Leblanc, Europhys. Letts. **27**, 557 (1994).
- [5] B. Harms and Y. Leblanc, Ann. Phys. **242**, 265 (1995).
- [6] B. Harms and Y. Leblanc, *Proceedings of the Texas/PASCOS Conference, 92. Relativistic Astrophysics and Particle Cosmology*, eds. C.W. Ackerlof and M.A. Srednicki, Annals of the New York Academy of Sciences **688**, 454 (1993).
- [7] B. Harms and Y. Leblanc, *Supersymmetry and Unification of Fundamental Interactions*, ed. Pran Nath, World Scientific (1994) p.337.
- [8] B. Harms and Y. Leblanc, *Banff/CAP Workshop on Thermal Field Theory*, eds. F.C. Khanna, R. Kobes, G. Kunstatter and H. Umezawa, World Scientific (1994), p.387.
- [9] R. Casadio, B. Harms, Y. Leblanc and P.H. Cox, Phys. Rev. D **55**, 814 (1997).
- [10] The ratio Q/M for physical particles can be estimated using M.K.S. units once one reintroduces the conventional values of G and c . One then has $Q/M = \bar{Q}/(\bar{M} \sqrt{4\pi\epsilon_0 G})$ which is approximately equal to 10^{42} for an electron ($\bar{Q} \simeq 10^{-19}$ C and $\bar{M} \simeq 10^{-30}$ Kg) and to 10^{-2} for a particle of the same charge but of Planck mass. This implies that our approximation (Q/M small) does not apply to most known elementary particles, but can be valid for black holes produced during the gravitational collapse of massive stars and also for cosmological black holes of Planck mass size. On the other hand, the opposite approximation (Q/M large) corresponds to a naked singularity.
- [11] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, Oxford University Press, Oxford (1983).
- [12] C.W. Misner, K.S Thorne and J.A. Wheeler, *Gravitation*, W.H. Freeman and Co., San Francisco, 1973.
- [13] N. Straumann, *General Relativity and Relativistic Astrophysics*, Springer-Verlag, Berlin(1984).
- [14] E. Newman and R. Penrose, J. Math. Phys. **6**, 566 (1964).
- [15] S.A. Teukolsky, Astroph. J. **185**, 635 (1973).
- [16] C. Flammer, *Spheroidal Wave Functions*, Stanford University Press, Stanford (1957).

Figure caption

The shaded region represents the interior of the black hole ($r \leq r_+$). The region between r_+ and r_d is the one denoted by \mathcal{R} in the text and corresponds to the domain where sufficiently strong dilatonic waves interact with the background electromagnetic fields to produce Maxwell (outgoing) radiation. For $r > r_d$ the latter propagates freely toward the observer ($r \rightarrow \infty$).

