

COALESCING BINARIES AND DOPPLER EXPERIMENTS

A. Vecchio

*Max Planck Institut für Gravitationsphysik, Albert-Einstein-Institut
14473 Potsdam, Germany*

B. Bertotti

*Dipartimento di Fisica Nucleare e Teorica, Università di Pavia
27100 Pavia, Italy*

L. Iess

*Dipartimento di Ingegneria Aerospaziale, Università "La Sapienza"
00158 Roma, Italy*

We discuss the sensitivity of the CASSINI experiments to gravitational waves emitted by the in-spiral of compact binaries. We show that the maximum distance reachable by the instrument is ~ 100 Mpc. In particular, CASSINI can detect massive black hole binaries with chirp mass $\gtrsim 10^6 M_\odot$ in the Virgo Cluster with signal-to-noise ratio between 5 and 30 and possible compact objects of mass $\gtrsim 30 M_\odot$ orbiting the massive black hole that our Galactic Centre is likely to harbour.

The Doppler tracking¹ of interplanetary spacecraft is the only technique presently available to search for gravitational waves (GW) in the low frequency regime. During the past decade, a number of experiments, lasting from some day to a month, have been carried out, the most recent involving the spacecraft GALILEO, MARS-OBSERVER and ULYSSES^{2,3}. The CASSINI probe, due to be launched on October 6th 1997 and equipped with an improved radio link (in X and K_a band, 8.4 and 34 GHz, respectively), will perform three 40-days data acquisition runs between 2000 and 2004. Here we discuss the expected sensitivity of these experiments to GWs emitted during the in-spiral of binaries of compact objects.

In a Doppler experiment a GW of amplitude h produces at the detector output a signal $\tilde{s}(f)$, that, in the convenient Fourier domain, reads¹

$$\tilde{s}(f) = \tilde{r}_\theta(f; T) \tilde{h}(f), \quad (1)$$

where $\tilde{r}_\theta(f; T)$ is the characteristic *three-pulses response*, which depends on T , the round-trip light-time of the radio link out to the distance of the probe, and on θ , the angle between the spacecraft and the source. Focusing attention on the radiation emitted during the in-spiral of a binary toward its final

coalescence, \tilde{h} reads⁵

$$\tilde{h}(f) = \left(\frac{5}{96\pi^{4/3}} \right)^{1/2} Q \frac{\mathcal{M}^{5/6}}{D} f^{-7/6} e^{i\Psi(f)}, \quad (2)$$

where \mathcal{M} is the chirp mass of the binary, D the distance and Q a function of the orientation of the source and the GW polarization; $\Psi(f)$ is the GW phase. Eq. (2) is valid up to some frequency f_{isco} , after which the final coalescence sets off with emission of a burst, whose structure is still quite unknown; indeed, we cut-off the signal (somehow arbitrarily) at $f = f_{\text{isco}}$.

The signal-to-noise ratio (SNR) ρ measured at the output of the optimal matched filter reads:

$$\rho^2 = 4 \int_{f_b}^{f_e} \frac{|\tilde{s}(f)|^2}{S_n(f)} df = \left(\frac{5Q^2}{24\pi^{4/3}} \right) \frac{\mathcal{M}^{5/3}}{D^2} \int_{f_b}^{f_e} \frac{f^{-7/3} |\tilde{r}_\theta(f)|^2}{S_n(f)} df, \quad (3)$$

where $S_n(f)$ is the noise spectral density of the detector; f_b and $f_e = \min[f_b(1 - T_1/t_n)^{-3/8}, f_{\text{isco}}]$ are the instantaneous frequencies of $h(t)$ at the beginning ($t = 0$, for convention) and at the end ($t = T_1$) of the data set; here T_1 is the time of observation and t_n the time to coalescence from f_b (in general $t_n \gg T_1$). In the ideal case in which we neglect the finite bandwidth of the signal and of the instrument ($f_b \rightarrow 0$, $f_e \rightarrow \infty$) and the different frequency response due to the structure of the noise and of the three-pulses filter ($S_n = \text{const.}$, $\theta = \pi/2$), we can derive an *ideal* SNR, ρ_{id} , that depends only on fundamental parameters of the source and the detector; for CASSINI typical values it reads

$$\rho_{\text{id}} \simeq 3 \left(\frac{D}{100 \text{ Mpc}} \right)^{-1} \left(\frac{\mathcal{M}}{10^7 M_\odot} \right)^{5/6} \left(\frac{T}{10^4 \text{ sec}} \right)^{2/3} \left(\frac{S_n}{9 \times 10^{-26} \text{ Hz}^{-1}} \right)^{-1/2}. \quad (4)$$

This result, although obtained in a simplified (and optimistic) case, clearly shows that the instrument is sensitive to binaries out to the distance of the Virgo Cluster ($D \simeq 17 \text{ Mpc}$) and beyond. Of course the reference value (4) is not attained in a real experiment, due to the limited bandwidth (f_b, f_e), the strong oscillations of $\tilde{r}_\theta(f; T)$, that, depending on θ and (f_b, f_e) can suppress the signal at the detector output, and the increase of $S_n(f)$ at low frequencies. We can rigorously describe the degradation of SNR, with respect to ρ_{id} , introducing the *detector efficiency function*

$$\Upsilon(f_b T, f_e T, \theta) \equiv \frac{\rho}{\rho_{\text{id}}} \leq 1. \quad (5)$$

Its main features are the following⁵: (i) Υ is peaked at $f \sim 1/T$; as a consequence, for f_b larger than a few $1/T$, Υ is severely depressed; searches for signals in past Doppler data³ have always concentrate on the band $f > 1/T$, therefore missing the most sensitive portion of the observable spectrum; (ii) only broad band searches (say $f_e \gtrsim 2 f_b$) of chirping signals can produce $\Upsilon \rightarrow 1$; in order to reach distant sources, one can confine the analysis to binaries with mass $\mathcal{M} \gtrsim 3.3 \times 10^7 (f_b/10^{-5} \text{ Hz})^{-8/5} (T_1/40 \text{ days})^{-3/5} M_\odot$; (iii) the optimal direction of observation θ , for given \mathcal{M} and D , depends on the frequency band (f_b, f_e) swept by the signal.

We apply now the previous results to the CASSINI experiments for two specific astrophysical targets: the Virgo Cluster and our Galactic Centre. In both cases the sensitivity of the three data sets turns out to be quite comparable and we discuss only the second one. The results are summarized in Fig. 1.

For the Virgo Cluster, binaries with $\mathcal{M} \gtrsim 2 \times 10^6 M_\odot$ and $f_b \lesssim 7 \times 10^{-5} \text{ Hz}$ can be detected at $\text{SNR} \geq 5$, and in the region of the plane (\mathcal{M}, f_b) around $(10^{-5} \text{ Hz}, 5 \times 10^7 M_\odot)$ signals produce $\text{SNR} \simeq 30$. These results hold for systems of comparable mass; decreasing the mass ratio m_2/m_1 , the sensitivity region in the plane (\mathcal{M}, f_b) shrinks and for $m_2/m_1 < 0.01$ the instrument is not able to detect any source (at $D = 17 \text{ Mpc}$). The detector can therefore reach, at $\text{SNR} = 5$, a distance $\simeq 100 \text{ Mpc}$ for $m_1 \sim m_2 \sim 10^8 M_\odot$. According to current estimates of the event rate of super-massive black hole binary coalescence⁴, we can conclude that even with this new generation of Doppler instruments the chance of detection is probably low; however (i) when accurate templates for the signals emitted during the final black hole merger will be available, the search depth can considerably increase; (ii) the results reported in Fig. 1 have been obtained using the conservative estimate $S_n \propto f^{-2}$ for $f < 10^{-4} \text{ Hz}$; improvements in the data pre-processing could lead to a flatter noise curve at very low frequencies, enhancing the sensitivity in that very crucial band; (iii) T_1 could be extended.

We consider now galactic observations. Based on evidences of a massive black hole in our Galactic Centre⁶, we investigate the ability of CASSINI of picking up signals emitted by a compact object of mass m_2 orbiting the central black hole $m_1 = 2 \times 10^6 M_\odot$. The instrument turns out to be a rather sensitive galactic monitor; in fact it would be able to detect secondary black holes with $m_2 \gtrsim 30 M_\odot$ at $\text{SNR} > 5$, and for $m_2 \gtrsim 10^3 M_\odot$ with a remarkable $\text{SNR} \sim 100$ or higher (but the detection probability remains low^{5,7}). Furthermore, most of the systems visible during one experiment would be detectable (independently) also in one of the other two data sets, as the time to coalescence for the relevant range of masses and frequencies is $\simeq 2995 (f_b/5 \times 10^{-4} \text{ Hz})^{-8/3} (m_2/10^2 M_\odot)^{-1} (m_1/2 \times 10^6 M_\odot)^{-2/3} \text{ days}$.




Figure 1: Detectable binaries in the Virgo Cluster (solid line; contour plots of $\text{SNR} = 1, 5, 10$ in the plane (\mathcal{M}, f_b) ; $m_1 = m_2$, $\cos \theta \simeq -0.04$, $D = 17 \text{ Mpc}$) and our Galactic Centre (dot-dash line; contour plots of $\text{SNR} = 1, 5, 100$ in the plane (m_2, f_b) for $m_1 = 2 \times 10^6 M_\odot$; $\cos \theta \simeq -0.85$, $D = 8 \text{ kpc}$) during CASSINI second experiment ($T \simeq 6796 \text{ s}$, $T_1 = 40 \text{ days}$).

We can therefore conclude that CASSINI represents a significant improvement with respect to previous Doppler missions³, and although can not guarantee a positive detection of GWs, it is definitely able to provide valuable astrophysical information on massive black holes, which are inaccessible to any generation of Earth-based detectors.

References

1. F. B. Estabrook and H. R. Wahlquist, Gen. Rel. Grav. **6**, 439 (1975)
2. B. Bertotti, R. Ambrosini, J. Armstrong, S.W. Asmar, G. Comoretto, G. Giampieri, Y. Koyama, L. Iess, A. Messeri, A. Vecchio and H.D.

- Wahlquist, *Astron. and Astroph.* **296**, 13, (1995).
3. B. Bertotti, in *Proceedings of the 14th International Conference on General Relativity and Gravitation*, edited by M. Francaviglia, G. Longhi, L. Lusanna and E. Sorace (World Scientific, Singapore, 1997), pp. 79-102.
 4. A. Vecchio, *Class. and Quantum Grav.* **14**, 1431 (1997).
 5. B. Bertotti, A. Vecchio and L. Iess, *Gravitational waves from coalescing binaries and Doppler experiments*, in preparation.
 6. A. Eckart and R. Genzel, *Nature* **383**, 405 (1996).
 7. S. Sigurdsson, *Class. and Quantum Grav.* **14**, 1425 (1997).