

Gravitational field of a stationary circular cosmic string loop

A.A.Sen¹ and N.Banerjee

Relativity and Cosmology Research Center

Department of Physics, Jadavpur University

Calcutta 700032, India

Abstract

Gravitational field of a stationary circular cosmic string loop has been studied in the context of full nonlinear Einstein's theory of gravity. It has been assumed that the radial and tangential stresses of the loop are equal to the energy density of the string loop. An exact solution for the system has been presented which has a singularity at a finite distance from the axis, but is regular for any other distances from the axis of the loop.

PACS NO: 04.20Jb, 98.80.Cq

¹e-mail: anjan@juphys.ernet.in

Phase transition in the early universe might have produced some topological defects [1]. One of these defects, cosmic strings, have attracted a lot of interests amongst cosmologists for various reasons[2]. Cosmic strings are capable of producing observational effects such as the double images of quasars and also they are considered to be possible candidates as seeds for the galaxy formation. Moreover, strings are believed to sustain the cosmological evolutions unlike other defects.

In some gauge models, strings do not have any end, and thus they are either of infinite extent form or closed circular loops. One of the most notable features of the gravitational field of a straight infinite cosmic string is the presence of an "angular deficit" in an otherwise Minkowskian spacetime having a magnitude related to the linear energy density μ of the string by the equation $\delta\phi = 8\pi G\mu$. In fact, this angular deficit plays the key role for the production of the double images of a quasars[3]-[6].

The deficit angle model is widely believed to be a good approximation for describing a spacetime exterior to the string core. Frolov, Israel and Unruh (FIU)[7] used this approximation for studying a closed circular cosmic string loop at a moment of time symmetry. With the help of the initial value formulation[8], they produced a family of momentarily stationary circular cosmic strings, which are considered as thin loops either at the time of formation or at the turning point of expansion and collapse. An important assumption in this work is that all points on the circular string be conical singularities with angular deficit equal to that of a infinite straight string of equal linear energy density. Hughes, McManus, and Vandyck[9] investigated further the problem of angular deficit of circular string. They considered a weak field stationary solutions of Einstein's field equations for a thin circular string and also established that radial stress should be introduced to support the string loop against possible gravitational collapse. They determined the form of the radial stress from the stress energy conservation relations. The main result of their study is that under weak field assumption, circular string produces the same angular deficit as straight infinitely long string having the same linear energy density. In another work, Mc-

Manus and Vandyck[10] considered a string loop with a rotation of the loop which provides the necessary centrifugal reaction partially or fully in order to avoid the possible gravitational collapse.

In the present work, we have assumed the form of the energy momentum tensor for circular string as that proposed by Hughes et.al[9]. But unlike the work of Hughes et.al, we have taken the nonzero components of T_ν^μ to be function of both r and z . With some simplified assumptions, we have been able to produce a set of exact analytical solutions of the full nonlinear Einstein's equations for a circular string. The spacetime, we have obtained, have a singularity at a finite distance from the axis, but is regular for any other distances from the axis of the loop.

A circular cosmic string loop produces an axially symmetric gravitational field, and the most general static metric of this type may be written as [11]

$$ds^2 = -e^{2\nu} dt^2 + e^{2(\xi-\nu)} r^2 d\phi^2 + e^{2(\eta-\nu)} (dr^2 + dz^2) \quad (1)$$

where the three functions ν, ξ, η are functions of r and z and (t, ϕ, r, z) denotes the cylindrical coordinates.

We have taken the form of the energy momentum tensor for the circular string loop as prescribed by Hughes et.al[9]:

$$\begin{aligned} T_t^t &= -\alpha(r, z) \\ T_\phi^\phi &= -\beta(r, z) \\ T_r^r &= -\Delta(r, z) \end{aligned} \quad (2)$$

and all other components of T_ν^μ are zero. For a circular string T_ϕ^ϕ plays the same role as that of the longitudinal stress T_z^z for a straight string. T_r^r is the external radial stress, required to support the loop against collapse.

In order to solve the full nonlinear Einstein's equations ($G_\nu^\mu = 8\pi T_\nu^\mu$) (we have assumed $G = c = 1$) for the system given by (1) and (2), we make an assumption that all the nonzero components of T_ν^μ are equal, i.e.

$$T_t^t = T_\phi^\phi = T_r^r = -\sigma(r, z) \quad (3)$$

With this assumption and with the line element (1), the Einstein's equations become:

$$\ddot{\xi} - 2\ddot{\nu} + \ddot{\eta} + \frac{2}{r}(\dot{\xi} - \dot{\nu}) + (\dot{\xi} - \dot{\nu})^2 + \xi'' - 2\nu'' + \eta'' + (\xi' - \nu')(\xi' - \eta') = 8\pi\sigma e^{2(\eta-\nu)} \quad (4a)$$

$$\ddot{\eta} + \dot{\nu}^2 + \eta'' + \nu'^2 = 8\pi\sigma e^{2(\eta-\nu)} \quad (4b)$$

$$\frac{\dot{\eta}}{r} + \dot{\eta}\dot{\xi} + \dot{\nu}^2 + \xi'' + \xi'^2 + \nu'^2 - \xi'\eta' = 8\pi\sigma e^{2(\eta-\nu)} \quad (4c)$$

$$\ddot{\xi} + \frac{1}{r}(2\dot{\xi} - \dot{\eta}) + \xi^2 + \dot{\nu}^2 - 2\dot{\xi}\dot{\nu} + \xi'\eta' - \nu'^2 = 0 \quad (4d)$$

$$\frac{1}{r}(\xi' - \eta') + \dot{\xi}(\eta' - \xi') + \xi'\dot{\eta} - 2\dot{\nu}\nu' = 0 \quad (4e)$$

In the above equations an overhead dot represents differentiation w.r.t r and prime represents differentiation w.r.t z .

The conservation equations ($T_{\nu;\mu}^\mu = 0$) give the relations

$$\dot{\sigma} + \sigma(\dot{\eta} - \dot{\nu}) = 0 \quad (5)$$

and

$$\sigma(\xi' + \eta' - \nu') = 0 \quad (6)$$

We see immediately from (6) that, in order to have the energy density σ to be nonzero, we should have,

$$\xi' = \nu' - \eta'. \quad (7)$$

Equation (5) can be integrated to give

$$\sigma = C(z)e^{(\nu-\eta)} \quad (8)$$

where $C(z)$ is an arbitrary function of z .

Now we make the simplifying assumption that the metric coefficients are separable functions of their arguments i.e. $\xi = \xi_1(r) + \xi_2(z)$, $\nu = \nu_1(r) + \nu_2(z)$ and $\eta = \eta_1(r) + \eta_2(z)$. Equations (8) then shows that σ will be also separable function of its arguments.

In what follows we will find the possible solutions of the full nonlinear Einstein's equations given by (4a)-(4e) with these assumptions.

Equations (4a)-(4e) can now be written as,

$$\ddot{\xi}_1 - 2\ddot{\nu}_1 + \ddot{\eta}_1 + \frac{2}{r}(\dot{\xi}_1 - \dot{\nu}_1) + (\dot{\xi}_1 - \dot{\nu}_1)^2 + \xi_2'' - 2\nu_2'' + \eta_2'' + (\xi_2' - \nu_2')(\xi_2' - \eta_2') = 8\pi\sigma e^{2(\eta-\nu)} \quad (9a)$$

$$\ddot{\eta}_1 + \dot{\nu}_1^2 + \eta_2'' + \nu_2'^2 = 8\pi\sigma e^{2(\eta-\nu)} \quad (9b)$$

$$\frac{\dot{\eta}_1}{r} + \dot{\eta}_1\dot{\xi}_1 + \dot{\nu}_1^2 + \xi_2'' + \xi_2'^2 + \nu_2'^2 - \xi_2'\eta_2' = 8\pi\sigma e^{2(\eta-\nu)} \quad (9c)$$

$$\ddot{\xi}_1 + \frac{1}{r}(2\dot{\xi}_1 - \dot{\eta}_1) + \dot{\xi}_1^2 + \dot{\nu}_1^2 - \dot{\xi}_1\dot{\eta}_1 + \xi_2'\eta_2' - \nu_2'^2 = 0 \quad (9d)$$

$$\frac{1}{r}(\xi_2' - \eta_2') + \dot{\xi}_1(\eta_2' - \xi_2') + \xi_2'\dot{\eta}_1 - 2\dot{\nu}_1\nu_2' = 0 \quad (9e)$$

As the right hand side of both equations (9a) and (9b) are equal, one can equate the r part and z part of these equations.

By equating r part of (9a) and (9b), one can get after some simple calculations

$$\dot{\xi}_1 + \frac{1}{r} = 2\dot{\nu}_1 \quad (10)$$

With the assumption of separability of the metric coefficients, the z part of equation (7) reads like

$$\xi_2' = \nu_2' - \eta_2' \quad (11)$$

With (11) and (10), one can have from (9e), the relation,

$$\xi_2'(4\nu_1 - \eta_1) = 0 \quad (12)$$

From eqn(12), it is easy to calculate that either $\xi_2' = 0$ or $4\nu_1 = \eta_1$.

In this work, we present the solutions where $\xi_2' = 0$ and $4\nu_1 \neq \eta_1$.

Therefore,

$$e^{2\xi_2} = C_0, \quad (13)$$

C_0 being a constant.

From eqn(11), with the help of (13) one gets,

$$\nu_2' = \eta_2' \quad (14)$$

With eqn(13), eqn(9d) becomes,

$$\ddot{\xi}_1 + \frac{1}{r}(2\dot{\xi}_1 - \dot{\eta}_1) + \dot{\xi}_1^2 + \dot{\nu}_1^2 - \dot{\xi}_1\dot{\eta}_1 = \nu_2'^2 = K \quad (15)$$

where K is the constant of separation.

With (10),(13),(14) and (15), (9a) and (9b) reduce to a single equation,

$$\ddot{\eta}_1 + \dot{\nu}_1^2 + K = 8\pi\sigma e^{2(\eta_1 - \nu_1)} \quad (16)$$

and eqn(9c) becomes

$$\frac{\dot{\eta}_1}{r} + \dot{\eta}_1\dot{\xi}_1 + \dot{\nu}_1^2 + K = 8\pi\sigma e^{2(\eta_1 - \nu_1)}. \quad (17)$$

From the z part of eqn(15) and using (14) we have

$$e^{\eta_2} = e^{\nu_2} = A_0 e^{\sqrt{K}z} \quad (18)$$

where A_0 is an arbitrary constant of integration.

Combining eqn(16) and (17), we get

$$\ddot{\eta}_1 - \frac{\dot{\eta}_1}{r} - \dot{\eta}_1\dot{\xi}_1 = 0$$

which, with the use of eqn(10) becomes

$$\ddot{\eta}_1 - 2\dot{\nu}_1\dot{\eta}_1 = 0, \quad (19)$$

for which $\dot{\eta}_1 = 0$ is a solution which is used in our subsequent calculations. The other case is $\frac{\ddot{\eta}_1}{\dot{\eta}_1} = 2\dot{\nu}_1$, and this is not considered in the present work.

Hence for our case

$$e^{2\eta_1} = \text{constant} = m. \quad (20)$$

Using (20) and (10) in the r part of eqn(15), we get on integration

$$e^{\nu_1} = e^{a(r-r_0)} [1 - e^{-5a(r-r_0)}]^{2/5} \quad (21)$$

where $a^2 = K/5$ and r_0 is an arbitrary integration constant.

Eqn(10) and (21) together yield the solution of ξ_1 as

$$e^{\xi_1} = \frac{1}{r} e^{2a(r-r_0)} [1 - e^{-5a(r-r_0)}]^{4/5}. \quad (22)$$

where a constant of integration has been ignored without any loss of generality.

Hence the complete line element for a circular cosmic string loop with radial and tangential stress equal to the energy density, is given by

$$ds^2 = e^{2a(r-r_0)} [1 - e^{-5a(r-r_0)}]^{4/5} (-A_0^2 e^{2nz} dt^2 + C_0 A_0^{-2} e^{-2nz} d\phi^2) \\ + m e^{-2a(r-r_0)} [1 - e^{-5a(r-r_0)}]^{-4/5} (dr^2 + dz^2) \quad (23)$$

where $n = \sqrt{K}$ and $a^2 = K/5$.

One can calculate the energy density σ from either (16) or (17) which becomes,

$$\sigma = \frac{K}{8\pi m} e^{2a(r-r_0)} [1 - e^{-5a(r-r_0)}]^{4/5} \left[\frac{(e^{-5a(r-r_0)} + 1)^2}{5(e^{-5a(r-r_0)} - 1)^2} + 1 \right] \quad (24)$$

One can see from (24) that σ is regular everywhere except at $r = r_0$ and hence $r = r_0$ is identified to correspond to the radius of the loop.

We have a circular ring at $r = r_0$, $z = 0$ momentarily at rest at time $t = 0$. At this point it is convenient to pass from the coordinates r, z to toroidal coordinates α, ψ defined by

$$r = r_0 N^{-2} \sinh \alpha, z = r_0 N^{-2} \sin \psi \quad (25a)$$

with

$$N(\alpha, \psi) = (\cosh \alpha - \cos \psi)^{1/2} \quad (0 \leq \alpha \leq \infty, -\pi \leq \psi \leq \pi) \quad (25b)$$

We note that the circular loop $r = r_0, z = 0$ corresponds to the limiting torus $\alpha \rightarrow \infty$.

The metric (1) is now transformed to toroidal coordinates as

$$ds^2 = -e^{2\nu} dt^2 + e^{2(\xi-\nu)} r^2 d\phi^2 + r_0^2 N^{-4} e^{2(\eta-\nu)} (d\alpha^2 + d\psi^2) \quad (26)$$

where ν, ξ, η, r are functions of α and ψ .

To check the metric (26) for elementary flatness on the loop i.e. for $\alpha \rightarrow \infty$ [7, 9] we must examine circle $\alpha = \alpha_0$ at constant t, ϕ around the ring and calculate the ratio of the proper perimeter to proper radius in the proper radius tends to zero.

For t, ϕ constant, the metric (26) using (18) and (20) becomes

$$ds^2 = r_0^2 m N^{-4} e^{-2\nu_1} (d\alpha^2 + d\psi^2) \quad (27)$$

To be elementary flat on the loop like Hughes et.al [9] $F = \left| \lim_{\alpha_0 \rightarrow \infty} \frac{\int_{-\pi}^{\pi} d\psi N^{-2} e^{-\nu_1} |_{\alpha=\alpha_0}}{\int_{\alpha_0}^{\infty} N^{-2} e^{-\nu_1} d\alpha} \right|$ should be a constant.

Using (21) and (25) one can calculate that

$$e^{-\nu_1} \rightarrow b_0 e^{4\alpha/5} \quad (28)$$

for $\alpha \rightarrow \infty$ where $b_0 = (-10ar_0)^{-2/5}$.

From (25b)

$$N^{-2} \rightarrow 2e^{-\alpha} \quad (29)$$

for $\alpha \rightarrow \infty$.

Using (28) and (29) one can see that

$$F \rightarrow 2\pi/5 \quad (30)$$

for $\alpha \rightarrow \infty$. And hence one can calculate the angular deficit near the loop to be $d\psi = 8\pi/5$.

In Conclusion we have presented the exact line element for a circular loop in the context of a full set of nonlinear Einstein's theory of gravity. The solutions presented here are, however, not the most general ones, as we have assumed that both the radial and the tangential stresses are equal to the energy density of the loop and also assumed that the metric components are separable as products of functions of r and z , but this is perhaps the first attempts for obtaining the solutions of the full nonlinear Einstein's equations for a circular cosmic string loop. The earlier solution

for the spacetime for a circular string loop , given by Hughes et.al [9] are based on a linearized approximation of the field equations. In the present solution the energy density for the loop blows at a certain $r = r_0$ which evidently corresponds to the site of the string loop.

The authors are grateful to the referee for his valuable comments and suggestions and also for pointing out some serious typographical errors. This work is partially supported by the D.S.T., Govt.of India. One of the author (A.A.S) is thankful to the University Grants Commission, India for financial support.

References

- [1] T.W.Kibble, J.Phys.A **9**,1387 (1976).
- [2] A.Vilenkin, Phys.Rep. **121**,263 (1985).
- [3] A.Vilenkin, Phys.Rev.D **24**, 2082 (1981).
- [4] J.R.Gott, Astrophys.J **288**, 422 (1985).
- [5] T.Vachaspati and A.Vilekin, Phys.Rev.D **31**, 3052 (1985).
- [6] R.Brandenberger, A.Albrecht and N.Turok, Nucl.Phys, **B277**, 605 (1986).
- [7] V.P.Frolov, W.Israel and W.G.Unruh, Phys.Rev.D **39**, 1084 (1989).
- [8] R.Arnowitt, S.Deser and C.W.Misner,in: **Gravitation:An Introduction to current research**, edited by L.witten(Willy, N.Y.1962).
- [9] S.J.Hughes, D.J.McManus and M.A.Vandyck, Phys.Rev.D **47**, 468 (1993).
- [10] D.J.McManus and M.A.Vandyck, Phys.Rev.D **47**, 1491 (1993).
- [11] J.L.Synge, **Relativity: The General Theory**, (North Holland, Amsterdam, 1960).