

GRAVITATION GAUGE GROUP

G.T.Ter-Kazarian¹

Byurakan Astrophysical Observatory, Armenia 378433

June 3, 1996

Abstract

Suggested theory involves a drastic revision of a role of local internal symmetries in physical concept of curved geometry. Under the reflection of fields and their dynamics from Minkowski to Riemannian space a standard gauge principle of local internal symmetries is generalized. The gravitation gauge group is proposed, which is generated by hidden local internal symmetries. In all circumstances, it seemed to be of the most importance for understanding of physical nature of gravity. The most promising aspect in our approach so far is the fact that the energy -momentum conservation laws of gravitational interacting fields are formulated quite naturally by exploiting whole advantage of auxiliary *shadow fields* on flat *shadow space*. The developed mechanism enables one to infer Einstein 's equation of gravitation, but only with strong difference from Einstein 's theory at the vital point of well -defined energy -momentum tensor of gravitational field and conservation laws. The gravitational interaction as well as general distortion of manifold $G(2.2.3)$ with hidden group $U^{loc}(1)$ was considered.

Key words: Gravitation-Internal Symmetries

1 Introduction

In spite of unrivaled by its simplicity and beautiful features, Einstein 's classical theory of gravitation clashes with some basic principle of field theory. This state of affairs has not much changed up to present and proposed abundant models of gravitation are not conducive to provide unartificial and unique recipe for resolving controversial problems of energy-momentum conservation laws of gravitational interacting fields, localization of energy of gravitation waves and also severe problems involved in quantum gravity. It may seem foolhardy to think the rules governing such issues in scope of the theory accounting for gravitation entirely in terms of intricate Riemannian geometry. The difficulties associated with this step are notorious, however, these difficulties are technical. In the main, they stem from the fact that Riemannian geometry, in general, has not admitted a group of isometries. So, the Poincare transformations no longer act as isometries. For example, it is not possible to define energy -momentum as Noether currents related to exact symmetries and so on. On the other hand, the concepts of local internal symmetries and gauge fields [1-3] have become a powerful tool and successfully utilized for the study of electroweak and strong interactions. The intensive attempts have been made for

¹E-mail address:gago@bao.sci.am

constructing a gauge theory of gravitation. But they are complicated sure enough and not generally acceptable. Since none of the solutions of the problems of determination of the gauge group of gravitation and Lagrangian of gravitational field proposed up to the present seems to be wholly convincing, one might gain insight into some of unknown features of phenomena of local internal symmetries and gravity by investigating a new approaches in hope of resolving of such issues. Effecting a reconciliation it is the purpose of present article to explore a number of fascinating features of *generating the gauge group of gravitation by hidden local internal symmetry*. This is the guiding principle framing our discussion. In line with this we feel it of the most importance to generalize standard gauge principle of local internal symmetries. While, a second trend emerged as a formalism of reflection of physical fields and their dynamics from Minkowski space to Riemannian. There is an attempt to supply some of the answers in a concise form to the crucial problems of gravitation and to trace some of the major currents of thoughts under a novel view-point. Exploiting a whole advantage of the field theory in terms of flat space, a particular emphasis will be placed just on the formalism of reflection. This is not a final report on a closed subject, but it is hoped that suggested theory will serve as useful introduction and that it will thereby add the knowledge on the role of local internal symmetries in physical concept of curved geometry. Of course, much remains to be done before one can determine whether this approach can ever contribute to the larger goal of gaining new insight into the theory of gravity.

2 Formulation of Principle

In standard picture, suppose a massless gauge field $\mathbf{B}_l(x_f) = T^a B_l^a(x_f)$ with the values in Lie algebra of group G is a local form of expression of connection in principle bundle with a structure group G . Collection of matter fields are defined as the sections of vector bundles associated with G by reflection $\Phi_f : M^4 \rightarrow E$ that $p \Phi_f(x_f) = x_f$, where $x_f \in M^4$ is a space-time point of Minkowski flat space specified by index (f) . The Φ_f is a column vector denoting particular component of field taking values in standard fiber F_{x_f} upon $x_f : p^{-1}(U^{(f)}) = U^{(f)} \times F_{x_f}$, where $U^{(f)}$ is a region of base of principle bundle upon which an expansion into direct product $p^{-1}(U^{(f)}) = U^{(f)} \times G$ is defined. The various suffixes of Φ_f are left implicit. The fiber is Hilbert vector space on which a linear representation $U_f(x_f) = \exp(-T^a \theta_f^a(x_f))$ of group G with structure constants C^{abc} is given. This space regarded as Lie algebra of group G upon which Lie algebra acts according to law of adjoint representation: $\mathbf{B} \leftrightarrow ad \mathbf{B} \Phi_f \rightarrow [\mathbf{B}, \Phi_f]$. To facilitate writing, we shall consider, in the main, the most important fields of spin $0, \frac{1}{2}, 1$. But developed method may be readily extended to the field of arbitrary spin s , since latter will be treated as a system of $2s$ fermions of half -integral spin. In order to generalize standard gauge principle below we proceed with preliminary discussion.

a. As a starting point we shall assume that under gauge field \mathbf{B}_f the basis vectors e_f^l transformed into four vector fields

$$\hat{e}^\mu(\kappa \mathbf{B}_f) = \hat{D}_l^\mu(\kappa \mathbf{B}_f) e_f^l, \quad (2.1)$$

where \hat{D}_l^μ are real -valued matrix -functions of \mathbf{B}_f , κ is universal coupling constant by which gravitational constant will be expressed (see eq.(4.7)). The double occurrence

of dummy indices, as usual, will be taken to denote a summation extended over their all values. Explicit form of \hat{D}_l^μ will not concern us here, which must be defined under concrete physical considerations [4-8] (sec. 5,6). However, two of the most common restrictions will be placed upon these functions. As far as all known interactions are studied on the base of commutative geometry, then, first of all, the functions \hat{D}_l^μ will be diagonalized at given (μ, l) . As a real -valued function of Hermitian matrix \mathbf{B}_f , each of them is Hermitian too and may be diagonalized by proper unitary matrix. Hence $D_l^\mu(\kappa \mathbf{B}_f) = \text{diag}(\lambda_{l1}^\mu, \lambda_{l2}^\mu, \lambda_{l3}^\mu, \lambda_{l4}^\mu)$, provided with eigen -values λ_{li}^μ as the roots of polynomial characteristic equation

$$C(\lambda_l^\mu) = \det(\lambda_l^\mu I - \hat{D}_l^\mu) = 0. \quad (2.2)$$

Thus

$$e^\mu(\kappa \mathbf{B}_f) = D_l^\mu(\kappa \mathbf{B}_f) e_f^l \quad (2.3)$$

lead to commutative geometry. Consequently, if $e_\nu(\kappa \mathbf{B}_f)$ denotes the inverse matrix - vector $\|D\| \neq 0$, then $\langle e^\mu, e_\nu \rangle = D_l^\mu D_\nu^l = \delta_\nu^\mu$, $D_l^\mu D_\mu^k = \delta_l^k$. A bilinear form on vector fields of sections τ of tangent bundle of R^4 : $\hat{g} : \tau \otimes \tau \rightarrow C^\infty(R^4)$, namely the metric $(g_{\mu\nu})$, is a section of conjugate vector bundle $S^2\tau^*$ (symmetric part of tensor degree) with corresponding components in basis $e_\mu(\kappa \mathbf{B}_f)$

$$g_{\mu\nu} = \langle e_\mu, e_\nu \rangle = \langle e_\nu, e_\mu \rangle = g_{\nu\mu}, \quad e_\mu = g_{\mu\nu} e^\nu. \quad (2.4)$$

In holonomic basis $\hat{g} = g_{\mu\nu} dx^\mu \otimes dx^\nu$. One now has to impose a second restriction on the functions D_l^μ by placing stringent condition upon the tensor $\omega_l^m(x_f)$ implying

$$\partial_l^f D_k^\mu(\kappa \mathbf{B}_f) = \partial_k^f (\omega_l^m(x_f) D_m^\mu(\kappa \mathbf{B}_f)), \quad (2.5)$$

where $\partial_l^f = \frac{\partial}{\partial x_f^l}$, that the $\omega = \omega_l^l$ should be Lorentz scalar function of the trace of curvature form Ω of connection \mathbf{B}_l with the values in Lie algebra of group G. That is

$$\omega = \omega(\text{tr } \Omega) = \omega(d \text{tr } \mathbf{B}), \quad (2.6)$$

provided

$$\begin{aligned} \Omega &= \sum_{l < k} \mathbf{F}_{lk}^f dx_f^l \wedge dx_f^k, \quad \mathbf{B} = \mathbf{B}_l^f dx_f^l, \\ \mathbf{F}_{lk}^f &= \partial_l^f \mathbf{B}_k - \partial_k^f \mathbf{B}_l - ig[\mathbf{B}_l, \mathbf{B}_k]. \end{aligned} \quad (2.7)$$

There up on a functional ω has a null variational derivative $\frac{\delta \omega(\text{tr } \Omega)}{\delta \mathbf{B}_l} = 0$ at local variations of connection $\mathbf{B}_l \rightarrow \mathbf{B}_l + \delta \mathbf{B}_l$, namely, ω is invariant under Lorentz(Λ) and G-gauge groups $\omega = \text{inv}(\Lambda, G)$.

b. Constructing a diffeomorphism $x^\mu(x_f^l) : M^4 \rightarrow R^4$, the holonomic functions $x^\mu(x_f^l)$ satisfy defining relation

$$e_\mu \psi_l^\mu = e_l^f + \chi_l^f(\mathbf{B}_f), \quad (2.8)$$

where

$$\chi_l^f(\mathbf{B}_f) = e_\mu \chi_l^\mu = -\frac{1}{2} e_\mu \int_0^{x_f^f} (\partial_k^f D_l^\mu - \partial_l^f D_k^\mu) dx_f^k. \quad (2.9)$$

The following notational conventions will be used throughout: $\psi_l^\mu = \partial_l^f x^\mu$, where the indices $\mu, \nu, \lambda, \tau, \sigma, \kappa$ stand for variables in R^4 , when l, k, m, n, i, j refer to M^4 . A closer

examination of eq.(2.9) shows that covector $\chi_l^f(\mathbf{B}_f)$ realizes coordinates x^μ by providing a criteria of integration

$$\partial_k^f \psi_l^\mu = \partial_l^f \psi_k^\mu \quad (2.10)$$

and undegeneration $\|\psi\| \neq 0$ [9,10]. Due to eq.(2.6) and eq.(2.9), one has Lorentz scalar gauge invariant functions

$$\chi = \langle e_f^l, \chi_l^f \rangle = \frac{\omega}{2} - 2, \quad S = \frac{1}{4} \psi_\mu^l D_l^\mu = \frac{1}{2} \left(1 + \frac{\omega}{4}\right). \quad (2.11)$$

So, out of a set of arbitrary curvilinear coordinates in R^4 the *real -curvilinear* coordinates may be distinguished, which satisfy eq.(2.8) under all possible Lorentz and gauge transformations. There is a single -valued conformity between corresponding tensors with various suffixes on R^4 and M^4 . While, each co- or contra -variant index transformed incorporating respectively with functions ψ_l^μ or ψ_μ^l . Each transformation of real -curvilinear coordinates $x'^{\mu'} \rightarrow x^\mu$ was generated by some Lorentz and gauge transformations

$$\frac{\partial x'^{\mu'}}{\partial x^\mu} = \psi_{\nu'}^{\mu'}(B_f') \psi_\mu^l(B_f) \Lambda_l^{\nu'}. \quad (2.12)$$

There would then exist preferred systems and group of transformations of real -curvilinear coordinates in R^4 . The wider group of transformations of arbitrary curvilinear coordinates in R^4 would then be of no importance for the field dynamics. If an inverse function ψ_μ^l meets condition

$$\frac{\partial \psi_\mu^l}{\partial x^\nu} \neq \Gamma_{\mu\nu}^\lambda \psi_\lambda^l, \quad (2.13)$$

where $\Gamma_{\mu\nu}^\lambda$ is the usual Christoffel symbol agreed with a metric $g_{\mu\nu}$, then a curvature of R^4 is not vanished [11].

In pursuing the original problem further we are led to the principle point of drastic change of standard gauge scheme to assume hereafter that *single -valued, smooth, double-sided reflection of fields* $\Phi_f \rightarrow \Phi(\Phi_f) : F_{x_f} \rightarrow F_x$, namely $\Phi_f^{l \cdots m}(x_f)$, takes place under local group G , where $\Phi_f \subset F_{x_f}$, $\Phi \subset F_x$, F_x is the fiber upon $x : p^{-1}(U) = U \times F_x$, U is the region of base R^4 . The tensor suffixes were only put forth in illustration of a point at issue. Then

$$\Phi^{\mu \cdots \delta}(x) = \psi_l^\mu \cdots \psi_m^\delta R(\mathbf{B}_f) \Phi_f^{l \cdots m}(x_f) \equiv (R_\psi)_{l \cdots m}^{\mu \cdots \delta} \Phi_f^{l \cdots m}(x_f), \quad (2.14)$$

where $R(\mathbf{B}_f)$ is a reflection matrix. The idea of *general gauge principle* may be framed into requirement of *invariance of physical system of fields* $\Phi(x)$ under the finite local gauge transformations $U_R = R'_\psi U_f R_\psi^+$ of the Lie group of gravitation G_R generated by G , where $R'_\psi = R_\psi(\mathbf{B}'_f)$ if gauge field $\mathbf{B}_f(x_f)$ was transformed under G in standard form. While the corresponding transformations of fields $\Phi(x)$ and their covariant derivatives are written

$$\begin{aligned} \Phi'(x) &= U_R(x) \Phi(x), \\ (g^\mu(x) \nabla_\mu \Phi(x))' &= U_R(x) (g^\mu(x) \nabla_\mu \Phi(x)). \end{aligned} \quad (2.15)$$

The solution of eq.(2.15) may be readily obtained as the reflection of covariant derivatives

$$g^\nu(x) \nabla_\nu \Phi^{\mu \cdots \delta}(x) = S(B_f) \psi_l^\mu \cdots \psi_m^\delta R(\mathbf{B}_f) \gamma^k D_k \Phi_f^{l \cdots m}(x_f), \quad (2.16)$$

where $D_l = \partial_l^f - ig\mathbf{B}_l(x_f)$, $S(B_f)$ is gauge invariant Lorentz scalar, ∇_μ is covariant derivative in R^4 : $\nabla_\mu = \partial_\mu + \Gamma_\mu$, provided with connection [11] $\Gamma_\mu(x) = \frac{1}{2}\Sigma^{\alpha\beta}V_\alpha^\nu(x)\partial_\mu V_{\beta\nu}(x)$, the $\Sigma^{\alpha\beta}$ are generators of Lorentz group, $V_\alpha^\mu(x)$ are the components of affine tetrad vectors e^α in used coordinate net x^μ : $V_\alpha^\mu(x) = \langle e^\mu, e_\alpha \rangle$; one has $g^\mu(x) \Rightarrow e^\mu(x)$ and $\gamma^l \Rightarrow e_f^l$ for fields ($s = 0, 1$); but $g^\mu(x) = V_\alpha^\mu(x)\gamma^\alpha$ for spinor field ($s = \frac{1}{2}$), where γ^α are Dirac's matrices. Since the fields $\Phi_f(x_f)$ no longer hold, the reflected ones $\Phi(x)$ will be regarded as the real physical fields. But a conformity eq.(2.14) and eq.(2.16) enables $\Phi_f(x_f)$ to serve as an auxiliary *shadow fields* on *shadow flat space* M^4 . These notions arise basically from the most important fact that a Lagrangian $L(x)$ of fields $\Phi(x)$ may be obtained under the reflection from a Lagrangian $L_f(x_f)$ of corresponding shadow fields and vice versa. Certainly, the $L(x)$ is also an invariant under the wider group of arbitrary curvilinear transformations $x \rightarrow x'$ in R^4

$$J_\psi L(x)|_{inv(G_R; x \rightarrow x')} = L_f(x_f)|_{inv(\Lambda; G)}, \quad (2.17)$$

where $J_\psi = \|\psi\| \sqrt{1 + 2\|\langle e_f^l, \chi_k^f \rangle\| + \|\langle \chi_l^f, \chi_k^f \rangle\|}^{1/2}$ (see eq.(4.1)). While, *the internal gauge symmetry G remained hidden symmetry, since it screened by gauge group of gravitation G_R* . We note that the tetrad e^α and basis e_f^l vectors meet a condition

$$\begin{aligned} \rho_l^\alpha(x, x_f) &= \langle e^\alpha, e_f^l \rangle = V_\mu^\alpha(x) D_l^\mu(\kappa \mathbf{B}_f), \\ \langle e_f^\alpha, e_f^\beta \rangle &= \langle e^\alpha, e^\beta \rangle = \delta_\beta^\alpha. \end{aligned} \quad (2.18)$$

So, Minkowskian metric is written $\eta^{\alpha\beta} = \text{diag}(1, -1, -1, -1) = \langle e_f^\alpha, e_f^\beta \rangle = \langle e^\alpha, e^\beta \rangle$.

3 Reflection Matrix

One interesting offshoot of general gauge principle is a formalism of reflection. A straightforward calculation for fields $s = 0, 1$ gives the explicit unitary reflection matrix

$$R(x, x_f) = R_f(x_f)R_g(x) = \exp(-i\Theta_f(x_f) - \Theta_g(x)), \quad (3.1)$$

provided

$$\Theta_f(x_f) = g \int_0^{x_f} \mathbf{B}_l(x_f) dx_f^l, \quad \Theta_g(x) = \int_0^x [R_f^+ \Gamma_\mu R_f + \psi^{-1} \partial_\mu \psi] dx^\mu, \quad \psi \equiv (\psi_l^\mu), \quad (3.2)$$

where $\Theta_g = 0$ for scalar field and $\Theta_g + \Theta_g^+ = 0$ for vector field, because of $\Gamma_\mu + \Gamma_\mu^+ = 0$ and $\partial_\mu(\psi^{-1}\psi) = 0$. The function $S(B_f)$ has a form eq.(2.11), since $S(B_f) = \frac{1}{4}R^+(\psi_\mu^l D_l^\mu)R = \frac{1}{4}\psi_\mu^l D_l^\mu$. The infinitesimal gauge transformation $U_f \approx 1 - iT^a \theta_f^a(x_f)$ yields

$$U_R = \exp\left(igC^{abc}T^a \int_0^{x_f} \theta_f^b B_l^c dx_f^l + \Theta_g' - \Theta_g\right), \quad (3.3)$$

where the infinitesimal transformation $B_l'^a = B_l^a + C^{abc}\theta_f^b B_l^c - \frac{1}{g}\partial_l^f \theta_f^a$ was used. For example, a Lagrangian of *isospinor-scalar* shadow field φ_f is in the form

$$\begin{aligned} L_f(x_f) &= J_\psi L(x) = J_\psi [(e^\mu \nabla_\mu \varphi)^+ (e^\mu \nabla_\mu \varphi) - m^2 \varphi^+ \varphi] = \\ &= J_\psi [S(B_f)^2 (D_l \varphi_f)^+ (D_l \varphi_f) - m^2 \varphi_f^+ \varphi_f]. \end{aligned} \quad (3.4)$$

A Lagrangian of *isospinor-vector* Maxwell's shadow field arises in a straightforward manner

$$\begin{aligned} L_f(x_f) &= J_\psi L(x) = \\ &= -\frac{1}{4} J_\psi F_{\mu\nu}^+ F^{\mu\nu} = -\frac{1}{4} J_\psi S(B_f)^2 (F_{\mu\nu}^{(f)})^+ F_{(f)}^{\mu\nu}, \end{aligned} \quad (3.5)$$

provided

$$F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu, \quad F_{\mu\nu}^{(f)} = (\psi_\nu^k D_\mu^f - \psi_\mu^k D_\nu^f) A_k^f, \quad D_\mu^f = D_\mu^l D_l, \quad (3.6)$$

with an additional gauge violating term

$$\begin{aligned} L_G^f(x_f) &= J_\psi L_G(x) = -\frac{1}{2} \zeta_0^{-1} J_\psi (\nabla_\mu A^\mu)^+ \nabla_\nu A^\nu = \\ &= -\frac{1}{2} \zeta_0^{-1} J_\psi S(B_f)^2 (\lambda_m^l D_l A_f^m)^+ \lambda_n^k D_k A_f^n, \quad \lambda_m^l = \frac{1}{2} (\delta_m^l + \omega_m^l), \end{aligned} \quad (3.7)$$

where ζ_0^{-1} is gauge fixation parameter. Finally, a Lagrangian of *isospinor-ghost* fields is written

$$L_{gh}^f(x_f) = J_\psi L_{gh}(x) = J_\psi < (e^\mu \partial_\mu C)^+, e^\nu \partial_\nu C > = J_\psi S(B_f)^2 (D_l C_f)^+ D_l C_f. \quad (3.8)$$

Continuing along this line we come to a discussion of the reflection of spinor field $\Psi_f(x_f)(s = \frac{1}{2})$ as a solution of eq.(2.15)

$$\begin{aligned} \Psi(x) &= R(\mathbf{B}_f) \Psi_f(x_f), \quad \bar{\Psi}(x) = \bar{\Psi}_f(x_f) \tilde{R}^+(\mathbf{B}_f), \\ g^\mu(x) \nabla_\mu \Psi(x) &= S(B_f) R(\mathbf{B}_f) \gamma^l D_l \Psi_f(x_f), \\ (\nabla_\mu \bar{\Psi}(x)) g^\mu(x) &= S(B_f) (D_l \bar{\Psi}_f(x_f)) \gamma^l \tilde{R}^+(\mathbf{B}_f), \end{aligned} \quad (3.9)$$

where $\tilde{R} = \gamma^0 R \gamma^0$, $\Sigma^{\alpha\beta} = \frac{1}{4} [\gamma^\alpha, \gamma^\beta]$, $\Gamma_\mu(x) = \frac{1}{4} \Delta_{\mu, \alpha\beta}(x) \gamma^\alpha \gamma^\beta$, $\Delta_{\mu, \alpha\beta}(x)$ are Ricci's rotation coefficients. The reflection matrix R is in the form eq.(3.1), provided we make change

$$\Theta_g(x) = \frac{1}{2} \int_0^x R_f^+ \{g^\mu \Gamma_\mu R_f, g_\nu dx^\nu\}, \quad (3.10)$$

the $\{, \}$ is an anticommutator. A calculation now gives

$$S(B_f) = \frac{1}{8K} \psi_\mu^l \{ \tilde{R}^+ g^\mu R, \gamma_l \} = inv, \quad (3.11)$$

where

$$\begin{aligned} K &= \tilde{R}^+ R = \tilde{R}_g^+ R_g = \\ &= \exp \left(-\frac{1}{2} \int_0^x \left(\{ R_f^+ \tilde{\Gamma}_\mu^+ g^\mu, g_\nu dx^\nu \} R_f + R_f^+ \{ g^\mu \Gamma_\mu R_f, g_\nu dx^\nu \} \right) \right), \quad \tilde{\Gamma}_\mu^+ = \gamma^0 \Gamma_\mu^+ \gamma^0. \end{aligned} \quad (3.12)$$

Taking into account that $[R_f, g_\nu] = 0$ and substituting [12]

$$\tilde{\Gamma}_\mu^+ g^\nu + g^\nu \Gamma_\mu = -\nabla_\mu g^\nu = 0 \quad (3.13)$$

into eq.(3.12), we get

$$K = 1. \quad (3.14)$$

Since

$$\tilde{U}_R^+ U_R = \tilde{R} U_f^+ \tilde{R}'^+ R' U_f R^+ = \tilde{R} U_f^+ \tilde{R}^+ R U_f R^+, \quad (3.15)$$

where $\tilde{R}'^+ R' = \tilde{R}^+ R = 1$, then

$$\tilde{U}_R^+ U_R = \gamma^0 U_R^+ \gamma^0 U_R = 1. \quad (3.16)$$

A Lagrangian of *isospinor-spinor* shadow field may be written

$$\begin{aligned} L_f(x_f) &= J_\psi L(x) = \\ &= J_\psi \left\{ \frac{i}{2} \left[\bar{\Psi}(x) g^\mu(x) \nabla_\mu \Psi(x) - (\nabla_\mu \bar{\Psi}(x)) g^\mu(x) \Psi(x) \right] - m \bar{\Psi}(x) \Psi(x) \right\} = \\ &= J_\psi \left\{ S(B_f) \frac{i}{2} \left[\bar{\Psi}_f \gamma^l D_l \Psi_f - (D_l \bar{\Psi}_f) \gamma^l \Psi_f \right] - m \bar{\Psi}_f \Psi_f \right\}. \end{aligned} \quad (3.17)$$

In special case if curvature tensor $R_{\mu\nu\tau}^\lambda = 0$, the eq.(2.8) may be satisfied globally in M^4 by putting

$$\psi_l^\mu = D_l^\mu = V_l^\mu = \frac{\partial x^\mu}{\partial \xi^l}, \quad \|D\| \neq 0, \quad \chi_l^f = 0, \quad (3.18)$$

where ξ^l are inertial coordinates. So, $S = J_\psi = 1$, which means that one simply has constructed local G-gauge theory in M^4 both in curvilinear as well as inertial coordinates.

4 Action Principle

Field equations may be inferred from an invariant action

$$S = S_{B_f} + S_\Phi = \int L_{B_f}(x_f) d^4 x_f + \int \sqrt{L_\Phi(x)} d^4 x. \quad (4.1)$$

A Lagrangian $L_{B_f}(x_f)$ of gauge field $\mathbf{B}_l(x_f)$ defined on M^4 is invariant under Lorentz as well as G -gauge groups. But a Lagrangian of the rest of fields $\Phi(x)$ defined on R^4 is invariant under the gauge group of gravitation G_R . Consequently, the whole action eq.(4.1) is G -gauge invariant, since G_R generated by G. Field equations followed at once in terms of Euler-Lagrang variations respectively in M^4 and R^4

$$\begin{aligned} \frac{\delta^f L_{B_f}}{\delta^f B_l^a} &= J_a^l = -\frac{\delta^f L_{\Phi_f}}{\delta^f B_l^a} = -\frac{\partial g^{\mu\nu}}{\partial B_l^a} \frac{\delta(\sqrt{L_\Phi})}{\delta g^{\mu\nu}} = \\ &= -\frac{1}{2} \frac{\partial g^{\mu\nu}}{\partial B_l^a} \frac{\sqrt{D_{k\mu} \delta(\sqrt{L_\Phi})}}{\|D\| \delta D_k^\nu} = -\frac{\sqrt{\partial g^{\mu\nu}}}{2} \frac{\partial g^{\mu\nu}}{\partial B_l^a} T_{\mu\nu}, \\ \frac{\delta L_\Phi}{\delta \Phi} &= 0, \quad \frac{\delta L_\Phi}{\delta \Phi} = 0, \end{aligned} \quad (4.2)$$

where $T_{\mu\nu}$ is the energy -momentum tensor of fields $\Phi(x)$. Making use of Lagrangian of corresponding shadow fields $\Phi_f(x_f)$, in generalized sense, one may readily define the energy -momentum conservation laws and also exploit whole advantage of field theory in terms of flat space in order to settle or mitigate the difficulties whenever they arise including the quantization of gravitation, which will not concern us here. Meanwhile, of course, one is free to carry out an inverse reflection to R^4 whenever it will be needed. To render our discussion here more transparent, below we clarify a relation between

gravitational and coupling constants. So, we consider the theory at the limit of Newton's non-relativistic law of gravitation in the case of static weak gravitational field given by Poisson equation [11]

$$\nabla^2 g_{00} = 8\pi G T_{00}. \quad (4.3)$$

In linear approximation

$$g_{00}(\kappa \mathbf{B}_f) \approx \left(1 + 2\kappa B_k^a(x_f)\theta_a^k\right), \quad \theta_a^k = \left(\frac{\partial D_0^0(\kappa \mathbf{B}_f)}{\partial \kappa B_k^a}\right)_0 = \text{const}, \quad (4.4)$$

then

$$\theta_a^k \nabla^2 \kappa B_k^a = 4\pi G T_{00}. \quad (4.5)$$

Since, the eq.(4.2) must match onto eq.(4.5) at considered limit, then the right-hand sides of both equations should be in the same form

$$\kappa \theta_a^k \frac{\delta^f L_{B_f}}{\delta^f B_k^a} = \kappa \theta_a^k J_k^a \approx -\frac{\kappa^2}{2} \frac{\theta_a^k \partial g^{00}}{\partial (\kappa B_k^a)} T_{00} = \kappa^2 (\theta_a^k \theta_k^a) T_{00}. \quad (4.6)$$

With this final detail carried for one gets

$$G = \frac{\kappa^2}{4\pi \theta_c^2}, \quad \theta_c^2 \equiv (\theta_a^k \theta_k^a). \quad (4.7)$$

At this point we emphasize that Weinberg's argumentation [13], namely, a prediction of attraction between particle and antiparticle, and repulsion between the same kind of particles, which is valid for pure vector theory, no longer holds in suggested theory of gravitation. Although the $\mathbf{B}_l(x_f)$ is vector gauge field, the eq.(4.7) just furnished the prove that only gravitational attraction is existed. One final observation is worth recording. A fascinating opportunity has turned out in the case if one utilizes reflected Lagrangian of Einstein's gravitational field

$$L_{B_f}(x_f) = J_\psi L_E(x) = \frac{J_\psi R}{16\pi G}, \quad (4.8)$$

where R is a scalar curvature. Hence, one readily gets the field equation

$$\left(R_\lambda^\mu - \frac{1}{2}\delta_\lambda^\mu R\right) D_\mu^\lambda \frac{\partial D_l^\lambda}{\partial B_k^a} = -8\pi G U_a^k(\kappa \mathbf{B}_f) = -8\pi G T_{\lambda\nu} D^{\nu l} \frac{\partial D_l^\lambda}{\partial B_k^a}, \quad (4.9)$$

which obviously leads to Einstein's equation. Of course in these circumstances it is straightforward to choose $g^{\mu\nu}$ as the characteristic of gravitational field without referring to gauge field $B_l^a(x_f)$. However, in this case the energy-momentum tensor of gravitational field is well-defined. At this vital point suggested theory strongly differs from Einstein's classical theory. In line with this, one should use the real-curvilinear coordinates, in which the gravitational field was well-defined in the sense that it cannot be destroyed globally by coordinate transformations. While, taking into account general rules eq.(2.12), the energy-momentum tensor of gravitational field may be readily obtained by expressing the energy-momentum tensor of vector gauge field $B_l^a(x_f)$ in terms of metric tensor and its derivatives

$$\begin{aligned} T_{\mu\nu} &= g_{\mu\lambda} T_\nu^\lambda = g_{\mu\lambda} \psi_k^\lambda \psi_\nu^i T_{(f)i}^k = \\ &= g_{\mu\lambda} \psi_k^\lambda \psi_\nu^i \left(\frac{\partial B_l^a}{\partial g^{\mu'\nu'}} \psi_i^\sigma \partial_\sigma g^{\mu'\nu'} \frac{\partial (\partial_\tau g^{\lambda'\tau'})}{\partial (\partial_k^f B_l^a)} \frac{\partial (J_\psi L_E)}{\partial (\partial_\tau g^{\lambda'\tau'})} - \delta_i^k J_\psi L_E \right). \end{aligned} \quad (4.10)$$

At last, we should note that eq.(4.8) is not the simplest one among gauge invariant Lagrangians. Moreover, it must be the same in all cases including eq.(3.18) too. But in last case it contravenes the standard gauge theory. There is no need to contemplate such a drastic revision of physics. So, for our part we prefer G -gauge invariant Lagrangian in terms of M^4

$$L_{B_f}(x_f) = -\frac{1}{4} < \mathbf{F}_{lk}^f(x_f), \mathbf{F}_f^{lk}(x_f) >_K, \quad (4.11)$$

where $\mathbf{F}_{lk}^f(x_f)$ is in the form eq.(2.7), $<, >_K$ is the Killing undegenerate form on the Lie algebra of group G for adjoint representation. Certainly, an explicit form of functions $D_l^\mu(\kappa \mathbf{a}_f)$ should be defined.

5 Gravitation at $G = U^{loc}(1)$

The gravitational interaction with hidden Abelian local group $G = U^{loc}(1) = SO^{loc}(2)$ and one-dimensional trivial algebra $\hat{g} = R^1$ was considered in [4-8], wherein the explicit form of transformation function D_l^μ is defined by making use of principle bundle $p : E \rightarrow G(2.3)$. It is worthwhile to consider the major points of it anew in concise form and make it complete by calculations, which will be adjusted to fit the outlined here theory. We start with a very brief recapitulation of structure of flat manifold $G(2.3) = {}^*R^2 \otimes R^3 = R_+^3 \oplus R_-^3$ provided with the basis vectors $e_{(\lambda\alpha)}^0 = O_\lambda \otimes \sigma_\alpha$, where $< O_\lambda, O_\tau > = {}^*\delta_{\lambda\tau} = 1 - \delta_{\lambda\tau}$, $< \sigma_\alpha, \sigma_\beta > = \delta_{\alpha\beta}$ ($\lambda, \tau = \pm : \alpha, \beta = 1, 2, 3$), δ is Kronecker symbol. A bilinear form on vector fields of sections τ of tangent bundle of $G(2.3)$: $\hat{g}^0 : \tau \otimes \tau \rightarrow C^\infty(G(2.3))$, namely, the metric $(g_{(\lambda\alpha)(\tau\beta)}^0)$ is a section of conjugate vector bundle with components $\hat{g}^0(e_{(\lambda\alpha)}^0, e_{(\tau\beta)}^0)$ in basis $(e_{(\lambda\alpha)}^0)$. The $G(2.3)$ decomposes into three-dimensional ordinary (R_f^3) and time (T_f^3) flat spaces $G(2.3) = R_f^3 \oplus T_f^3$ with signatures $sgn(R_f^3) = (+ + +)$ and $sgn(T_f^3) = (- - -)$. Since all directions in T_f^3 are equivalent, then by notion *time* one implies the projection of time-coordinate on fixed arbitrary universal direction in T_f^3 . By this reduction $T_f^3 \rightarrow T_f^1$ the transition $G(2.3) \rightarrow M^4 = R_f^3 \oplus T_f^1$ may be performed whenever it will be needed. Under massless gauge field $a_{(\lambda\alpha)}(\eta_f)$ associating with $U^{loc}(1)$ the basis $e_{(\lambda\alpha)}^0$ transformed at a point $\eta_f \in G(2.3)$ according to eq.(2.3)

$$e_{(\lambda\alpha)} = D_{(\lambda\alpha)}^{(\tau\beta)} e_{(\tau\beta)}^0. \quad (5.1)$$

While, the matrix D is in the form $D = C \otimes R$, where the distortion transformations $O_{(\lambda\alpha)} = C_{(\lambda\alpha)}^\tau O_\tau$ and $\sigma_{(\lambda\alpha)} = R_{(\lambda\alpha)}^\beta O_\beta$ are defined. Thereby $C_{(\lambda\alpha)}^\tau = \delta_\lambda^\tau + \kappa a_{(\lambda\alpha)} {}^*\delta_\lambda^\tau$, but R is a matrix of the group $SO(3)$ of all ordinary rotations of the planes, each of which involves two arbitrary basis vectors of R_λ^3 , around the orthogonal axes. The angles of permissible rotations will be determined throughwith a special constraint imposed upon distortion transformations, namely, a sum of distortions of corresponding basis vectors O_λ and σ_β has to be zero at given λ :

$$< O_{(\lambda\alpha)}, O_\tau >_{\tau \neq \lambda} + \frac{1}{2} \varepsilon_{\alpha\beta\gamma} \frac{< \sigma_{(\lambda\beta)}, \sigma_\gamma >}{< \sigma_{(\lambda\beta)}, \sigma_\beta >} = 0, \quad (5.2)$$

where $\varepsilon_{\alpha\beta\gamma}$ is an antisymmetric unit tensor. There up on $\tan \theta_{(\lambda\alpha)} = -\kappa a_{(\lambda\alpha)}$, where $\theta_{(\lambda\alpha)}$ is the particular rotation around the axis σ_α of R_λ^3 . Inasmuch as the R should be independent of sequence of rotation axes, then it implies the mean value $R = \frac{1}{6} \sum_{i \neq j \neq k} R^{(ijk)}$, where

$R^{(ijk)}$ the matrix of rotations carried out in sequence (ijk) ($i, j, k = 1, 2, 3$). As it was seen at the outset the field $a_{(\lambda\alpha)}$ was generated by the distortion of basis pseudo-vector O_λ , when the distortion of σ_α has followed from eq(5.2).

Certainly, the whole theory outlined in sections (1-4) will then hold provided we simply replace each single index μ of variables by the pair $(\lambda\alpha)$ and so on. Following to standard rules, next we construct the diffeomorphism $\eta^{(\lambda\alpha)}(\eta_f^{(\tau\beta)}) : G(2.3) \rightarrow G(23)$ and introduce the action eq.(4.1) for the fields. In the sequel, a transition from six-dimensional curved manifold $G(23)$ to four-dimensional Riemannian geometry R^4 is straightforward by making use of reduction of three time-components $e_{0\alpha} = \frac{1}{\sqrt{2}}(e_{(+\alpha)} + e_{(-\alpha)})$ of basis six-vectors

$e_{(\lambda\alpha)}$ to single e_0 in fixed universal direction. Actually, since Lagrangian of fields on R^4 is a function of scalars, namely, $A_{(\lambda\alpha)}B^{(\lambda\alpha)} = A_{0\alpha}B^{0\alpha} + A_\alpha B^\alpha$, so taking into account that $A_{0\alpha}B^{0\alpha} = A_{0\alpha} < e^{0\alpha}, e^{0\beta} > B_{0\beta} = A_0 < e^0, e^0 > B_0 = A_0 B^0$, one readily may perform a required transition. The gravitation field equation is written

$$\partial_f^{(\tau\beta)} \partial_{(\tau\beta)}^f a^{(\lambda\alpha)} - (1 - \zeta_0^{-1}) \partial_f^{(\lambda\alpha)} \partial_{(\tau\beta)}^f a^{(\tau\beta)} = -\frac{1}{2} \sqrt{-g} \frac{\partial g^{(\tau\beta)(\mu\gamma)}}{\partial a_{(\lambda\alpha)}} T_{(\tau\beta)(\mu\gamma)}. \quad (5.3)$$

To render our discussion more transparent, below we consider in detail a solution of spherical-symmetric static gravitational field $a_{(+\alpha)} = a_{(-\alpha)} = \frac{1}{\sqrt{2}} a_{0\alpha}(r_f)$. So, $\theta_{(+\alpha)} = \theta_{(-\alpha)} = -\arctan(\frac{\kappa}{\sqrt{2}} a_{0\alpha})$ and $\sigma_{(+\alpha)} = \sigma_{(-\alpha)}$. It is convenient to make use of spherical coordinates $\sigma_{(+1)} = \sigma_r, \sigma_{(+2)} = \sigma_\theta, \sigma_{(+3)} = \sigma_\varphi$. The transition $G(2.3) \rightarrow M^4$ is performed by choosing in T_f^3 the universal direction along radius-vector: $x_f^{0r} = t_f, x_f^{0\theta} = x_f^{0\varphi} = 0$. Then, from eq.(5.1) one gets

$$e_0 = D_0^0 e_0^0, \quad e_r = D_r^r e_r^0, \quad e_\theta = e_\theta^0, \quad e_\varphi = e_\varphi^0, \quad (5.4)$$

provided $D = C \otimes I$ with components $D_0^0 = 1 + \frac{\kappa}{\sqrt{2}} a_0, D_r^r = 1 - \frac{\kappa}{\sqrt{2}} a_0, a_0 \equiv a_{0r}$, where $e_0^0 = \xi_0 \otimes \sigma_r, e_r^0 = \xi \otimes \sigma_r, e_\theta^0 = \xi \otimes \sigma_\theta, e_\varphi^0 = \xi \otimes \sigma_\varphi, \xi_0 = \frac{1}{\sqrt{2}}(O_+ + O_-)$ and $\xi = \frac{1}{\sqrt{2}}(O_+ - O_-)$. The coordinates $x^\mu(t, r, \theta, \varphi)$ implying $x^\mu(x_f^l) : M^4 \rightarrow R^4$ exist in the whole region $p^{-1}(U) \in R^4$

$$\frac{\partial x^\mu}{\partial x_f^l} = \psi_l^\mu = \frac{1}{2}(D_l^\mu + \omega_l^m D_m^\mu), \quad (5.5)$$

where according to eq.(5.4) one has $x^{0r} = t, x^{0\theta} = x^{0\varphi} = 0$. A straightforward calculation gives non-vanishing components

$$\begin{aligned} \psi_0^0 &= \frac{1}{2} D_0^0, \quad \psi_1^0 = \frac{1}{2} t_f \partial_r^f D_0^0, \quad \psi_1^1 = D_r^r, \quad \psi_2^2 = \psi_3^3 = 0, \\ \omega_1^1 &= \omega_2^2 = \omega_3^3 = 1, \quad \omega_1^0 = t_f \partial_r^f D_0^0. \end{aligned} \quad (5.6)$$

Although $\partial_r \psi_0^0 = \Gamma_{01}^0 \psi_1^0$ and $\partial_r \psi_1^1 = \Gamma_{11}^1 \psi_1^1$, but

$$\partial_t \psi_1^0 = \psi_0^0 \partial_r^f D_0^0 \neq \Gamma_{10}^0 \psi_0^0 = \psi_0^0 \partial_r \ln D_0^0, \quad (5.7)$$

where according to eq.(5.4), the non-vanishing components of Christoffel symbol are written $\Gamma_{01}^0 = \frac{1}{2}g^{00}\partial_r g_{00}$, $\Gamma_{00}^1 = -\frac{1}{2}g^{11}\partial_r g_{00}$, $\Gamma_{11}^1 = \frac{1}{2}g^{11}\partial_r g_{11}$. So, the condition eq.(2.13) holds, namely, the curvature of R^4 is not zero. The curved space R^4 has the group of motions $SO(3)$ with two-dimensional space-like orbits S^2 where the standard coordinates are θ and φ . The stationary subgroup of $SO(3)$ acts isotropically upon the tangent space at the point of sphere S^2 of radius r . So, the bundle $p : R^4 \rightarrow R^2$ has the fiber $S^2 = p^{-1}(x)$, $x \in R^2$ with a trivial connection on it, where R^2 is the factor-space $R^4/SO(3)$. In outside of the distribution of matter with the total mass M , the eq.(5.3) in Feynman gauge reduced to $\nabla_f^2 a_0 = 0$, which has the solution $\frac{\kappa}{\sqrt{2}}a_0 = -\frac{GM}{r_f} = -\frac{r_g}{2r_f}$ (see eq.(4.7)). So, the line element is written

$$ds^2 = (1 - \frac{r_g}{2r_f})^2 dt^2 - (1 + \frac{r_g}{2r_f})^2 dr^2 - r^2(\sin^2 \theta d\varphi^2 + d\theta^2), \quad (5.8)$$

provided by eq.(5.5) and eq.(5.6). Finally, for example, the explicit form of unitary matrix U_R of gravitation group G_R in the case of scalar field reads

$$U_R = R' U_f R^+ = e^{-i[t_f \partial_r^f \theta_f(r_f) + \theta_f(r_f)]}, \quad (5.9)$$

where

$$R = R_f = e^{-\frac{igr_g}{2r_f} t_f}, \quad g(a'_0 - a_0) = \partial_r^f \theta_f(r_f). \quad (5.10)$$

A Lagrangian of charged scalar shadow field eq.(3.4) is in the form

$$L_f(x_f) = \frac{1}{2} \left[\left(\frac{7}{8} \right)^2 (D_l \varphi_f)^* D_l \varphi_f - m^2 \varphi_f^* \varphi_f \right], \quad (5.11)$$

where, according to eq.(2.11) and eq.(5.6), one has $J_\Psi = \frac{1}{2}(1 + \|\omega_l^m\|) = \frac{1}{2}$, $S = \frac{7}{8}$.

6 Distortion of Flat Manifold $G(2.2.3)$ at $G = U^{loc}(1)$

Following to [4-8], the foregoing theory can be readily generalized for the distortion of 12-dimensional flat manifold $G(2.2.3) = {}^*R^2 \otimes {}^*R^2 \otimes R^3 = G_\eta(2.3) \oplus G_u(2.3) = \sum_{\lambda, \mu=1}^2 \oplus R_{\lambda\mu}^3 =$

$R_x^3 \oplus T_x^3 \oplus R_u^3 \oplus T_u^3$ with corresponding basis vectors $e_{(\lambda, \mu, \alpha)} = O_{\lambda, \mu} \otimes \sigma_\alpha \subset G(2.2.3)$, $e_{i(\lambda\alpha)}^0 = O_{i\lambda} \otimes \sigma_\alpha \subset G(2.3)$, where $O_{i+} = \frac{1}{\sqrt{2}}(O_{1,1} + \varepsilon_i O_{2,1})$, $O_{i-} = \frac{1}{\sqrt{2}}(O_{1,2} + \varepsilon_i O_{2,2})$, $\varepsilon_\eta = 1$, $\varepsilon_u = -1$ and $\langle O_{\lambda, \mu}, O_{\tau, \nu} \rangle = {}^*\delta_{\lambda\tau} {}^*\delta_{\mu\nu}$ ($\lambda, \mu, \tau, \nu = 1, 2$), $\langle \sigma_\alpha, \sigma_\beta \rangle = \delta_{\alpha\beta}$. There up on $\langle O_{i\lambda}, O_{i\tau} \rangle = \varepsilon_i \delta_{ij} {}^*\delta_{\lambda\tau}$. The massless gauge field of distortion $a_{(\lambda, \mu, \alpha)}(\zeta_f)$ with the values in Lie algebra of $U^{loc}(1)$ is a local form of expression of connection in principle bundle $p : E \rightarrow G(2.2.3)$ with a structure group $U^{loc}(1)$. Collection of matter fields $\Phi_f(\zeta_f)$ are the sections of vector bundles associated with $U^{loc}(1)$ by reflection $\Phi_f : G(2.2.3) \rightarrow E$ that $p\Phi_f(\zeta_f) = \zeta_f$, where the coordinates $\zeta^{(\lambda, \mu, \alpha)}$ exist in the whole region $p^{-1}(U^{(f)}) \in G(2.2.3)$.

Outlined theory will then hold, while each pair of indices $(\lambda\alpha)$ will be replaced by the (λ, μ, α) . So the basis $e_{(\tau, \nu, \beta)}$ transformed

$$e_{(\lambda\mu\alpha)} = D_{(\lambda\mu\alpha)}^{(\tau, \nu, \beta)} e_{(\tau, \nu, \beta)}, \quad (6.1)$$

provided $D = C \otimes R$, where $O_{(\lambda\mu\alpha)} = C_{(\lambda\mu\alpha)}^{\tau, \nu} O_{\tau, \nu}$, $\sigma_{(\lambda\mu\alpha)} = R_{(\lambda\mu\alpha)}^\beta \sigma_\beta$. The matrices C generate the group of distortion transformations of bi-pseudo-vectors $O_{\tau, \nu}$: $C_{(\lambda\mu\alpha)}^{\tau, \nu} = \delta_\lambda^\tau \delta_\mu^\nu + \kappa a_{(\lambda, \mu, \alpha)}^* \delta_\lambda^\tau \delta_\mu^\nu$, but the matrices R are the elements of the group $SO(3)_{\lambda\mu}$ of ordinary rotations of the planes of corresponding basis vectors of $R_{\lambda\mu}^3$. A special constraint eq.(5.2) holds for basis vectors of each $R_{\lambda\mu}^3$. Thus, the gauge field $a_{(\lambda, \mu, \alpha)}$ was generated by the distortion of bi-pseudo-vectors $O_{\tau, \nu}$. While the rotation transformations R follow due to eq.(5.2) and R implies, as usual, the mean value with respect to sequence of rotation axes. The angles of permissible rotations are $\tan \theta_{(\lambda, \mu, \alpha)} = -\kappa a_{(\lambda, \mu, \alpha)}(\zeta_f)$. The action eq.(4.1) now reads

$$S = S_{a_f} + S_\Phi = \int L_{a_f}(\zeta_f) d\zeta^{(1,1,1)} \wedge \dots \wedge d\zeta^{(2,2,3)} + \int \sqrt{g} L_\Phi(\zeta) d\zeta^{(111)} \wedge \dots \wedge d\zeta^{(223)}, \quad (6.2)$$

where g is the determinant of metric tensor on $G(223)$ and $\zeta^{(\lambda\mu\alpha)}(\zeta^{(\tau, \nu, \beta)}) : G(2.2.3) \rightarrow G(223)$ was constructed according to eq.(2.8)-eq.(2.10), provided

$$\frac{\partial \zeta^{(\lambda\mu\alpha)}}{\partial \zeta^{(\tau, \nu, \beta)}} = \psi_{(\tau, \nu, \beta)}^{(\lambda\mu\alpha)} = \frac{1}{2} \left(D_{(\tau, \nu, \beta)}^{(\lambda\mu\alpha)} + \omega_{(\tau, \nu, \beta)}^{(\rho, \omega, \gamma)} D_{(\rho, \omega, \gamma)}^{(\lambda\mu\alpha)} \right). \quad (6.3)$$

Right through the variational calculations one infers the field equations eq.(4.2), while

$$\begin{aligned} & \partial^{(\tau, \nu, \beta)} \partial_{(\tau, \nu, \beta)} a^{(\lambda, \mu, \alpha)} - (1 - \zeta_0^{-1}) \partial^{(\lambda, \mu, \alpha)} \partial_{(\tau, \nu, \beta)} a^{(\tau, \nu, \beta)} = J^{(\lambda, \mu, \alpha)} = \\ & = -\frac{1}{2} \sqrt{g} \frac{g^{(\tau\nu\beta)(\rho\omega\gamma)}}{\partial a_{(\lambda, \mu, \alpha)}} T_{(\tau\nu\beta)(\rho\omega\gamma)}, \end{aligned} \quad (6.4)$$

provided

$$T_{(\tau\nu\beta)(\rho\omega\gamma)} = \frac{2\delta(\sqrt{g}L_\Phi)}{\sqrt{g}\delta g^{(\tau\nu\beta)(\rho\omega\gamma)}}. \quad (6.5)$$

The curvature of manifold $G(2.3) \rightarrow G(23)$ (sec. 5), which leads to four-dimensional Riemannian geometry R^4 , is a familiar distortion

$$a_{(1,1,\alpha)} = a_{(2,1,\alpha)} \equiv \frac{1}{\sqrt{2}} a_{\eta(+\alpha)}, \quad a_{(1,2,\alpha)} = a_{(2,2,\alpha)} \equiv \frac{1}{\sqrt{2}} a_{\eta(-\alpha)}, \quad (6.6)$$

when $\sigma_{(11\alpha)} = \sigma_{(21\alpha)} \equiv \sigma_{(+\alpha)}$, $\sigma_{(12\alpha)} = \sigma_{(22\alpha)} \equiv \sigma_{(-\alpha)}$. Hence $e_{i(\lambda\alpha)} = O_{i(\lambda\alpha)} \otimes \sigma_{(\lambda\alpha)}$, where $O_{i(\lambda\alpha)} = O_{i\lambda} + \frac{\kappa}{\sqrt{2}} \varepsilon_i a_{\eta(\lambda\alpha)}^* \delta_\lambda^\tau O_{i\tau}$. In the aftermath $G(223) = G_\eta(23) \oplus G_u(23)$. The other important case of inner-distortion

$$a_{(1,1,\alpha)} = -a_{(2,1,\alpha)} \equiv \frac{1}{\sqrt{2}} a_{u(+\alpha)}, \quad a_{(1,2,\alpha)} = -a_{(2,2,\alpha)} \equiv \frac{1}{\sqrt{2}} a_{u(-\alpha)} \quad (6.7)$$

leads to $\sigma_{(11\alpha)} = -\sigma_{(21\alpha)} \equiv \sigma_{(+\alpha)}$, $\sigma_{(12\alpha)} = -\sigma_{(22\alpha)} \equiv \sigma_{(-\alpha)}$. Hence, $O_{i(\lambda\alpha)} = O_{i\lambda} + \frac{\kappa}{\sqrt{2}} \varepsilon_i a_{u(\lambda\alpha)}^* \delta_\lambda^\tau O_{i\tau}$, where the $e_{i(\lambda\alpha)} = O_{i(\lambda\alpha)} \otimes \sigma_{(\lambda\alpha)}$ is the basis in inner-distorted manifold $G(23)$.

7 Discussion and Conclusions

If we collect together the results just established we have finally arrived at quite promising, if not entirely satisfactory, proposal of generating the group of gravitation by hidden local internal symmetries. While, under the reflection of shadow fields from Minkowski flat space to Riemannian, one framed the idea of general gauge principle of local internal symmetries G into requirement of invariance of physical system of reflected fields on R^4 under the Lie group of gravitation G_R of local gauge transformations generated by G . This yields the invariance under wider group of arbitrary curvilinear coordinate transformations in R^4 . While the energy -momentum conservation laws are well -defined. The fascinating prospect emerged for resolving or mitigating a shortage of controversial problems of gravitation including its quantization by exploiting whole advantage of field theory in terms of flat space. In the aftermath, one may carry out an inverse reflection into R^4 whenever it will be needed. It was proved that only gravitational attraction exists. One may easily infer Einstein's equation of gravitation, but with strong difference at the vital point of well -defined energy -momentum tensor of gravitational field and conservation laws. Nevertheless, for our part we prefer standard gauge invariant Lagrangian eq.(4.11), while the functions eq.(2.3) ought to be defined. We considered the gravitational interaction with hidden Abelian group $G = U^{loc}(1)$ with the base $G(2.3)$ of principle bundle as well as the distortion of manifold $G(2.2.3)$, which yields both the curvature and inner-distortion of space and time. However, we believe we have made good headway by presenting a reasonable framework whereby one will be able to verify the basic ideas and illustrate main features of drastic generalization of standard gauge principle of local internal symmetries in order to gain new insight into physical nature of gravity.

Acknowledgements

It is pleasure to express my gratitude to K.Yerknapetian and A.Vardanian for support.

References

- [1] Weil H., Sitzungsber. d. Berl. Akad., 1918, 465.
- [2] Yang C.N., Mills R.L., Phys. Rev., 1954, v. 96, 191.
- [3] Utiyama R., Phys. Rev., 1956, v.101, 1597.
- [4] Ter-Kazarian G.T., Selected Questions of Theoretical and Mathematical Physics, 1986, VINITI, N5322-B86, Moscow.
- [5] Ter-Kazarian G.T.,Comm. Byurakan Obs., 1989,v.62, 1.
- [6] Ter-Kazarian G.T., Astrophys. and Space Sci., 1992, v.194, 1.
- [7] Ter-Kazarian G.T.,I.C.T.P.-Preprint, IC/94/290, Trieste, Italy.
- [8] Ter-Kazarian G.T.,1995, hep-th/9510110.
- [9] Dubrovin B.A., Novikov S.P., Fomenko A.T., Contemporary Geometry, 1986, Nauka, Moscow.

- [10] Pontryagin L.S., Continuous Groups, 1984, Nauka, Moscow.
- [11] Weinberg S., 1972, Gravitation and Cosmology, J.W. and Sons, New York.
- [12] Fock V.A., Zeitsch. fur Phys., 1929, v.57, 261.
- [13] Weinberg S., 1964, Phys. Rev. b133, 1318.