

Can we obtain Conserved Currents in Supersymmetric Quantum Cosmology?

P.V. Moniz

*DAMTP, University of Cambridge
Silver Street, Cambridge, CB3 9EW, UK*

e-mail: prlv10@amtp.cam.ac.uk; paul.vargasmoniz@ukonline.co.uk

URL: <http://www.damtp.cam.ac.uk/user/prlv10>

In this paper we report whether conserved currents can be sensibly defined in supersymmetric minisuperspaces. Our analysis deals with a $k = +1$ FRW model. Supermatter in the form of scalar supermultiplets is included. We show that conserved currents *cannot* be adequately established except for some very simple scenarios. More details can be found in ref. ^{3,5}.

N=1 supergravity ¹ constitutes a “square-root” ² of gravity: in finding a physical state Ψ , it is sufficient to solve the Lorentz and supersymmetry constraints; Ψ will then consequently obey the Hamiltonian constraints^a. Supersymmetry thus induces advantageous features: in many cases we have to solve simple *first-order* differential equations in the bosonic variables³. This contrasts with the situation without supersymmetry: a *second-order* (Wheeler-DeWitt) equation has to be solved, employing boundary conditions ⁴. Therefore, it is quite tempting to address from a supersymmetric point of view the issue of probability densities for a quantum state Ψ and conservation equations of the type $\nabla_a J^a = 0$ (see also ref. ⁵).

The approach that we employ here is based on a differential operator representation for the fermionic variables. *This constitutes the correct procedure*: it is totally consistent with the existence of second-class constraints and subsequent Dirac brackets in supergravity theories. These then imply that fermionic variables and their Hermitian conjugates are intertwined within a canonical coordinate-momentum relation. It should also be pointed out that other authors have pursued objectives similar to ours but with *different* methods ^{6,7,8}. In particular, *rigid* supersymmetry was used⁶, other approach⁷ was *not* supersymmetric. Furthermore, a wave function arranged as a vector was used⁸ where the Lorentz constraints were severely restricted, but an improved approach seems to have been found⁹.

Let us consider the action of the more general theory of N=1 supergravity in the presence of gauged supermatter¹. Our physical variables include the tetrad $e^{AA'}_\mu$ and the gravitinos which are represented by $\psi^A_\mu, \bar{\psi}^{A'}_\mu$. The “overline” denotes Hermitian conjugation. The tetrad for a $k = +1$ FRW model can be written as $e_{a\mu} = \text{diag}[N(\tau), aE_{\hat{a}i}]$, where \hat{a} and i run from 1 to 3 and $E_{\hat{a}i}$ is a basis of left-invariant 1-forms on the unit S^3 with volume $\sigma^2 = 2\pi^2$. We take ψ^A_0 and $\bar{\psi}^{A'}_0$ to be functions of time only and $\psi^A_i = e^{AA'}_i \bar{\psi}_{A'}$, $\bar{\psi}^{A'}_i = e^{AA'}_i \psi_A$. We have introduced the new spinors ψ_A and their Hermitian conjugate, $\bar{\psi}_{A'}$, which are also functions of time only. The scalar supermultiplet will consist of complex scalar

^aFor a review on canonical quantization of supersymmetric minisuperspaces see ref. ³.

fields $\phi(t) = re^{i\theta}$, $\bar{\phi}(t)$ and their spin- $\frac{1}{2}$ partners $\chi_A(t)$, $\bar{\chi}_{A'}(t)$.

Simple Dirac brackets are then obtained, namely

$$[\chi_A, \bar{\chi}_B]_D = -i\epsilon_{AB}, [\psi_A, \bar{\psi}_B]_D = i\epsilon_{AB}, [a, \pi_a]_D = 1, [\phi, \pi_\phi]_D = 1, [\bar{\phi}, \pi_{\bar{\phi}}]_D = 1, \quad (1)$$

and the rest of the brackets are zero. At this point we choose $(\chi_A, \psi_A, a, \phi, \bar{\phi})$ to be the coordinates of the configuration space and $(\bar{\chi}_A, \bar{\psi}_A, \pi_a, \pi_\phi, \pi_{\bar{\phi}})$ to be the momentum operators in this representation.

The Lorentz constraints take the form $J_{AB} = \psi_{(A}\bar{\psi}_{B)} - \chi_{(A}\bar{\chi}_{B)} = 0$, which implies that the most general form for the wave function of the universe is

$$\Psi = A + B\psi^C\psi_C + C\psi^C\chi_C + D\chi^C\chi_C + E\psi^C\psi_C\chi^D\chi_D, \quad (2)$$

where A, B, C, D, E are functions of $a, \phi, \bar{\phi}$, only. The bosonic coefficients present in eq. (2) satisfy attractive relations^b in a 3-dimensional minisuperspace:

$$\frac{\partial(A \cdot E)}{\partial a} + \frac{\partial(A \cdot E)}{\partial \theta} - ir \left(\frac{\partial E}{\partial r} A - \frac{\partial A}{\partial r} E \right) = 0, \quad (3)$$

$$D_a(B \cdot D) + \frac{\partial(B \cdot D)}{\partial \theta} - ir \left(\frac{\partial B}{\partial r} D - \frac{\partial D}{\partial r} B \right) = 0, \quad (4)$$

with $D_a = \partial_a - 6/a$. However, the presence of the terms $ir \left(\frac{\partial E}{\partial r} A - \frac{\partial A}{\partial r} E \right)$ and $ir \left(\frac{\partial B}{\partial r} D - \frac{\partial D}{\partial r} B \right)$ in eq. (3) and (4), respectively, clearly prevent us from obtaining conservation equations of the type $\nabla \cdot J = 0$. The reason can be identified with the variable θ no longer being a cyclical coordinate when supersymmetry is present (see eq. (5) below). To understand this argument, let us consider a FRW model with complex scalar fields in non-supersymmetric quantum cosmology¹⁰. The corresponding action implies that the conjugate momentum $\pi_\theta \sim r^2 a^3 \frac{\partial \theta}{\partial t}$ is a constant and θ constitutes a cyclical coordinate. However, the situation in the corresponding supersymmetric scenario is quite different. The π_θ takes the form

$$\begin{aligned} \pi_\theta &= \frac{2\sigma^2}{(1+r^2)^2} r^2 a^3 \frac{\partial \theta}{\partial t} + \frac{5\sigma^2 r^2 a^3}{\sqrt{2}(1+r^2)^3} n^{AA'} \bar{\chi}_{A'} \chi_A - \frac{3\sigma^2 r^2 a^3}{\sqrt{2}(1+r^2)} n^{AA'} \psi_A \bar{\psi}_{A'} \\ &+ \frac{ir\sigma^2 a^3 e^{-i\theta}}{\sqrt{2}(1+r^2)^2} 3n_{AA'} \chi^A \bar{\psi}^{A'} + \frac{ir\sigma^2 a^3 e^{i\theta}}{\sqrt{2}(1+r^2)^2} 3n_{AA'} \chi^{A'} \psi^A \\ &+ \frac{ir\sigma^2 a^3 e^{-i\theta}}{\sqrt{2}(1+r^2)^2} \chi^A \psi_{0A} - \frac{ir\sigma^2 a^3 e^{i\theta}}{\sqrt{2}(1+r^2)^2} \bar{\chi}_{A'} \bar{\psi}_0^{A'}. \end{aligned} \quad (5)$$

and thus notice there are terms in the action that do *not* allow θ to be a cyclical coordinate. So, π_θ would not be a constant. And this will imply the absence of satisfactory conserved currents. A similar situation^c would occur in usual quantum cosmology with a matter Lagrangian taken from the Wess-Zumino model, due to the non-trivial interaction with fermion fields.

^bObtained from the supersymmetry constraint equations — $S_A \Psi = 0$ and $\bar{S}_{A'} \Psi = 0$ — see ref. ^{3,5} for more details.

The author is grateful to S. Kamenshchik for having pointed out this to him.

Overall, our message in this paper is that *generic* conserved currents do not seem feasible to obtain *directly* from the supersymmetry constraints equations. Only for very simple scenarios does this become possible. Otherwise, conserved currents (and consistent probability densities) may only be obtained upon the use of subsequent Wheeler-DeWitt-like equations. These are derived through the associated supersymmetric algebra of constraints.

In our view, the fundamental reason for our conclusions is related with the following. A physical supersymmetric wave functional Ψ takes values in a Grassman algebra. Such algebra is formed by complex linear combinations of products of anti-commuting elements such as the gravitino ψ_i^A . Hence, $\Psi \left[e_\mu^{AA'}, \psi_\mu^A, \bar{\psi}_\mu^{A'}; \phi, \bar{\phi}, \chi^A, \bar{\chi}^{A'} \right]$ embodies more than a wave function arranged as a vector and satisfying a Dirac-like equation. Furthermore, the first-order differential equations derived from the supersymmetry, Lorentzian and Grassmanian-valued Ψ constitute more than a simple set of conditions: They rather represent the action of the supersymmetry constraints on *different* fermionic representations of Ψ , related by a (coordinate-momentum) fermionic Fourier transformation³.

Acknowledgements

This work was supported by the JNICT/PRAXIS-XXI Fellowship BPD/6095/95. The author is grateful to A. Yu. Kamenshchik for useful comments. Conversations with R. Graham, H. Luckock and O. Oubr  gon motivated some of the analysis discussed in this paper and ref.⁵.

References

1. P. van Nieuwenhuizen, Phys. Rep. **68** (1981) 189.
2. C. Teitelboim, Phys. Rev. Lett. **38** (1977) 1106.
3. P. Moniz, *Supersymmetric Quantum Cosmology — Shaken not Stirred*, (Review Report) Int. J. Mod. Phys.-**A11** (1996) 4321, and references therein.
4. See for example, J. Halliwell, in: *Proceedings of the Jerusalem Winter School on Quantum Cosmology and Baby Universes*, edited by T. Piran et al, World Scientific, (Singapore, 1990) and references therein.
5. P. Moniz, *Conserved Currents in Supersymmetric Quantum Cosmology?*, gr-qc/9606047, to be published in Int. J. Mod. Phys. — **D** (1997).
6. H. Luckock and C. Oliwa, Phys. Rev. **D. 51** (1995) 5883;
7. R. Mallett, Class. Quantum Grav. **12** (1994) L1;
8. O. Oubr  gon, private communications; communication presented at the GR14 conference, August 6-12 1995, Florence, Italy
9. O. Obregon and C. Ramirez, preprint; A. Macias, E. Mielke and J. Socorro, preprint
10. A. Yu. Kamenshchik, I. M. Khalatnikov and A. V. Toporensky, Phys. Lett. **B357** (1995) 36.