

Microcanonical Quantum Statistics of Schwarzschild Black Holes

Günter Scharf
 Institut für Theoretische Physik,
 Universität Zürich,
 Winterthurerstr. 190 , CH-8057 Zürich, Switzerland
 E-Mail: Scharf@physik.unizh.ch

Abstract: It is shown that a quantized Schwarzschild black hole, if described by a square root energy spectrum with exponential multiplicity, can be treated as a microcanonical ensemble without problem leading to the expected thermodynamical properties.

PACS: 04.60.Kz; 04.70.Dy; 97.60.Lf

Recently H.A.Kastrup [1-3] made the interesting observation that a Schwarzschild black hole with energy spectrum $E_n = \sigma\sqrt{n}E_P$, ($n = 1, 2, \dots$, E_P = Planck energy) and multiplicity g^n [4] leads to the expected thermodynamical properties if the divergent canonical partition function Z is analytically continued [2]. The analytic continuation in the complex g -plane gives a complex Z , the desired thermodynamics is obtained from its imaginary part. This situation has some similarity with Langer's nucleation theory [5]. Therefore, it is argued [3] that the complex canonical partition function might signal the instability of the black hole. On the other hand, it is known that the statistical ensembles are not always equivalent. In this case it is necessary to return to the microcanonical ensemble which has the widest range of validity. For a black hole this is even more appealing because it is hard to imagine a thermal bath coupled to it. We will quickly see that our system can be treated microcanonically without problem. This is even the simplest microcanonical calculation I know because it uses mathematics from high school, only.

Let energy eigenvalues E_n with multiplicities $\nu(n)$ be

$$E_n = b\sqrt{n}, \quad \nu(n) = g^n, \quad n = 1, 2, \dots \quad (1)$$

where $b = \sigma E_P$, $\sigma = O(1)$, $g > 1$ and $E_P = \sqrt{\hbar c^5/G}$ is Planck's energy. The microcanonical partition function $\Omega(E)$ is equal to the number of eigenvalues below the energy E , hence

$$\Omega(E) = \sum_{n=1}^{n(E)} g^n = \frac{g^{n(E)} - 1}{g - 1}, \quad (2)$$

where

$$n(E) = \frac{E^2}{b^2} \quad (3)$$

according to (1). To find the thermodynamics we have to compute

$$\omega = \frac{\partial \Omega}{\partial E} = \frac{2 \log g}{g-1} \frac{E}{b^2} g^{E^2/b^2}. \quad (4)$$

Then the microcanonical temperature is given by

$$k_B T = \frac{\Omega}{\omega} = \frac{g^{E^2/b^2} - 1}{g^{E^2/b^2}} \frac{b^2}{2E \log g}. \quad (5)$$

Its inverse is equal to

$$\beta = \frac{2E}{b^2} \log g \left[1 + \mathcal{O}\left(g^{-E^2/b^2}\right) \right]. \quad (6)$$

Substituting for E the rest energy $E = Mc^2$ and [2]

$$\sigma^2 = \frac{\log g}{4\pi}, \quad (7)$$

we get Hawking's temperature

$$\beta_H = \frac{8\pi Mc^2}{E_P^2}, \quad (8)$$

up to the exponentially small corrections in (6). One can turn the thing around and leave $n(E)$ in (2) unspecified. Then one finds $\beta = n'(E) \log g$. If one assumes that $\beta \sim Mc^2 = E$, one derives the square root spectrum $n(E) \sim E^2$, $E \sim \sqrt{n}$.

The entropy is given by

$$\frac{S}{k_B} = \log \Omega = \frac{\log g}{b^2} E^2 - \log(g-1) + \mathcal{O}\left(g^{-E^2/b^2}\right). \quad (9)$$

Neglecting the constant second term and the exponentially small correction we have

$$\frac{S}{k_B} = \frac{\log g}{\sigma^2} \left(\frac{E}{E_P} \right)^2 = \frac{\log g}{\sigma^2} \left(\frac{Mc^2}{E_P} \right)^2. \quad (10)$$

If we insert Planck's energy

$$E_P = \sqrt{\frac{\hbar c^5}{G}},$$

and the above value (7) of σ^2 we find

$$\frac{S}{k_B} = \frac{4GM^2}{\hbar c} = \frac{A}{4\pi l_P^2}, \quad (11)$$

where $l_P^2 = \hbar G/c^3$ is the Planck length squared and $A = 4\pi R_S^2 = 16\pi G^2 M^2/c^4$ is the area of the horizon (R_S = Schwarzschild radius).

The surprising consequence of these very simple results is that the black hole looks like an innocent microcanonical equilibrium state.

References

- [1] H.A. Kastrup, Physics Letters **B 385** (1996) 75
- [2] H.A. Kastrup, gr-qc/9707009
- [3] H.A. Kastrup, gr-qc/9710032
- [4] J.D. Bekenstein, V.F.Mukhanov, Physics Letters **B 360** (1995) 7
- [5] J.S. Langer, in: Systems Far from Equilibrium (Lecture Notes in Physics 132) p.12, L.Garrido ed. Springer Verlag, New York, Heidelberg, Berlin, 1980