

FROM TAUB NUMBERS TO THE BONDI MASS^a

E.N. Glass^b

Physics Department, University of Michigan, Ann Arbor, MI, USA

Abstract

Taub numbers are studied on asymptotically flat backgrounds with Killing symmetries. When the field equations are solved for a background spacetime and higher order functional derivatives (higher order variational derivatives of the Hilbert Lagrangean) are solved for perturbations from the background, such perturbed space-times admit zeroth, first, and second order Taub numbers. Zeroth order Taub numbers are Komar constants (upto numerical factors) or Penrose-Goldberg constants of the background. For a Killing symmetry of the background, first order Taub numbers give the contribution of the linearized perturbation to the associated background quantity, such as the perturbing mass. Second order Taub numbers give the contribution of second order perturbations to the background quantity. The Bondi mass is a sum of first and second order Taub numbers on a Minkowski background.

1 INTRODUCTION

To define the Bondi mass one needs an asymptotically flat manifold and the notion of future null infinity \mathcal{I}^+ . The Bondi metric¹ is axisymmetric and has 4 metric functions V, U, β, γ .

$$g_{\mu\nu}^{Bondi} dx^\mu dx^\nu = (V e^{2\beta}/r - r^2 U^2 e^{2\gamma}) du^2 + 2e^{2\beta} du dr + 2r^2 U e^{2\gamma} du d\theta - r^2 (e^{2\gamma} d\theta^2 + e^{-2\gamma} \sin^2 \theta d\phi^2)$$

where $\gamma = O(1/r)$, $\beta = O(1/r^2)$, $U = O(1/r^2)$. The asymptotic form of metric function V is $V = r - 2M(u, \theta) + O(1/r)$. $M(u, \theta)$ is the Bondi mass aspect

$$-2M(u, \theta) = \Psi_2^0 + \bar{\Psi}_2^0 + \partial_u(\sigma^0 \bar{\sigma}^0).$$

The Bondi mass is the 2-surface integral of the mass aspect over a topological 2-sphere at \mathcal{I}^+

$$M_{Bondi} = -\frac{1}{8\pi} \int_{S^2} [\Psi_2^0 + \bar{\Psi}_2^0 + \partial_u(\sigma^0 \bar{\sigma}^0)] d\Omega$$

2 EARLY CALCULATIONS OF THE BONDI MASS

The first computation of the Bondi mass was done by J.N. Goldberg² with use of the Einstein pseudotensor and von Freud superpotential: $(\sqrt{-g} Einstein)_\alpha^\beta = \partial_\nu (superpotential)_\alpha^{[\beta\nu]} + (pseudotensor)_\alpha^\beta$.

^aTo appear in Class. Quan. Grav. **14**, 1899 (1997)

^bPermanent address: Physics Department, University of Windsor, Ontario, CANADA

Calculating in an asymptotically Cartesian frame and transforming to null spherical coordinates, Goldberg integrated the superpotential over an asymptotic 2-sphere at future null infinity and, using the generator of asymptotic time translations, obtained the Bondi mass.

Winicour and Tamburino³ constructed a tensorial calculation by modifying the Komar superpotential. For the null surface $u = \text{const.}$ they added a term which eliminated off-surface derivatives. Using an asymptotic symmetry, the integral of the modified Komar superpotential, called a "linkage", yields the Bondi mass at \mathcal{I}^+ . Unfortunately the linkage construction does not arise from a variational principle.

3 TAUB METHOD

The Taub method is fully developed in^{4,5,6}. We write the Bondi metric in a perturbation expansion as $g_{\alpha\beta}^{Bondi}(\lambda) = \eta_{\alpha\beta} + \lambda h_{\alpha\beta}^{(1)} + \lambda^2 h_{\alpha\beta}^{(2)} + \dots$. To insure a single coordinate system for each tensor in the perturbation expansion, the Minkowski background metric is constructed from the Bondi metric. We solve the Newman-Penrose form of Einstein's equations with initial data $\Psi_0(u_0, r, \theta) = \Psi_1^0(u_0, \theta) = \Psi_2^0(u_0, \theta) = \sigma^0(u, \theta) = 0$. One obtains Bondi metric functions $V = r$, $U = \beta = \gamma = 0$, i.e. the Minkowski metric. The symmetry generator of mass, the background timelike Killing vector, $k_t = \partial_u$, is covariant constant. The Taub superpotential $U_{Taub}^{\alpha\beta}(k_t, h) = (-g)^{\frac{1}{2}} k_t^\nu (\delta_\nu^{[\alpha} h^{\beta]}{}_{;\mu} - \delta_\nu^{[\alpha} h^{;\beta]} + h_\nu^{[\alpha;\beta]})$ is used to compute Taub numbers where the n^{th} Taub number is $\tau_n(k_t, h^{(n)}) = -\frac{1}{8\pi} \int_{\partial\mathcal{N}} U_{Taub}^{\alpha\beta}(k_t, h^{(n)}) dS_{\alpha\beta}$.

Each of the $h^{(n)}$ and hence each of the τ_n derives from a variational derivative of the Hilbert action, the n^{th} number from the $n+1^{th}$ variation.

The Bondi mass results from the sum of first and second numbers: $M_{Bondi} = \tau_1(k_t, h^{(1)}) + \tau_2(k_t, h^{(2)})$. τ_1 has the non-radiative mass contribution and τ_2 contains the news function.

4 PERTURBATION CALCULATION

To find τ_1 solve $DEin(h^{(1)}) = 0$ for $h_{\mu\nu}^{(1)}$ using the Newman-Penrose equations on the Minkowski background with tetrad $l = du$, $n = \frac{1}{2}du + dr$, $m = -(r/\sqrt{2})(d\theta + i\sin\theta d\phi)$. Find the spin coefficients, metric components, and Weyl tensor components with initial data $\{\Psi_0(u_0, r, \theta), \Psi_1^0(u_0, \theta), \Psi_2^0(u_0, \theta), \sigma^0(u, \theta)\}$.

$$h_{\mu\nu}^{(1)} = [(\Psi_2^0 + \bar{\Psi}_2^0)/r] l_\mu l_\nu - (2\bar{\sigma}^0/r) m_\mu m_\nu - (2\sigma^0/r) \bar{m}_\mu \bar{m}_\nu - (\bar{\rho}\bar{\sigma}^0/r)(l_\mu m_\nu + m_\mu l_\nu) - (\bar{\rho}\sigma^0/r)(l_\mu \bar{m}_\nu + \bar{m}_\mu l_\nu).$$

Substitute into $U_{Taub}^{\alpha\beta}(k_t, h^{(1)})$ and obtain $\tau_1 = -\frac{1}{8\pi} \int_{S^2} (\Psi_2^0 + \bar{\Psi}_2^0) d\Omega$. This is M_{Bondi} when the news $\partial_u \sigma^0 = 0$. Vacuum Schwarzschild has $\sigma = 0$, $\Psi_2^0 = -m$ and so $M_{Bondi} = \tau_1 = m$.

To find τ_2 iterate the Newman-Penrose equations. The second order Bianchi identities for the Weyl tensor components have first order quantities as sources whereas the first order equations were source-free⁷.

$$U_{Tab}^{\alpha\beta}(k_t, h^{(2)}) = \sqrt{-g} l^{[\alpha} n^{\beta]} [-2 \frac{\partial_u(\sigma^0 \bar{\sigma}^0)}{r^2} + O_3].$$

$$\tau_2(k_t, h^{(2)}) = -\frac{1}{8\pi} \int_{S^2} [\partial_u(\sigma^0 \bar{\sigma}^0)] d\Omega.$$

$M_{Bondi} = \tau_1 + \tau_2$. Higher orders fall off faster and do not contribute to the Bondi mass

1. H. Bondi, M. van der Burg, and A. Metzner, Proc. Roy. Soc. (London) **A269**, 21 (1962).
2. J.N. Goldberg, Phys. Rev. **131**, 1367 (1963).
3. L. Tamburino and J. Winicour, Phys. Rev. **150**, 1039 (1966).
4. E.N. Glass, Phys. Rev. **D 47**, 474 (1993).
5. M.G. Naber and E.N. Glass, J. Math. Phys. **35**, 5969 (1994).
6. E.N. Glass and Mark Naber, J. Math. Phys. **35**, 1834 (1994).
7. W.E. Couch, R.J. Torrence, A.I. Janis, and E.T. Newman, J. Math. Phys. **9**, 484 (1968).