

Non-uniqueness of the third post-Newtonian binary point-mass dynamics

Piotr Jaranowski^{1,2} and Gerhard Schäfer³

¹*Institute of Physics, Białystok University
Lipowa 41, 15-424 Białystok, Poland **

²*Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik
Schlaatzweg 1, 14473 Potsdam, Germany*

³*Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität
Max-Wien-Platz 1, 07743 Jena, Germany †*

Abstract

It is shown that the recently found non-uniqueness of the third post-Newtonian binary point-mass ADM-Hamiltonian is related to the non-uniqueness at the third post-Newtonian approximation of the applied ADM-coordinate conditions.

PACS number(s): 04.25.Nx, 04.30.Db, 97.60.Jd, 97.60.Lf

In a recent paper [1] the authors reported on the non-uniqueness of the 3rd post-Newtonian ADM-Hamiltonian for binary point-mass systems. The term in the Hamiltonian which came out to be ambiguous, in the center-of-mass reference frame, is given by (see Eqs. (71) and (75) in [1])

$$\omega \frac{G^3 m_1 m_2}{c^6} p_{1i} p_{1j} \partial_{1i} \partial_{1j} \left(-\frac{1}{r_{12}} \right). \quad (1)$$

In Eq. (1) m_1 and m_2 denote the masses of the bodies 1 and 2, respectively, $r_{12} = |\mathbf{x}_1 - \mathbf{x}_2|$ is their relative coordinate distance, where \mathbf{x}_a ($a = 1, 2$) denotes the position of the a th body. For the momenta of the bodies $p_{1i} = -p_{2i}$ holds ($i = 1, 2, 3$); ∂_{ai} denotes the partial derivative with respect to x_a^i . G and c are the Newtonian gravitational constant and the speed of light, respectively. The ambiguity in the Hamiltonian is expressed by an unspecified finite number ω .

In our treatment we applied the following generalized isotropic ADM-coordinate conditions (see Eqs. (7-4.22) and (7-4.23) in [2], respectively Eqs. (3) and (4) in [1])

$$g_{ij} = \left(1 + \frac{1}{8}\phi \right)^4 \delta_{ij} + h_{ij}^{TT}, \quad (2)$$

$$\pi^{ii} = 0, \quad (3)$$

where g_{ij} denotes the 3-metric and π^{ij} the field momentum (canonical conjugate to g_{ij}), h_{ij}^{TT} is the transverse-traceless (with respect to the flat-space metric) part of $g_{ij} - \delta_{ij}$. The form of the isotropic part of Eq. (2) stems from the Schwarzschild metric in isotropic coordinates.

*E-mail: pio@alpha.fuw.edu.pl

†E-mail: gos@tpi.uni-jena.de

To leading order in powers of $1/c$ (this is enough for the following) the coordinate condition (3) reads ($x^0 = ct$)

$$2\partial_i g_{0i} - \partial_0 g_{ii} = 0. \quad (4)$$

Note that the coordinate conditions (2) can exactly be written as

$$3\partial_j g_{ij} - \partial_i g_{jj} = 0. \quad (5)$$

Let us define the following infinitesimal coordinate transformation

$$x'^\mu = x^\mu + \epsilon^\mu, \quad \epsilon^0 = 0, \quad \epsilon^i = \alpha \frac{G^3 m_1^2 m_2^2}{2c^6 M} \partial_i (r_1^{-1} + r_2^{-1}), \quad (6)$$

where $M = m_1 + m_2$ denotes the total mass of the system, $r_a = |\mathbf{x} - \mathbf{x}_a|$, and α is a pure number. This transformation induces the following leading order transformation in the metric coefficients, keeping the independent variables fixed (see, e.g., Eq. (16) in [3])

$$g'_{\mu\nu}(t, x^i, x_a^i(t), p_{ai}(t)) = g_{\mu\nu}(t, x^i, x_a^i(t), p_{ai}(t)) - g_{\mu\lambda} \partial_\nu \epsilon^\lambda - g_{\nu\lambda} \partial_\mu \epsilon^\lambda. \quad (7)$$

Hereof the transformation of the Hamiltonian results, plugging the expression (7) into the test-mass Hamiltonian (see, e.g., Eq. (5.2) in [4]) and identifying in turn the test mass with the source masses. This gives

$$H' = H + \alpha \frac{G^3 m_1 m_2}{c^6} p_{1i} p_{1j} \partial_{1i} \partial_{1j} \left(\frac{1}{r_{12}} \right). \quad (8)$$

Outside the mass points, the transformation (6) keeps invariant the equations (2) and (3), respectively (5) and (4). Towards spacelike infinity, the perturbation (6) dies out very fast, like $1/r^2$, implying an $1/r^3$ -decay for the metric perturbation (7).

The shift in equation (8) is identical with equation (1). This shows that the dynamical ambiguity found in [1] is related to the ambiguity in the coordinate system (in quantum field theory those gauge ambiguities are well known and result in the DeWitt-Faddeev-Popov ghost fields). A corresponding ambiguity is likely to exist also in harmonic coordinates as one can infer from [5], page 120. For extended bodies, neither the coordinate-system ambiguity arises nor is the dynamical ambiguity present.

Acknowledgments

This work was supported by the Max-Planck-Gesellschaft Grant No. 02160-361-TG74 (GS) and by the KBN grant No. 2 P303D 021 11 (PJ).

References

- [1] P. Jaranowski and G. Schäfer, gr-qc/9712075, submitted to Phys. Rev. D.
- [2] R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation: An Introduction to Current Research*, edited by L. Witten (John Wiley, New York, 1962), p. 227.
- [3] T. Damour and G. Schäfer, Gen. Rel. Grav. **17**, 879 (1985).
- [4] G. Schäfer, Ann. Phys. (N.Y.) **161**, 81 (1985).
- [5] T. Damour, in *Gravitational Radiation*, edited by N. Deruelle and T. Piran (North-Holland, Amsterdam, 1983), p. 59.