

Smooth sandwich gravitational waves

J. Podolský

*Department of Theoretical Physics, Faculty of Mathematics and Physics,
Charles University, V Holešovičkách 2, 180 00 Prague 8, Czech Republic*

Electronic address: podolsky@mbox.troja.mff.cuni.cz

Abstract

Gravitational waves which are smooth and contain two asymptotically flat regions are constructed from the homogeneous pp -waves vacuum solution. Motion of free test particles is calculated explicitly and the limit to an impulsive wave is also considered.

PACS number(s): 04.30.-w, 04.20.Jb, 98.80.Hw

The widely known class of pp -waves [1] is characterized by the existence of a quadruple Debever-Penrose null vector field which is covariantly constant. In vacuum the metric can be written as

$$ds^2 = 2 d\zeta d\bar{\zeta} - 2 du dv - (f + \bar{f}) du^2 , \quad (1)$$

where $f(u, \zeta)$ is an arbitrary function of u , analytic in ζ . The simplest case for which the metric describes gravitational waves arises when f is of the form

$$f(u, \zeta) = d(u)\zeta^2 , \quad (2)$$

where $d(u)$ is an *arbitrary* function of u ; such solutions are called homogeneous pp waves (or “plane” gravitational waves). Performing the transformation [2]

$$\zeta = \frac{1}{\sqrt{2}} (Px + iQy) , \quad v = \frac{1}{2} (t + z + PP'x^2 + QQ'y^2) , \quad u = t - z , \quad (3)$$

where real functions $P(u) \equiv P(t - z)$, $Q(u) \equiv Q(t - z)$ are solutions of differential equations

$$P'' + d(u)P = 0 , \quad Q'' - d(u)Q = 0 , \quad (4)$$

(here prime denotes the derivative with respect to u) the metric can be written in the form

$$ds^2 = -dt^2 + P^2 dx^2 + Q^2 dy^2 + dz^2 , \quad (5)$$

which is more suitable for physical interpretation. Considering free test particles standing at fixed x , y and z , their *relative* motion in the x -direction is given by the function $P(u)$ while it is given by $Q(u)$ in the y -direction. The motions are unaffected in the z -direction which demonstrate transversality of the waves.

Assuming “profile” functions $d(u)$ non-vanishing on some finite interval of u only, sandwich gravitational waves were constructed in [3]-[6] and elsewhere. Here we consider the function $d(u)$ of the form

$$d(u) = \frac{a}{\cosh^2(bu)} , \quad (6)$$

where a and b are arbitrary real positive constants. Since all derivatives of $d(u)$ are continuous the corresponding gravitational waves are *smooth*, contrary to “standard” sandwich waves explicitly presented in literature. On the other hand, the space-times (1) given by Eqs. (2) and (6) do not contain flat regions in front of the wave and behind it. They are curved everywhere, becoming flat only *asymptotically* as $u \rightarrow \pm\infty$. Therefore, we should call them “smooth asymptotic sandwich waves”.

In order to find the form (5) of the metric exhibiting naturally the particle motions we solve the equations (4). Introducing a substitution

$$\xi = -\tanh(bu) , \quad (7)$$

the equations take the form

$$(1 - \xi^2) \frac{d^2 R}{d\xi^2} - 2\xi \frac{dR}{d\xi} \pm \frac{a}{b^2} R = 0 , \quad (8)$$

where the upper sign is applied for $R = P$ and the lower sign for $R = Q$. Clearly, a general solution can be written as a linear combination of the Legendre functions of the first kind $P_\alpha(\xi)$ and the second kind $Q_\alpha(\xi)$ where $\alpha = (\sqrt{1 \pm 4a/b^2} - 1)/2$. It is natural to impose the conditions $R(u \rightarrow -\infty) \rightarrow 1$ and $R'(u \rightarrow -\infty) \rightarrow 0$ so that the metric (5) is written in explicit Minkowski form in the asymptotic region where $u \rightarrow -\infty$. These conditions are satisfied by particular solutions of the form

$$P(u) = P_\mu(\xi(u)) , \quad Q(u) = P_\nu(\xi(u)) , \quad (9)$$

where

$$\mu = (\sqrt{1 + 4a/b^2} - 1)/2 , \quad \nu = (\sqrt{1 - 4a/b^2} - 1)/2 . \quad (10)$$

Typical behavior of the functions P, Q is shown in Fig. 1. It can be observed that in both asymptotic regions $u \rightarrow \pm\infty$ the particles move uniformly. This can be shown analytically using the definition $P_\alpha(\xi) \equiv F(-\alpha, \alpha+1; 1; (1-\xi)/2)$ where F is the hypergeometric function. Clearly, $P \rightarrow 1, Q \rightarrow 1$ as $u \rightarrow -\infty$. For $u \rightarrow +\infty$ (corresponding to $\xi \rightarrow -1$) we use the identity 15.3.10. in [7] according to which

$$F\left(-\alpha, \alpha+1; 1; \frac{1-\xi}{2}\right) = -\frac{\sin \pi \alpha}{\pi} \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(n!)^2} \times \left[2\psi(n+1) - \psi(n-\alpha) - \psi(\alpha+n+1) - \ln\left(\frac{1+\xi}{2}\right)\right] \left(\frac{1+\xi}{2}\right)^n, \quad (11)$$

where ψ is the digamma function. Using the identity $\psi(-\alpha) = \psi(1-\alpha) + 1/\alpha$ and the limit $\ln[(1+\xi)/2] \rightarrow -2bu$ for $u \rightarrow +\infty$ we conclude that

$$P(u \rightarrow +\infty) = C_1 + C_2 u, \quad Q(u \rightarrow +\infty) = D_1 + D_2 u, \quad (12)$$

where

$$C_1 = \frac{\sin \pi \mu}{\pi} \left[\frac{1}{\mu} + \psi(1+\mu) + \psi(1-\mu) - 2\psi(1) \right], \quad C_2 = -\frac{2b}{\pi} \sin \pi \mu, \\ D_1 = \frac{\sin \pi \nu}{\pi} \left[\frac{1}{\nu} + \psi(1+\nu) + \psi(1-\nu) - 2\psi(1) \right], \quad D_2 = -\frac{2b}{\pi} \sin \pi \nu. \quad (13)$$

For particular values of μ corresponding to $\mu = m = 0, 1, 2, \dots$, the constant C_2 vanishes so that $P(u \rightarrow +\infty) = C_1$. These cases coincide with solutions $P(u)$ given by Legendre *polynomials* $P_m(\xi)$ and we easily get $C_1 = P_m(\xi = -1) = (-1)^m$. Explicit particular solutions of this type arise when $a/b^2 = m(m+1)$ so that $P(u) = -\tanh(\sqrt{a/2}u)$ for $m = 1$, $P(u) = [3\tanh^2(\sqrt{a/6}u) - 1]/2$ for $m = 2$, etc. Note also that there are no analogous solutions for $Q(u)$ since (10) admits $\nu \in \langle -1, 0 \rangle$ only. Therefore, the only non-negative integer (for which the constant D_2 vanishes) is $\nu = 0$ representing a trivial case $d(u) = 0$. For all other values of ν the constant D_2 is positive so that $Q(u) \rightarrow +\infty$ as $u \rightarrow +\infty$.

We can also use the above results for construction of impulsive gravitational waves. The sequence of smooth functions (6) approach the δ function (in a distributional sense) as $a \rightarrow 0$ if the second parameter is $b = 2a$ (so that the normalization condition $\int_{-\infty}^{+\infty} d(u) du = 1$ holds for arbitrary a). Considering this limit, indicated also in Fig. 1, $\alpha = (\sqrt{1 \pm 1/a} - 1)/2 \rightarrow 0$ and using Eqs. (13) we get

$$P(u) = 1 - u \Theta(u), \quad Q(u) = 1 + u \Theta(u), \quad (14)$$

where Θ is the Heaviside step function ($\Theta = 0$ for $u < 0$, $\Theta = 1$ for $u > 0$). Therefore, particle motion in the impulsive gravitational wave is the same as in the limit of “standard” sandwich waves [2].

Acknowledgments

I acknowledge the support of grants No. GACR-202/96/0206 and No. GAUK-230/1996 from the Czech Republic and Charles University.

References

- [1] Kramer D., Stephani H., MacCallum M.A.H., and Herlt H.: Exact Solutions of the Einstein's Field Equations, Cambridge University Press, Cambridge, 1980.
- [2] Penrose R.: in General Relativity (Ed. L.O'Raiheartaigh), Clarendon Press, Oxford, 1972.
- [3] Bondi H., Pirani F.A.E., and Robinson I.: Proc. Roy. Soc **A 251** (1959) 519.
- [4] Bondi H. and Pirani F.A.E.: Proc. Roy. Soc **A 421** (1989) 395.
- [5] Rindler W.: Essential Relativity, Springer, New York, 1977.
- [6] Podolský J. and Veselý K.: Czech. J. Phys. **48** (1998) 871.
- [7] Abramowitz M. and Stegun I.A.: Handbook of Mathematical Functions, Dover Publications, New York, 1964.

Figure Caption

Fig. 1. Typical behavior of the functions $P(u)$ and $Q(u)$ determining relative motion of free test particles (initially at rest) in x and y -directions, respectively, caused by smooth sandwich gravitational waves for different values of $a = \frac{1}{48}, \frac{2}{48}, \dots, \frac{12}{48}$, with $b = 2a$. The values $a = \frac{1}{8}, \frac{1}{24}, \frac{1}{48}$ correspond to $\mu = m = 1, 2, 3$, respectively, for which $P \rightarrow (-1)^m$ as $u \rightarrow +\infty$.

This figure "Smooth.gif" is available in "gif" format from:

<http://arxiv.org/ps/gr-qc/9805098v2>