

Quantum entropy of two-dimensional extreme charged dilaton black hole

Bin Wang^{1,20}, Ru-Keng Su^{1,20} and P.K.N.Yu³

¹ China Center of Advanced Science and Technology (World Laboratory), P.O.Box 8730,
Beijing 100080, P.R.China

² Department of Physics, Fudan University, Shanghai 200433, P.R.China

³ Department of Physics and Materials Science, City University of Hong Kong, Hong Kong

Abstract

By using Hawking's treatment as well as Zaslavskii's treatment respectively and the brick wall model, two different values of classical entropy and quantum entropy of scalar fields in the two-dimensional extreme charged dilaton black hole backgrounds have been obtained. A new divergent term emerges in the quantum entropy under the extreme limit for Zaslavskii's treatment and its connection with the phase transition has been addressed.

PACS number(s): 04.70.Dy, 04.20.Gz, 04.62.+v.

⁰Email: binwang@fudan.ac.cn, Permanent address: Dept. of Physics, Shanghai Normal Univ., Shanghai 200234, P.R.China

⁰Email: rksu@fudan.ac.cn

Two different treatments suggested by Hawking et al[1-3] and Zaslavskii [4] respectively have been employed to calculate the classical entropy of extremal black holes (EBH) recently. These different treatments start with different points and lead to very different results. Starting with the original four-dimensional (4D) Reissner-Nordström (RN) EBH, Hawking et al claimed that a RN EBH has zero entropy, infinite proper distance l between the horizon and any fixed point. It cannot be formed from its nonextremal counterpart [1,5]. However, in a grand canonical ensemble, if first taking the boundary limit $r_+ \rightarrow r_B$, where r_+ is the event horizon and r_B is the boundary of the cavity, and then the extremal limit $q \rightarrow m$, Zaslavskii [4] proved that a RN nonextremal black hole (NEBH) can approach the extreme state as closely as one likes in the topological sector of nonextremal configuration. Its entropy still obeys the Bekenstein-Hawking formula. Since these two different treatments approach to two different topological objects, we suggested that there are two kinds of EBHs in the nature. The first kind with zero entropy, zero Euler characteristic, is the original EBH; and the second kind with entropy $A/4$, non-zero Euler characteristic, can be formed from its nonextremal counterpart [6].

Obviously, if these two kinds of EBHs do exist in the nature, their different effects will not only be demonstrated on the classical level, but also be exhibited in the quantum study. Many calculations, including the brick wall model [7-10], Pauli-Villars regular theory [11] and etc., have been used to deal with the entropy of quantum scalar fields on black hole backgrounds under WKB approximation or one-loop approximation respectively. It is believed that fields propagating in the region just outside the horizon give the main quantum fluctuational contribution to the black hole entanglement entropy [7-9,12-14]. Since these two kinds of EBHs' backgrounds are very different, one can predict that the quantum corrections of the scalar field in different kinds of EBHs will have considerable differences. This paper involves from an attempt to study the quantum fluctuational corrections of these two EBHs.

To illustrate our results transparently, for simplicity, we investigate two-dimensional (2D) charged dilaton black hole (CDBH), which is an asymptotically flat solution for the heterotic string [15,16]. The action has the form

$$I = - \int_M \sqrt{g} e^{-2\phi} [R + 4(\nabla\phi)^2 + \lambda^2 - \frac{1}{2}F^2] - 2 \int_{\partial M} e^{-2\phi} K \quad (1)$$

which has a black hole solution metric

$$ds^2 = -g(r)dt^2 + g^{-1}(r)dr^2 \quad (2)$$

$$g(r) = 1 - 2me^{-\lambda r} + q^2e^{-2\lambda r} \quad (3)$$

$$e^{-2\phi} = e^{-2\phi_0}e^{\lambda r}, \quad A_0 = \sqrt{2}qe^{-\lambda r} \quad (4)$$

where m and q are the mass and electric charge of the black hole respectively. The horizons are located at $r_{\pm} = (1/\lambda) \ln(m \pm \sqrt{m^2 - q^2})$.

Using the finite-space formulation of black hole thermodynamics, employing the grand canonical emsemble and putting the black hole into a cavity as usual[4,17], we calculate the free energy and entropy of the CDBH. To simplify our calculations, we introduce a coordinate transformation

$$r = \frac{1}{\lambda} \ln[m + \frac{1}{2}e^{\lambda(\rho+\rho_0^*)} + \frac{m^2 - q^2}{2}e^{-\lambda(\rho+\rho_0^*)}] \quad (5)$$

where ρ_0^* is an integral constant, and rewrite Eq.(2) to a particular gauge

$$ds^2 = -g_{00}(\rho)dt^2 + d\rho^2 \quad (6)$$

The event horizon locate at $\rho_+ = (1/\lambda) \ln \sqrt{m^2 - q^2} - \rho_0^*$. The Euclidean action takes the form

$$I = - \int_{\partial M} \sqrt{\frac{1}{g_{11}}} e^{-2\phi} \left(\frac{1}{2} \frac{\partial_1 g_{00}}{g_{00}} - 2\partial_1 \phi \right) \quad (7)$$

The dilaton charge is found to be

$$D = e^{-2\phi_0} \left(m + \frac{1}{2}e^x + \frac{m^2 - q^2}{2}e^{-x} \right) \quad (8)$$

$$x = \lambda(\rho + \rho_0^*) \quad (9)$$

The free energy, $F = I/\beta$, where β is the proper periodicity of Euclideanized time at a fixed value of the spatial coordinate and has the form $\beta = 1/T_w = \sqrt{g_{00}}/T_c$. T_c is the inverse periodicity of the Euclidean time at the horizon

$$T_c = \frac{\lambda \sqrt{m^2 - q^2}}{2\pi(m + \sqrt{m^2 - q^2})} \quad (10)$$

Using the formula of entropy $S = -(\partial F/\partial T_w)_D$, we obtain

$$S = \frac{2\pi e^{-2\phi} [m + \frac{e^x}{2} + \frac{(m^2 - q^2)e^{-x}}{2}] [1 + (m^2 - q^2)e^{-2x}] \sqrt{m^2 - q^2} (m + \sqrt{m^2 - q^2})}{(m^2 - q^2) + m [\frac{e^x}{2} + \frac{(m^2 - q^2)e^{-x}}{2}]} \quad (11)$$

We are now in a position to extend the above calculations to EBH. We are facing two limits, namely, the extreme limit $q \rightarrow m$ and the boundary limit

$$x \rightarrow x_+ = \lambda(\rho_+ + \rho_0^*) = \ln \sqrt{m^2 - q^2} \quad (12)$$

We can take the limits in different orders: (A) by first taking the extreme limit $q \rightarrow m$ and then the boundary limit. This corresponds to the treatment of Hawking et al. by starting with an original first kind of EBHs. (B) by first taking the boundary limit $x \rightarrow x_+$ and then the extreme limit $q \rightarrow m$. This corresponds to the treatment of Zaslavskii and obtains the second kind of EBHs. To do our limits procedures mathematically, we may take $x = x_+ + \epsilon, \epsilon \rightarrow 0^+$ and $m = q + \eta, \eta \rightarrow 0^+$, where ϵ and η are infinitesimal quantities with different orders of magnitude, and substitute them into Eq(11). It can easily be shown that in treatment (A)

$$S_{CL}(A) = 0 \quad (13)$$

which is in consistent with the result in [18]. However in treatment (B)

$$S_{CL}(A) = 4\pi m e^{-2\phi_0} \quad (14)$$

which is just the Bekenstein-Hawking entropy. These results support that there are two kinds of 2D charged dilaton EBHs [6] which have completely different values of classical entropy.

Extending our calculations to quantum cases, we suppose that the CDBH is enveloped by a scalar field, and the whole system, the hole and the scalar field, are filling into a cavity. The wave equation of the scalar field is

$$\frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \phi) - m^2 \phi = 0 \quad (15)$$

Substituting the metric Eq.(2) into Eq.(15), we find

$$E^2 (1 - 2me^{-\lambda r} + q^2 e^{-2\lambda r})^{-1} f + \frac{\partial}{\partial r} [(1 - 2me^{-\lambda r} + q^2 e^{-2\lambda r}) \frac{\partial f}{\partial r}] - m^2 f = 0 \quad (16)$$

Introducing the brick wall boundary condition[7]

$$\phi(x) = 0 \text{ at } r = r_+ + \epsilon$$

$$\phi(x) = 0 \text{ at } r = L$$

and calculating the wave number $K(r, E)$ and the free energy F , we get

$$K^2(r, E) = (1 - 2me^{-\lambda r} + q^2 e^{-2\lambda r})^{-1} [(1 - 2me^{-\lambda r} + q^2 e^{-2\lambda r})^{-1} E^2 - m^2] \quad (17)$$

$$F_{QM} = -\frac{\pi}{6\beta^2\lambda} \left[\frac{1}{2} \ln(R^2 - 2mR + q^2) + \frac{m}{2\sqrt{m^2 - q^2}} \ln \frac{R - m - \sqrt{m^2 - q^2}}{R - m + \sqrt{m^2 - q^2}} \right] \quad (18)$$

where $R = e^{\lambda(r_+ + \epsilon)}$, and $\epsilon \rightarrow 0$ is the coordinate cutoff parameter. To extend the above discussion to EBH, we are facing two limits $\epsilon \rightarrow 0$ and $q \rightarrow m$ again. For treatment (A), taking the extreme limit first and $\epsilon \rightarrow 0$ (i.e. $r \rightarrow r_+$) afterwards, we obtain

$$F_{QM}(A) = -\frac{\pi}{6\beta^2\lambda} \left(\frac{m}{m\lambda\epsilon} + \ln \frac{1}{m\lambda\epsilon} \right) \quad (19)$$

For treatment (B), adopting $\epsilon \rightarrow 0$ (i.e. $r \rightarrow r_+$) at first and the extreme limit afterwards, we get

$$F_{QM}(B) = -\frac{\pi}{6\beta^2\lambda} \left(\ln \frac{1}{m\lambda\epsilon} + \frac{m}{2\sqrt{m^2 - q^2}} \ln \frac{2\sqrt{m^2 - q^2}}{\lambda\epsilon(m + \sqrt{m^2 - q^2})} \right) \quad (20)$$

For the sake of following discussion, we leave the term $\sqrt{m^2 - q^2}$ in the last term of Eq(22) for the moment.

Using the formula $S = \beta^2(\partial F / \partial \beta)$, the entropy in two treatments can be obtained as

$$S_{QM}(A) = \frac{\pi}{3\beta\lambda} \left(\frac{m}{m\lambda\epsilon} + \ln \frac{1}{m\lambda\epsilon} \right) \quad (21)$$

$$S_{QM}(B) = \frac{\pi}{3\beta\lambda} \left(\ln \frac{1}{m\lambda\epsilon} + \frac{m}{2\sqrt{m^2 - q^2}} \ln \frac{2\sqrt{m^2 - q^2}}{\lambda\epsilon(m + \sqrt{m^2 - q^2})} \right) \quad (22)$$

Obviously, different treatments again lead to different entropy of the scalar field in the black hole background. These results support that there are two kinds of 2D charged dilaton EBHs from the quantum point of view.

It is worth noting, in addition to the usual ultraviolet divergence $\epsilon \rightarrow 0$, which has been found for both 4D and 2D NEBH cases [7-14] and was suggested capable of being overcome by different renormalization methods [9,19], a new divergent term for Zaslavskii treatment emerges in Eq(22). The last term of the right-hand-side of Eq(22) diverges by taking the limit $m \rightarrow q$. This term does not appear in Eq(21) for the treatment of Hawking et al. To explain these results, we go over the arguments of Hawking et al. and that of Zaslavskii again. Due to topological differences, Hawking et al. claimed that the first kind of EBH and its NEBH counterpart are completely different objects. They cannot be transformed from each other. This kind of EBH is the original EBH and can only be prepared at the

beginning of the universe [1]. Since we put $q = m$ first and then calculate its entropy by using the usual thermodynamical approach [17], naturally its quantum entropy includes ultraviolet divergence only. But for the second kind of EBH suggested by Zaslavskii, it can be transformed as closely as one likes from its nonextremal counterpart by taking the limit $m \rightarrow q$. Keeping in mind that the entropy of scalar field in the EBH background was obtained through WKB approximation, the divergent quantum entropy due to taking $q \rightarrow m$ reflects in fact the divergent quantum fluctuations of the entanglement entropy of the whole system including the EBH and the scalar field. According to the statistical physics, the infinite fluctuation breaks down the rigorous meanings of thermodynamical quantities and is just the characteristic of the point of phase transition. This conclusion is in good agreement with previous studies about the phase transition of black hole [20-23]. With the aid of Landau-Lifshitz order-disorder phase transition theory, in a series of previous papers [20-23], it has been found a phase transition will happen when NEBH approaches to its EBH counterpart. The new divergent term of the quantum fluctuation of entropy when $q \rightarrow m$ in Eq(22) supports the viewpoint that a phase transition will happen when a NEBH approaches to the second kind of EBH.

In summary, we have extended two different treatments from 4D studies to 2D CDBH and studied its classical entropy as well as quantum entropy. In addition to the zero classical entropy obtained by using Hawking treatment [18], a nonzero classical result has been derived for the second kind of EBH by Zaslavskii treatment. We have also calculated the quantum entropy of scalar field in the EBH background, and proved that these two different treatments again lead to two different results. Our results support that there are two kinds of 2D charged dilaton EBHs [6]. We have found a new fascinating divergent term in the quantum entropy due to $q \rightarrow m$ in the Zaslavskii's treatment, which corresponds to a phase transition and is in agreement with previous classical studies.

This work was supported in part by NNSF of China.

References

- [1] S.W.Hawking, G.Horowitz and S.Ross, Phys.Rev.D51,4302(1995)
- [2] C.Teitelboim, Phys.Rev.D51,4315(1995)
- [3] A.Ghosh and P.Mitra, Phys.Rv.Lett.78,1858(1997), ibid77,4848(1996),
ibid80,3413(1998)
- [4] O.B.Zaslavskii,
Phys.Rev.Lett.76,2211(1996), Phys.Rev.D56,2188(1997), Phys.Rev.Lett.80,3412(1998),
Phys.Rev.D56,6695(1997)
- [5] B.Wang, R.K.Su, P.K.N.Yu and E.C.M.Young, Phys.Rev.D57, 5284(1998)
- [6] B.Wang and R.K.Su, Phys.Lett.B (in press)
- [7] G. 't Hooft, Nucl.Phys.B 256,727(1985)
- [8] V.Frolov and I.Novikov, Phys.Rev.D48,4545(1993)
- [9] S.N.Solodukhin, Phys.Rev.D51,609,618(1995), ibid D54,3900(1996)
- [10] A.Ghosh and P.Mitra, Phys.Lett.B357,295(1995)
- [11] J.G.Demers, R.Lafrance and R.C.Myers, Phys.Rev.D52,2245(1995)
- [12] M.Srednicki, Phys.Rev.Lett71,666(1993)
- [13] C.Callen and F.Wilezek, Phys.Lett.B333,54(1994)
- [14] J.Ellis, N.E.Mavromato, E.Winstanley and D.V.Nanopoulos,
Mod.Phys.Lett.A12,243(1997)
- [15] C.Nappi and A.Pasquinucci, Mod.Phys.Lett.A7,3337(1992)
- [16] M.D.McGuigan, C.R.Nappi and S.A.Yost, Nucl.Phys.B375,121(1992)
- [17] H.W.Bradeen, J.D.Brown, B.F.Whiting and J.W.York, Phys.Rev.D42,3376(1990)
- [18] A.Kumar and K.Ray, Phys.Rev.D51,5954(1995)
- [19] L.Susskind and J.Uglum, Phys.Rev.D50,2700(1994)

- [20] R.K.Su, R.G.Cai and P.K.N.Yu, Phys.Rev.D50,2932(1994)
R.G.Cai, R.K.Su and P.K.N.Yu, Phys.Rev.D48,3473(1993), ibid D52,6186(1995)
- [21] O.Kaburaki, Phys.Lett.A217,316(1996)
- [22] C.O.Lousto, Phys.Rev.D51,1733(1995)