

Energy-momentum and gauge conservation laws

G.Giachetta, L.Mangiarotti,¹ and G.Sardanashvily²

¹*Department of Mathematics and Physics, University of Camerino, Camerino 62032, Italy*

²*Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia*

We treat energy-momentum conservation laws as particular gauge conservation laws when generators of gauge transformations are horizontal vector fields on fibre bundles. In particular, the generators of general covariant transformations are the canonical horizontal prolongations of vector fields on a world manifold. This is the case of the energy-momentum conservation in gravitation theories. We find that, in main gravitational models, the corresponding energy-momentum flows reduce to the generalized Komar superpotential. We show that the superpotential form of a conserved flow is the common property of gauge conservation laws if generators of gauge transformations depend on derivatives of gauge parameters. At the same time, dependence of a conserved flow on gauge parameters make gauge conservation laws form-invariant under gauge transformations.

PACS numbers: 03.50.-z, 04.20.Cv, 11.15.-q, 11.30.-g

I. INTRODUCTION

Energy-momentum in field theory is a vast subject which can be studied from different viewpoints. The problem lies in the fact that the canonical energy-momentum tensor fails to be a true tensor, while the metric one is appropriate only to theories with a background geometry. In gravitation theories, an energy-momentum flow reduces to a superpotential depending on a world vector field as a gauge parameter.

Analyzing the energy-momentum problem, we follow the general procedure of obtaining differential conservation laws in gauge theory [1-6]. Let $Y \rightarrow X$ be a fibre bundle over a world manifold X coordinated by (x^λ, y^i) , where x^λ are coordinates on X . By gauge transformations are generally meant automorphisms of $Y \rightarrow X$ [7]. To obtain differential conservation laws, it suffices to consider one-parameter groups of gauge transformations whose generators are projectable vector fields

$$u = u^\mu(x^\lambda)\partial_\mu + u^i(x^\lambda, y^j)\partial_i \quad (1)$$

on Y . If the Lie derivative $\mathbf{L}_u L$ of a Lagrangian L vanishes, we have the weak conservation law

$$d_\lambda T^\lambda \approx 0 \quad (2)$$

of the corresponding symmetry flow T^λ along a vector field u . If $u = u^i \partial_i$ is a vertical vector field, (2) is a familiar Nöther conservation law, where T^λ is a Nöther current. This is the case of internal symmetries. If u is a horizontal prolongation on Y of a vector field τ on X , called hereafter a world vector field, we have an energy-momentum conservation law [5, 8, 9]. Of course, different horizontal prolongations of vector fields τ lead to different energy-momentum flows T^λ .

Let us emphasize that the expression (2) is linear in the vector field u . Therefore, one can consider superposition of conservation laws along different vector fields. In particular, every vector field u (1) on Y projected onto a world vector field τ can be seen as the sum $u = u_\tau + v$ of a horizontal prolongation u_τ of τ on Y and a vertical vector field v on Y . It follows that every conservation law (2) can be represented as a superposition of a Nöther conservation law and an energy-momentum conservation law. Conversely, two horizontal prolongations u_τ and u'_τ of the same world vector field τ differ from each other in a vertical vector field $u'_\tau - u_\tau$. Hence, the energy-momentum flows along u_τ and u'_τ differ from each other in a Nöther current. One can not single out in a canonical way the Nöther part of an energy-momentum flow. Therefore, if internal symmetries are broken, an energy-momentum flow is not conserved in general.

In particular, a generic gravitational Lagrangian is invariant under general covariant transformations, but not under vertical gauge transformations of the general linear group $GL(4, \mathbf{R})$. As a consequence, only the energy-momentum flow, corresponding to the canonical horizontal prolongation of vector fields τ on X , i.e., the generators of general covariant transformations, is generally conserved in gravitation theory. This energy-momentum flow possesses the following two important peculiarities.

(i) It reduces to a superpotential, i.e.,

$$T^\lambda \approx d_\mu U^{\mu\lambda}, \quad U^{\mu\lambda} = -U^{\lambda\mu}. \quad (3)$$

We will see that this is a common property of gauge conservation laws if a vector field u depends on the derivatives of gauge parameters. Indeed, the canonical prolongation $\tilde{\tau}$ of a world vector field $\tau = \tau^\lambda \partial_\lambda$ depends on the derivatives $\partial_\mu \tau^\lambda$ of the components of τ which play the role of gauge parameters of infinitesimal general covariant transformations.

(ii) A gravitational superpotential (3) depends on the components τ^λ of a world vector field τ . This is also a common property of gauge conservation laws which make them gauge-covariant, i.e., form-invariant under gauge transformations. Only in the Abelian case of electromagnetic theory, an electric current is free from gauge parameters.

A direct computation shows that General Relativity, Palatini formalism, metric-affine gravitation theory, and gauge gravitation theory in the presence of fermion fields lead

to the same gravitational superpotential. This is the generalized Komar superpotential

$$U^{\mu\lambda}(\tau) = \left(\frac{\partial L}{\partial R_{\mu\lambda}{}^{\alpha}{}_{\nu}} - \frac{\partial L}{\partial R_{\lambda\mu}{}^{\alpha}{}_{\nu}} \right) (D_{\nu}\tau^{\alpha} + S_{\nu}{}^{\alpha}{}_{\sigma}\tau^{\sigma}), \quad (4)$$

where $R_{\mu\lambda}{}^{\alpha}{}_{\nu}$ and $S_{\nu}{}^{\alpha}{}_{\sigma}$ are the curvature and the torsion of a world connection $K_{\nu}{}^{\alpha}{}_{\sigma}$, and D_{ν} is the corresponding covariant derivative. In the particular case of the Hilbert–Einstein Lagrangian L and a Levi–Civita connection K , the superpotential (4) reduces to the familiar Komar superpotential.

Throughout the paper, we follow the convention where the covariant derivative reads

$$D_{\nu}\tau^{\alpha} = \partial_{\nu}\tau^{\alpha} - K_{\nu}{}^{\alpha}{}_{\mu}\tau^{\mu}.$$

II. GAUGE CONSERVATION LAWS

To obtain gauge conservation laws, we follow the first variational formula of Lagrangian formalism.

Given a gauge field system described by sections ϕ of a fibre bundle $Y \rightarrow X$, its space of fields and their first order partial derivatives is the finite-dimensional first order jet manifold J^1Y of Y . Its elements are fields ϕ identified by their values and the values of their first order derivatives at points of X (see, e.g., [9] for a detailed exposition). This space is provided with the adapted coordinates $(x^{\lambda}, y^i, y_{\lambda}^i)$, where y_{λ}^i are the derivative coordinates such that $y_{\lambda}^i \circ \phi = \partial_{\lambda}\phi^i$. The transformation law of the derivative coordinates is

$$y_{\lambda}^i = \left(\frac{\partial y^i}{\partial y^j} y_{\mu}^j + \frac{\partial y^i}{\partial x^{\mu}} \right) \frac{\partial x^{\mu}}{\partial x'^{\lambda}}.$$

A Lagrangian L of a field system on a fibre bundle $Y \rightarrow X$ is defined as a density

$$L = L(x^{\lambda}, y^i, y_{\lambda}^i) d^4x$$

on the space J^1Y . We will use the notation

$$\begin{aligned} \pi_i^{\lambda} &= \partial_i^{\lambda} L, \\ \omega &= d^4x, \quad \omega_{\lambda} = \partial_{\lambda} \rfloor \omega, \quad \omega_{\mu\lambda} = \partial_{\mu} \rfloor \omega_{\lambda}. \end{aligned}$$

Let u be a projectable vector field (1) on a fibre bundle Y , treated as the generator of a one-parameter group of gauge transformations. Its prolongation on the space J^1Y reads

$$J^1u = u^{\lambda}\partial_{\lambda} + u^i\partial_i + (d_{\lambda}u^i - y_{\mu}^i\partial_{\lambda}u^{\mu})\partial_i^{\lambda}$$

where $d_\lambda = \partial_\lambda + y_\lambda^i \partial_i$ denote the total derivatives. The first variational formula provides the canonical decomposition of the Lie derivative

$$\mathbf{L}_u L \omega = [\partial_\lambda u^\lambda L + (u^\lambda \partial_\lambda + u^i \partial_i + (d_\lambda u^i - y_\mu^i \partial_\lambda u^\mu) \partial_i^\lambda) L] \omega \quad (5)$$

of a Lagrangian L in accordance with the variational problem. This decomposition reads

$$\begin{aligned} \partial_\lambda u^\lambda L + [u^\lambda \partial_\lambda + u^i \partial_i + (d_\lambda u^i - y_\mu^i \partial_\lambda u^\mu) \partial_i^\lambda] L = \\ (u^i - y_\mu^i u^\mu) \delta_i L - d_\lambda T^\lambda, \end{aligned} \quad (6)$$

where

$$\delta_i L = (\partial_i - d_\lambda \partial_i^\lambda) L, \quad d_\lambda = \partial_\lambda + y_\lambda^i \partial_i + y_{\lambda\mu}^i \partial_i^\mu, \quad (7)$$

are the variational derivatives and

$$T = T^\lambda \omega_\lambda, \quad T^\lambda = \pi_i^\lambda (u^\mu y_\mu^i - u^i) - u^\lambda L, \quad (8)$$

is said to be a symmetry flow along the vector field u .

It should be emphasized that the flow (8) is defined modulo the terms

$$d_\mu (c_i^{\mu\lambda} (y_\nu^i u^\nu - u^i)),$$

where $c_i^{\mu\lambda}$ are arbitrary skew-symmetric functions on Y [9]. Here, we leave aside these boundary terms which are independent of a Lagrangian, but they may be essential if one examines integral conservation laws.

On the shell

$$\delta_i L = (\partial_i - d_\lambda \partial_i^\lambda) L = 0$$

where solutions of the Euler–Lagrange equations live, the first variational formula (6) leads to the weak identity

$$\begin{aligned} \partial_\lambda u^\lambda L + [u^\lambda \partial_\lambda + u^i \partial_i + (d_\lambda u^i - y_\mu^i \partial_\lambda u^\mu) \partial_i^\lambda] L \approx \\ -d_\lambda [\pi_i^\lambda (u^\mu y_\mu^i - u^i) - u^\lambda L]. \end{aligned} \quad (9)$$

If the Lie derivative $\mathbf{L}_u L$ (5) vanishes, i.e., the Lagrangian L is invariant under the corresponding one-parameter group of gauge transformations, the weak identity (9) is brought into the weak conservation law

$$0 \approx -d_\lambda [\pi_i^\lambda (u^\mu y_\mu^i - u^i) - u^\lambda L], \quad (10)$$

of the flow T (8) along the vector field u .

The weak identity (10) leads to the differential conservation law

$$d_\lambda T^\lambda(\phi) = 0$$

on solutions ϕ of the Euler–Lagrange equations

$$\delta_i L(\phi) = 0.$$

This differential conservation law implies the integral conservation law

$$\int_{\partial M} T^\lambda(\phi) \omega_\lambda = 0, \quad (11)$$

where M is a compact 4-dimensional sub-manifold of X with a boundary ∂M .

We will see that, in gauge and gravitation models, a conserved flow takes the special form

$$T^\lambda = W^\lambda + d_\mu U^{\mu\lambda} \quad (12)$$

where the term W is expressed in the variational derivatives (7), i.e., $W \approx 0$, and T on-shell reduces to a superpotential $U^{\mu\lambda}$ (3). In this case, the integral conservation law (11) becomes tautological, but the superpotential form (12) of T implies the following integral relation

$$\int_{N^3} T^\lambda(\phi) \omega_\lambda = \int_{\partial N^3} U^{\mu\lambda}(\phi) \omega_{\mu\lambda}, \quad (13)$$

where N^3 is a compact oriented 3-dimensional submanifold of X with the boundary ∂N^3 and ϕ is a solution of the Euler–Lagrange equations. One can think of this relation as being a part of the Euler–Lagrange equations written in an integral form.

III. NÖTHER CONSERVATIONS LAWS

Here, we touch on briefly the well-known Nöther conservation laws in gauge theory of internal symmetries in order to emphasize some common properties of gauge conservation laws.

Let $P \rightarrow X$ be a principal bundle with a structure Lie group G . The corresponding gauge model is formulated on the bundle product

$$Y = E \times_X C \quad (14)$$

of a P -associated vector bundle E of matter fields and the bundle $C = J^1 P/G \rightarrow X$ whose sections A are principal connections on the principal bundle P . The bundle C is equipped with the coordinates (x^λ, a_λ^m) such that $a_\lambda^m \circ A = A_\lambda^m$ are gauge potentials.

The bundle Y (14) is coordinated by $(x^\lambda, y^i, a_\lambda^m)$. Let us consider a one-parameter group of vertical automorphisms of the principal bundle P . They yield the corresponding one-parameter group of gauge transformations of the product (14). Its generator is the vertical vector field

$$\xi = (\partial_\mu \xi^r + c_{qp}^r a_\mu^q \xi^p) \partial_r^\mu + \xi^p I_p^i \partial_i,$$

where c_{qp}^r are structure constants of the group G , I_p^i are generators of this group on the typical fibre of $Y \rightarrow X$, and $\xi^p(x^\lambda)$ are gauge parameters, transformed by the coadjoint representation [5, 9, 10]. Let us use the compact notation

$$\begin{aligned} \xi &= (u_p^{A\mu} \partial_\mu \xi^p + u_p^A \xi^p) \partial_A, \\ u_p^{A\mu} \partial_A &= \delta_p^r \partial_r^\mu, \quad u_p^A \partial_A = c_{qp}^r a_\mu^q \partial_r^\mu + I_p^i \partial_i. \end{aligned} \quad (15)$$

If a Lagrangian L is gauge invariant, the first variational formula (6) leads to the strong equality

$$0 = (u_p^A \xi^p + u_p^{A\mu} \partial_\mu \xi^p) \delta_A L + d_\lambda [(u_p^A \xi^p + u_p^{A\mu} \partial_\mu \xi^p) \pi_A^\lambda], \quad (16)$$

where $\delta_A L$ are the variational derivatives of L and

$$d_\lambda = \partial_\lambda + a_{\lambda\mu}^p \partial_p^\mu + y_\lambda^i \partial_i.$$

Due to the arbitrariness of gauge parameters $\xi^p(x^\lambda)$, this equality is equivalent to the following system of strong equalities:

$$u_p^A \delta_A L + d_\mu (u_p^A \pi_A^\mu) = 0, \quad (17a)$$

$$u_p^{A\mu} \delta_A L + d_\lambda (u_p^{A\mu} \pi_A^\lambda) + u_p^A \pi_A^\mu = 0, \quad (17b)$$

$$u_p^{A\lambda} \pi_A^\mu + u_p^{A\mu} \pi_A^\lambda = 0. \quad (17c)$$

Substituting (17b) and (17c) in (17a), we obtain the well-known constraint conditions

$$u_p^A \delta_A L - d_\mu (u_p^{A\mu} \delta_A L) = 0$$

of the variational derivatives of a gauge invariant Lagrangian.

On-shell, the first variational formula (16) leads to the weak conservation law

$$0 \approx d_\lambda [(u_p^A \xi^p + u_p^{A\mu} \partial_\mu \xi^p) \pi_A^\lambda] \quad (18)$$

of the Nöther current

$$T^\lambda = -(u_p^A \xi^p + u_p^{A\mu} \partial_\mu \xi^p) \pi_A^\lambda. \quad (19)$$

Accordingly, the equalities (17a) – (17c) on-shell lead to the equivalent system of Nöther identities

$$d_\mu(u_p^A \pi_A^\mu) \approx 0, \quad (20a)$$

$$d_\lambda(u_p^{A\mu} \pi_A^\lambda) + u_p^A \pi_A^\mu \approx 0, \quad (20b)$$

$$u_p^{A\lambda} \pi_A^\mu + u_p^{A\mu} \pi_A^\lambda = 0 \quad (20c)$$

for a gauge invariant Lagrangian L [2]. They are equivalent to the weak equality (18) due to the arbitrariness of the gauge parameters $\xi^p(x^\lambda)$.

The weak identities (20a) – (20c) play the role of the necessary and sufficient conditions in order that the weak conservation law (18) be gauge-covariant. This means that, if the equality (18) takes place for gauge parameters ξ , it does so for arbitrary deviations $\xi + \delta\xi$ of ξ . Then the conservation law (18) is form-invariant under gauge transformations, when gauge parameters are transformed by the coadjoint representation.

The equalities (20a) – (20c) are not independent, for (20a) is a consequence of (20b) and (20c). This reflects the fact that, in accordance with the strong equalities (17b) and (17c), the Nöther current (19) is brought into the superpotential form

$$T^\lambda = \xi^p u_p^{A\lambda} \delta_A L + d_\mu U^{\mu\lambda}, \quad U^{\mu\lambda} = -\xi^p u_p^{A\mu} \pi_A^\lambda, \quad (21)$$

where the superpotential $U^{\mu\lambda}$ does not depend on matter fields. It is readily observed that the superpotential form of the Nöther current (19) is caused by the fact that the vector fields (15) depend on derivatives of gauge parameters.

On solutions of the Euler–Lagrange equations, we have the corresponding integral relation (13), which reads

$$\int_{N^3} T^\lambda(\phi) \omega_\lambda = \int_{\partial N^3} \xi^p \pi_p^{\mu\lambda}(\phi) \omega_{\mu\lambda}. \quad (22)$$

Since the Nöther superpotential (21) does not contain matter fields, one can think of (22) as being the integral relation between the Nöther current (19) and the gauge field generated by this current. Due to the presence of gauge parameters, the relation (22) is gauge covariant.

In electromagnetic theory, the similar relation between an electric current and the electromagnetic field generated by this current is well known, but it is free from gauge parameters due to the peculiarity of Abelian gauge models. Let us consider electromagnetic theory, where $G = U(1)$ and $I^j(y) = iy^j$. In this case, a gauge parameter ξ is not changed under gauge transformations. Therefore, one can put, e.g., $\xi = 1$. Then the Nöther current (19) takes the form

$$T^\lambda = -u^A \pi_A^\lambda.$$

Since the group G is Abelian, this current does not depend on gauge potentials and it is invariant under gauge transformations. We have

$$T^\lambda = -iy^j \pi_j^\lambda. \quad (23)$$

It is easy to see that T (23), under the sign change, is the familiar electric current of matter fields, while the Nöther conservation law (18) is precisely the equation of continuity. The corresponding integral equation of continuity (11) reads

$$\int_{\partial M} (y^j \pi_j^\lambda)(\phi) \omega_\lambda = 0.$$

Though the Nöther current T (23) takes the superpotential form

$$T^\lambda = -\delta^\lambda L + d_\mu U^{\mu\lambda},$$

the equation of continuity is not tautological. This equation is independent of an electromagnetic field generated by the electric current (23) and it is therefore treated as the strong conservation law of an electric charge. When $\xi = 1$, the electromagnetic superpotential takes the form

$$U^{\mu\lambda} = -\frac{1}{4\pi} F^{\mu\lambda},$$

where F is the electromagnetic strength. Accordingly, the integral equality (22) is the integral form of the Maxwell equations. In particular, the well-known relation between the flux of an electric field through a closed surface and the total electric charge inside this surface is restated.

IV. ENERGY-MOMENTUM IN GRAVITATION THEORIES

From now on by a world manifold X is meant a 4-dimensional orientable noncompact parallelizable manifold. As a consequence, it admits a pseudo-Riemannian metric and a spin structure. Accordingly, a linear connection and a fibre metric on the tangent and cotangent bundles TX and T^*X of X are said to be a world connection and a world metric, respectively.

Gravitation theories are formulated on natural bundles $Y \rightarrow X$, e.g., tensor bundles which admit the canonical horizontal prolongations of any vector field τ on X . These prolongations are the generators of general covariant transformations, where the components of a vector field τ play the role of gauge parameters. By the reason we have explained above, we will investigate the energy-momentum conservation laws associated with these prolongations.

A. Tensor fields

We start from tensor fields which clearly illustrate the main peculiarities of energy-momentum conservation laws on natural bundles. Let

$$Y = (\bigotimes^m TX) \otimes_X (\bigotimes^k T^*X) \quad (24)$$

be a tensor bundle equipped with the holonomic coordinates $(x^\lambda, \dot{x}_{\beta_1 \dots \beta_k}^{\alpha_1 \dots \alpha_m})$.

Given a vector field $\tau = \tau^\lambda \partial_\lambda$, we have its canonical prolongation

$$\tilde{\tau} = \tau^\mu \partial_\mu + [\partial_\nu \tau^{\alpha_1} \dot{x}_{\beta_1 \dots \beta_k}^{\nu \alpha_2 \dots \alpha_m} + \dots - \partial_{\beta_1} \tau^\nu \dot{x}_{\nu \beta_2 \dots \beta_k}^{\alpha_1 \dots \alpha_m} - \dots] \frac{\partial}{\partial \dot{x}_{\beta_1 \dots \beta_k}^{\alpha_1 \dots \alpha_m}} \quad (25)$$

on the tensor bundle (24) and, in particular, its prolongations

$$\tilde{\tau} = \tau^\mu \partial_\mu + \partial_\nu \tau^\alpha \dot{x}^\nu \frac{\partial}{\partial \dot{x}^\alpha} \quad (26)$$

on the tangent bundle TX and

$$\tilde{\tau} = \tau^\mu \partial_\mu - \partial_\beta \tau^\nu \dot{x}_\nu \frac{\partial}{\partial \dot{x}_\beta}$$

on the cotangent bundle T^*X .

Of course, one can consider the horizontal prolongation

$$\tau_K = \tau^\lambda (\partial_\lambda + K_\lambda^\beta{}_\alpha \dot{x}^\alpha \frac{\partial}{\partial \dot{x}^\beta}) \quad (27)$$

of a world vector field τ on TX and tensor bundles by means of any world connection K . This is the generator of a 1-parameter group of nonholonomic automorphisms of these bundles. These automorphisms are met with in gauge theory of the general linear group $GL(4, \mathbf{R})$ [3], but a generic gravitational Lagrangian is not invariant under these transformations. Note that the prolongations (26) and (27) were treated as generators of the gauge group of translations in the pioneer gauge gravitation models (see [11, 12] for a survey).

Let us use the compact notation $y^A = \dot{x}_{\beta_1 \dots \beta_k}^{\alpha_1 \dots \alpha_m}$ such that the canonical prolongation $\tilde{\tau}$ (25) reads

$$\tilde{\tau} = \tau^\lambda \partial_\lambda + u^{A\beta}{}_\alpha \partial_\beta \tau^\alpha \partial_A. \quad (28)$$

This expression is the general form of the canonical prolongation of a world vector field τ on a natural bundle Y , when this prolongation depends only on the first order partial

derivatives of the components of τ . Therefore, the results obtained below for tensor fields are also true for every such natural bundle Y .

Let a Lagrangian L of tensor fields be invariant under general covariant transformations, i.e., its Lie derivative (5) vanishes for any world vector field τ . Then we have the strong equality

$$\partial_\alpha(\tau^\alpha L) + u^{A\beta}_\alpha \partial_\beta \tau^\alpha \partial_A L + d_\mu(u^{A\beta}_\alpha \partial_\beta \tau^\alpha) \pi_A^\mu - y^A_\alpha \partial_\beta \tau^\alpha \pi_A^\beta = 0. \quad (29)$$

The corresponding weak identity (10) takes the form

$$0 \approx -d_\lambda[\pi_A^\lambda(y^A_\alpha \tau^\alpha - u^{A\beta}_\alpha \partial_\beta \tau^\alpha) - \tau^\lambda L]. \quad (30)$$

Due to the arbitrariness of the gauge parameters τ^α , the equality (29) is equivalent to the system of strong equalities

$$\partial_\lambda L = 0, \quad (31a)$$

$$\delta_\alpha^\beta L + u^{A\beta}_\alpha \delta_A L + d_\mu(u^{A\beta}_\alpha \pi_A^\mu) = y^A_\alpha \pi_A^\beta, \quad (31b)$$

$$u^{A\beta}_\alpha \pi_A^\mu + u^{A\mu}_\alpha \pi_A^\beta = 0, \quad (31c)$$

where $\delta_A L$ are the variational derivatives.

Substituting the relations (31b) and (31c) in (30), we obtain the energy-momentum conservation law

$$0 \approx -d_\lambda[u^{A\lambda}_\alpha \delta_A L \tau^\alpha + d_\mu(u^{A\lambda}_\alpha \pi_A^\mu \tau^\alpha)]. \quad (32)$$

A glance at this expression shows that, on-shell, the corresponding energy-momentum flow reduces to a superpotential, i.e.,

$$T^\lambda = \tau^\alpha u^{A\lambda}_\alpha \delta_A L + d_\mu U^{\mu\lambda}, \quad U^{\mu\lambda} = u^{A\lambda}_\alpha \pi_A^\mu \tau^\alpha. \quad (33)$$

It is readily seen that the energy-momentum superpotential (33) emerges from the dependence of the canonical prolongation $\tilde{\tau}$ (28) on the derivatives of the components of the vector field τ . This dependence guarantees that the energy-momentum conservation law (32) is maintained under general covariant transformations.

B. General Relativity

A pseudo-Riemannian metric g on a world manifold X is represented by a section of the quotient

$$\Sigma = LX/O(1, 3) \rightarrow X, \quad (34)$$

where LX is the bundle of linear frames in the tangent spaces to X . It is called the metric bundle. The linear frame bundle LX is a principal bundle with the structure

group $GL(4, \mathbf{R})$. For the sake of simplicity, we will identify the metric bundle with a sub-bundle of the tensor bundle

$$\Sigma \subset \overset{2}{\vee} TX, \quad (35)$$

coordinated by $(x^\lambda, g^{\mu\nu})$. In General Relativity, it is more convenient to consider the metric bundle as a sub-bundle of the tensor bundle

$$\Sigma \subset \overset{2}{\vee} T^*X \quad (36)$$

equipped with the coordinates $(x^\lambda, g_{\mu\nu})$.

The second order Hilbert–Einstein Lagrangian L_{HE} of General Relativity is defined on the second order jet manifold $J^2\Sigma$ of Σ coordinated by $(x^\lambda, g_{\alpha\beta}, g_{\lambda\alpha\beta}, g_{\mu\lambda\alpha\beta})$. It reads

$$L_{HE} = -\frac{1}{2\kappa} g^{\alpha\nu} g^{\beta\mu} R_{\mu\nu\alpha\beta} \sqrt{-g} \omega, \quad g = \det(g_{\mu\nu}), \quad (37)$$

where

$$\begin{aligned} R_{\mu\nu\alpha\beta} &= -\left[\frac{1}{2}(g_{\mu\beta\alpha\nu} + g_{\nu\alpha\beta\mu} - g_{\nu\beta\alpha\mu} - g_{\mu\alpha\beta\nu}) + \right. \\ &\quad \left. g_{\varepsilon\gamma}(\{_\mu^\varepsilon_\beta\}\{_\nu^\gamma_\alpha\} - \{_\nu^\varepsilon_\beta\}\{_\mu^\gamma_\alpha\})\right], \\ \{_\nu^\alpha_\mu\} &= -\frac{1}{2}g^{\alpha\beta}(g_{\nu\beta\mu} + g_{\mu\beta\nu} - g_{\beta\mu\nu}). \end{aligned}$$

Let τ be a vector field on X and

$$\tilde{\tau} = \tau^\lambda \partial_\lambda - (g_{\nu\beta} \partial_\alpha \tau^\nu + g_{\alpha\nu} \partial_\beta \tau^\nu) \partial^{\alpha\beta} \quad (38)$$

its canonical prolongation (25) on the metric bundle Σ (36). Since the Lagrangian L_{HE} (37) is invariant under general covariant transformations, its Lie derivative along the vector field $\tilde{\tau}$ (38) vanishes. Then the first variational formula for second order Lagrangians [9, 13] leads to the energy-momentum conservation law

$$\begin{aligned} 0 &\approx -d_\lambda \{2g_{\mu\alpha} \tau^\mu \delta^{\alpha\lambda} L_{HE} + d_\mu [\frac{1}{2\kappa} \sqrt{-g} (g^{\lambda\nu} \nabla_\nu \tau^\mu - g^{\mu\nu} \nabla_\nu \tau^\lambda)]\}, \\ \nabla_\nu \tau^\mu &= \partial_\nu \tau^\mu - \{_\nu^\mu_\alpha\} \tau^\alpha. \end{aligned}$$

A glance at this conservation law shows that, on-shell, the energy-momentum flow reduces to the well-known Komar superpotential [14]:

$$U^{\mu\lambda} = \frac{1}{2\kappa} \sqrt{|g|} (g^{\lambda\nu} \nabla_\nu \tau^\mu - g^{\mu\nu} \nabla_\nu \tau^\lambda). \quad (39)$$

C. Metric-affine gravitation theory

In metric-affine gravitation theory, gravity is described by a pseudo-Riemannian metric g and a world connection K on X . Since world connections are associated with principal connections on the linear frame bundle LX , there is one-to-one correspondence between the world connections and the sections of the quotient

$$C_K = J^1 LX / GL(4, \mathbf{R}) \rightarrow X. \quad (40)$$

This bundle is provided with the coordinates $(x^\lambda, k_\lambda^\nu)$ so that, for any section K of $C_K \rightarrow X$,

$$k_\lambda^\nu \circ K = K_\lambda^\nu$$

are the coefficients of the world connection K . The bundle C_K (40) admits the canonical horizontal prolongation

$$\tilde{\tau}_K = \tau^\mu \partial_\mu + [\partial_\nu \tau^\alpha k_\mu^\nu{}_\beta - \partial_\beta \tau^\nu k_\mu^\alpha{}_\nu - \partial_\mu \tau^\nu k_\nu^\alpha{}_\beta + \partial_{\mu\beta} \tau^\alpha] \frac{\partial}{\partial k_\mu^\alpha{}_\beta} \quad (41)$$

of vector fields τ on X . We will use the compact notation

$$\tilde{\tau}_K = \tau^\lambda \partial_\lambda + (u^{A\varepsilon}_\gamma \partial_\varepsilon \tau^\gamma + u^{A\varepsilon\sigma}_\gamma \partial_{\varepsilon\sigma} \tau^\gamma) \partial_A,$$

where

$$\begin{aligned} y^A &= k_\mu^\alpha{}_\beta, \\ u_\mu^\alpha{}_\beta{}^\varepsilon &= k_\mu^\varepsilon{}_\beta \delta_\gamma^\alpha - k_\mu^\alpha{}_\gamma \delta_\beta^\varepsilon - k_\gamma^\alpha{}_\beta \delta_\mu^\varepsilon, \\ u_\mu^\alpha{}_\beta{}^{\varepsilon\sigma} &= \delta_\mu^\varepsilon \delta_\beta^\sigma \delta_\gamma^\alpha. \end{aligned} \quad (42)$$

Metric-affine gravitation theory is formulated on the bundle product

$$Y = \Sigma \times_X C_K, \quad (43)$$

coordinated by $(x^\lambda, g^{\alpha\beta}, k_\mu^\alpha{}_\beta)$, where Σ is the sub-bundle (35). The corresponding space $J^1 Y$ is equipped with the coordinates

$$(x^\lambda, g^{\alpha\beta}, k_\mu^\alpha{}_\beta, g_{\lambda}^{\alpha\beta}, k_{\lambda\mu}^\alpha{}_\beta).$$

We will assume that a metric-affine Lagrangian L_{MA} factorizes through the curvature

$$R_{\lambda\mu}^\alpha{}_\beta = k_{\lambda\mu}^\alpha{}_\beta - k_{\mu\lambda}^\alpha{}_\beta + k_\mu^\alpha{}_\varepsilon k_\lambda^\varepsilon{}_\beta - k_\lambda^\alpha{}_\varepsilon k_\mu^\varepsilon{}_\beta,$$

and does not depend on the derivative coordinates $g_{\lambda}^{\alpha\beta}$ of a world metric. Then the following relations take place:

$$\pi^{\lambda\nu}{}_{\alpha}{}^{\beta} = -\pi^{\nu\lambda}{}_{\alpha}{}^{\beta}, \quad (44)$$

$$\frac{\partial L_{MA}}{\partial k_{\nu}{}^{\alpha}{}_{\beta}} = \pi^{\lambda\nu}{}_{\alpha}{}^{\sigma} k_{\lambda}{}^{\beta}{}_{\sigma} - \pi^{\lambda\nu}{}_{\sigma}{}^{\beta} k_{\lambda}{}^{\sigma}{}_{\alpha}. \quad (45)$$

We also have the equalities

$$\pi_A^{\lambda} u^{\alpha\beta\mu}_{\alpha} = \pi^{\lambda\mu}{}_{\alpha}{}^{\beta}, \quad \pi_A^{\varepsilon} u^{\alpha\beta}_{\alpha} = -\partial^{\varepsilon}{}_{\alpha}{}^{\beta} L_{MA} - \pi^{\varepsilon\beta}{}_{\sigma}{}^{\gamma} k_{\alpha}{}^{\sigma}{}_{\gamma}.$$

Given a vector field τ on a world manifold X , its canonical prolongation on the product (43) reads

$$\begin{aligned} \tilde{\tau} = & \tau^{\lambda} \partial_{\lambda} + (g^{\nu\beta} \partial_{\nu} \tau^{\alpha} + g^{\alpha\nu} \partial_{\nu} \tau^{\beta}) \partial_{\alpha\beta} + \\ & (u^{\alpha\beta}_{\alpha} \partial_{\beta} \tau^{\alpha} + u^{\alpha\beta\mu}_{\alpha} \partial_{\beta\mu} \tau^{\alpha}) \partial_A. \end{aligned}$$

Let a metric-affine Lagrangian L_{MA} be invariant under general covariant transformations, i.e.,

$$\mathbf{L}_{\tilde{\tau}} L_{MA} = 0 \quad (46)$$

for any world vector field τ . Then, on-shell, the first variational formula (6) leads to the weak conservation law

$$0 \approx -d_{\lambda} [\pi_A^{\lambda} (y_A^{\alpha} \tau^{\alpha} - u^{\alpha\beta}_{\alpha} \partial_{\beta} \tau^{\alpha} - u^{\alpha\beta\gamma}_{\alpha} \partial_{\beta\gamma} \tau^{\alpha}) - \tau^{\lambda} L_{MA}] \quad (47)$$

where

$$T^{\lambda} = \pi_A^{\lambda} (y_A^{\alpha} \tau^{\alpha} - u^{\alpha\beta}_{\alpha} \partial_{\beta} \tau^{\alpha} - u^{\alpha\beta\gamma}_{\alpha} \partial_{\beta\gamma} \tau^{\alpha}) - \tau^{\lambda} L_{MA} \quad (48)$$

is the energy-momentum flow of the metric-affine gravity.

It is readily observed that, in the local gauge where the vector field τ is constant, the energy-momentum flow (48) leads to the canonical energy-momentum tensor

$$T^{\lambda} = (\pi^{\lambda\mu}{}_{\beta}{}^{\nu} k_{\alpha\mu}{}^{\beta}{}_{\nu} - \delta_{\alpha}^{\lambda} L_{MA}) \tau^{\alpha}.$$

This tensor was suggested in order to describe the energy-momentum complex in the Palatini model [15].

Due to the arbitrariness of the gauge parameters τ^{λ} , the equality (46) is equivalent to the system of strong equalities

$$\begin{aligned} \partial_{\lambda} L &= 0, \\ \delta_{\alpha}^{\beta} L_{MA} + 2g^{\beta\mu} \delta_{\alpha\mu} L_{MA} + u^{\alpha\beta}_{\alpha} \delta_A L_{MA} + d_{\mu} (\pi_A^{\mu} u^{\alpha\beta}_{\alpha}) - y_{\alpha}^A \pi_A^{\beta} &= 0, \end{aligned} \quad (49)$$

$$(u^{\alpha\beta\gamma}_{\gamma} \partial_A + u^{\alpha\beta}_{\gamma} \partial_A^{\sigma}) L_{MA} \partial_{\sigma\varepsilon} \tau^{\gamma} = 0, \quad (50)$$

$$\pi^{(\lambda\varepsilon}{}_{\gamma}{}^{\sigma)} = 0, \quad (51)$$

where $\delta_{\alpha\mu}L_{MA}$ and δ_AL_{MA} are the corresponding variational derivatives. It is readily observed that the equality (50) holds owing to the relation (45), while the equality (51) does due to the relation (44).

Substituting the term $y_\alpha^A\pi_A^\beta$ from the expression (49) in the energy-momentum conservation law (47), we bring this conservation law into the form

$$0 \approx -d_\lambda[2g^{\lambda\mu}\tau^\alpha\delta_{\alpha\mu}L_{MA} + u^{A\lambda}_\alpha\tau^\alpha\delta_AL_{MA} - \pi_A^\lambda u^{A\beta}_\alpha\partial_\beta\tau^\alpha + d_\mu(\pi^{\lambda\mu}_\alpha{}^\beta)\partial_\beta\tau^\alpha + d_\mu(\pi_A^\mu u^{A\lambda}_\alpha\tau^\alpha - d_\mu(\pi^{\lambda\mu}_\alpha{}^\beta\partial_\beta\tau^\alpha)]. \quad (52)$$

After separating the variational derivatives, the energy-momentum conservation law (52) of the metric-affine gravity takes the superpotential form

$$0 \approx -d_\lambda[2g^{\lambda\mu}\tau^\alpha\delta_{\alpha\mu}L_{MA} + (k_\mu{}^\lambda{}_\gamma\delta^\mu{}_\alpha{}^\gamma L_{MA} - k_\mu{}^\sigma{}_\alpha\delta^\mu{}_\sigma{}^\lambda L_{MA} - k_\alpha{}^\sigma{}_\gamma\delta^\lambda{}_\sigma{}^\gamma L_{MA})\tau^\alpha + \delta^\lambda{}_\alpha{}^\mu L_{MA}\partial_\mu\tau^\alpha - d_\mu(\delta^\mu{}_\alpha{}^\lambda L_{MA})\tau^\alpha + d_\mu(\pi^{\mu\lambda}_\alpha{}^\nu(\partial_\nu\tau^\alpha - k_\sigma{}^\alpha{}_\nu\tau^\sigma))],$$

where the energy-momentum flow on-shell reduces to the generalized Komar superpotential

$$U^{\mu\lambda} = \frac{\partial L_{MA}}{k_{\mu\lambda}{}^\alpha{}_\nu}(\partial_\nu\tau^\alpha - k_\sigma{}^\alpha{}_\nu\tau^\sigma) \quad (53)$$

that we have written in the form (4) [16].

In particular, let us consider the Hilbert–Einstein Lagrangian density

$$L_{HE} = -\frac{1}{2\kappa}R\sqrt{-g}\omega, \quad R = g^{\lambda\nu}R_{\lambda\alpha}{}^\alpha{}_\nu,$$

in the framework of metric-affine gravitation theory. Then the generalized Komar superpotential (4) comes to the Komar superpotential (39) if we substitute the Levi–Civita connection $k_\nu{}^\alpha{}_\sigma = \{\nu{}^\alpha{}_\sigma\}$. One may generalize this example by considering the Lagrangian

$$L = f(R)\sqrt{-g}\omega,$$

where $f(R)$ is a polynomial of the scalar curvature R . In the case of a symmetric connection, we restate the superpotential

$$U^{\mu\lambda} = -\frac{\partial f}{\partial R}\sqrt{-g}(g^{\lambda\nu}D_\nu\tau^\mu - g^{\mu\nu}D_\nu\tau^\lambda)$$

of the Palatini model [17] as like as the superpotential in the recent work [18] where Lagrangians of the Palatini model factorize through the product $R^{\alpha\beta}R_{\alpha\beta}$.

D. Gauge gravitation theory

Turning to the energy-momentum conservation law in gauge gravitation theory, we meet with the problem that spinor bundles over a world manifold do not admit general covariant transformations. This difficulty can be overcome as follows [9, 19]. The linear frame bundle LX is the principal bundle $LX \rightarrow \Sigma$ over the metric bundle Σ (34) whose structure group is the Lorentz group. Let us consider a spinor bundle $S \rightarrow \Sigma$ associated with the Lorentz bundle $LX \rightarrow \Sigma$. For each gravitational field g , the restriction of this spinor bundle to $g(X) \subset \Sigma$ is isomorphic to the spinor bundle $S^g \rightarrow X$ whose sections describe Dirac fermion fields in the presence of a gravitational field g . Moreover, the spinor bundle $S \rightarrow \Sigma$ can be provided with the Dirac operator and the Dirac Lagrangian whose restrictions to $g(X)$ are the familiar Dirac operator and Dirac Lagrangian of fermion fields in the presence of a background gravitational field g . It follows that sections of the fibre bundle $S \rightarrow X$ describe the total system of fermion and gravitational fields on a world manifold X . At the same time, the fibre bundle $S \rightarrow X$ is not a spinor bundle, and it inherits general covariant transformations of the frame bundle LX . The corresponding horizontal prolongation on S of world vector fields on X can be constructed.

As a consequence, gauge gravitation theory of metric gravitational fields, world connections and Dirac fermion fields can be formulated on the bundle product

$$Y = C_K \times_{\Sigma} S, \quad (54)$$

coordinated by $(x^\mu, h_a^\mu, k_\mu^\alpha{}_\beta, \psi^A)$, where h_a^μ are tetrad coordinates on Σ , i.e.,

$$g^{\mu\nu} = h_a^\mu h_b^\nu \eta^{ab}$$

where η is the Minkowski metric. The corresponding space $J^1 Y$ is provided with the adapted coordinates

$$(x^\mu, h_a^\mu, k_\mu^\alpha{}_\beta, \psi^A, h_{\lambda a}^\mu, k_{\lambda \mu}^\alpha{}_\beta, \psi_\lambda^A).$$

The total Lagrangian on this space is the sum

$$L = L_{MA} + L_D \quad (55)$$

of the metric-affine Lagrangian L_{MA} in the previous Section and the Dirac Lagrangian

$$\begin{aligned} L_D = & \left\{ \frac{i}{2} h_q^\lambda [\psi_A^+ (\gamma^0 \gamma^q)^A{}_B (\psi_\lambda^B - \frac{1}{4} (\eta^{kb} h_\mu^a - \eta^{ka} h_\mu^b) (h_{\lambda k}^\mu - h_k^\nu k_\lambda^\mu{}_\nu) L_{ab}{}^B{}_C \psi^C) - \right. \\ & \left. (\psi_{\lambda A}^+ - \frac{1}{4} (\eta^{kb} h_\mu^a - \eta^{ka} h_\mu^b) (h_{\lambda k}^\mu - h_k^\nu k_\lambda^\mu{}_\nu) \psi_C^+ L_{ab}{}^{+C}{}_A) (\gamma^0 \gamma^q)^A{}_B \psi^B] - \right. \\ & \left. m \psi_A^+ (\gamma^0)^A{}_B \psi^B \right\} \sqrt{-g}, \quad L_{ab} = \frac{1}{4} [\gamma_a, \gamma_b]. \end{aligned}$$

Note that, in fact, the Dirac Lagrangian L_D depends only on the torsion $k_\lambda^\mu{}_\nu - k_\nu^\mu{}_\lambda$ of a world connection, while the pseudo-Riemannian part is given by the derivative coordinates $h_{\lambda k}^\mu$.

Given a vector field τ on a world manifold X , its horizontal prolongation on the product (54) is

$$\tilde{\tau}_Y = \tilde{\tau} + v, \quad (56)$$

$$\tilde{\tau} = \tilde{\tau}_K + \partial_\nu \tau^\mu h_c^\nu \frac{\partial}{\partial h_c^\mu},$$

$$v = \frac{1}{4} Q_k^\mu (\eta^{kb} h_\mu^a - \eta^{ka} h_\mu^b) [-L_{ab}{}^d{}_c h_d^\nu \frac{\partial}{\partial h_c^\nu} + L_{ab}{}^A{}_B \psi^B \partial_A + L_{ab}^{+A}{}_B \psi_A^+ \partial^B],$$

where $\tilde{\tau}_K$ is the vector field (41), $L_{ab}{}^d{}_c$ are generators of the Lorentz group in the Minkowski space, and the terms Q_c^μ obey the condition

$$(Q_a^\mu h_b^\nu + Q_a^\nu h_b^\mu) \eta^{ab} = 0$$

(see [9, 19] for a detailed exposition). The horizontal part $\tilde{\tau}$ of the vector field (56) is the generator of a one-parameter group of general covariant transformations of the fibre bundle (54), whereas the vertical one v is the generator of a one-parameter group of vertical Lorentz automorphisms of the spinor bundle $S \rightarrow \Sigma$. By construction, the total Lagrangian L (55) obeys the relations

$$\mathbf{L}_v L_D = 0. \quad (57)$$

$$\mathbf{L}_{\tilde{\tau}} L_{MA} = 0, \quad \mathbf{L}_{\tilde{\tau}} L_D = 0. \quad (58)$$

The relation (57) results in the Nöther conservation law, while the equalities (58) lead to the energy-momentum one [4, 6].

Using the compact notation (42), let us rewrite the horizontal part $\tilde{\tau}$ of the vector field (56) in the form

$$\tilde{\tau} = \tau^\mu \partial_\mu + \partial_\nu \tau^\mu h_a^\nu \frac{\partial}{\partial h_a^\mu} + (u^{A\beta}{}_\alpha \partial_\beta \tau^\alpha + u^{A\varepsilon\beta}{}_\alpha \partial_{\varepsilon\beta} \tau^\alpha) \partial_A.$$

Due to the arbitrariness of the functions τ^α , the conditions (58) lead to the strong equalities

$$\delta_\alpha^\beta L_{MA} + 2h^{\beta\mu} \delta_{\alpha\mu} L_{MA} + u^{A\beta}{}_\alpha \delta_A L_{MA} + d_\mu (\pi_A^\mu u^{A\beta}{}_\alpha) = y_\alpha^A \partial_A^\beta L_{MA}, \quad (59)$$

$$\delta_\alpha^\beta L_D + \sqrt{-g} t_\alpha^\beta + \frac{\partial L_D}{\partial h_{\lambda c}^\alpha} h_{\lambda c}^\beta + \partial_A L_D u_\alpha^{A\beta} = \quad (60)$$

$$\frac{\partial L_D}{\partial h_{\beta c}^\mu} h_{\alpha c}^\mu + \frac{\partial L_D}{\partial \psi_\beta^A} \psi_\alpha^A + \frac{\partial L_D}{\partial \psi_{\beta A}^+} \psi_{\alpha A}^+,$$

where

$$\sqrt{-g}t_\alpha^\beta = h_a^\beta \frac{\partial L_D}{\partial h_a^\alpha}$$

is the metric energy-momentum tensor of fermion fields. We also have the relations (44), (45) and

$$\frac{\partial L_D}{\partial k_{\lambda}{}^\mu{}_\nu} = -\frac{\partial L_D}{\partial k_\nu{}^\mu{}_\lambda} = \frac{\partial L_D}{\partial h_{\lambda c}^\mu} h_c^\nu. \quad (61)$$

The corresponding energy-momentum conservation law reads

$$0 \approx -d_\lambda [\partial_A^\lambda L_{MA} (y_\alpha^A \tau^\alpha - u^{A\beta}_\alpha \partial_\beta \tau^\alpha - u^{A\varepsilon\beta}_\alpha \partial_{\varepsilon\beta} \tau^\alpha) - \frac{\partial L_D}{\partial h_{\lambda c}^\alpha} (\partial_\beta \tau^\alpha h_c^\beta - h_{\mu c}^\alpha \tau^\mu) + \frac{\partial L_D}{\partial \psi_\lambda^A} \psi_\alpha^A \tau^\alpha + \frac{\partial L_D}{\partial \psi_{\lambda A}^+} \psi_{\alpha A}^+ \tau^\alpha - \tau^\lambda L]. \quad (62)$$

Substituting the term $y_\alpha^A \partial_A^\beta L_{MA}$ from (59) and the term

$$\frac{\partial L_D}{\partial h_{\beta c}^\mu} h_{\alpha c}^\mu + \frac{\partial L_D}{\partial \psi_\beta^A} \psi_\alpha^A + \frac{\partial L_D}{\partial \psi_{\beta A}^+} \psi_{\alpha A}^+$$

from (60) in (62), we bring this conservation law into the superpotential form

$$0 \approx -d_\lambda [2h^{\lambda\mu} \tau^\alpha \delta_{\alpha\mu} L + (k_\mu{}^\lambda{}_\gamma \delta^\mu{}_\alpha{}^\gamma L - k_\mu{}^\sigma{}_\alpha \delta^\mu{}_\sigma{}^\lambda L - k_\alpha{}^\sigma{}_\gamma \delta^\lambda{}_\sigma{}^\gamma L) \tau^\alpha + \delta^\lambda{}_\alpha{}^\mu L \partial_\mu \tau^\alpha - d_\mu (\delta^\mu{}_\alpha{}^\lambda L) \tau^\alpha + d_\mu (\pi^{\mu\lambda}{}_\alpha{}^\nu (\partial_\nu \tau^\alpha - k_\sigma{}^\alpha{}_\nu \tau^\sigma))] - d_\lambda \left[\left(\frac{\partial L_D}{\partial h_{\mu a}^\alpha} h_a^\lambda + \frac{\partial L_D}{\partial h_{\lambda a}^\alpha} h_a^\mu \right) \partial_\mu \tau^\alpha \right]. \quad (63)$$

By virtue of the relations (61), the last term in the expression (63) vanishes, i.e., fermion fields do not contribute to the superpotential. It follows that the energy-momentum conservation law (62) of gauge gravitation theory takes the superpotential form (12), where $U^{\mu\lambda}$ is the generalized Komar superpotential (53).

V. CONCLUSIONS

The above energy-momentum conservation laws in gravitation theories are derived from the condition of invariance of gravitational Lagrangians under general covariant transformations. We have shown that they possess the generic properties of gauge conservation laws. Since the generators of general covariant transformations depend on the derivatives of gauge parameters, i.e., components of a world vector field τ , the corresponding energy-momentum flow T^λ takes the superpotential form. As in electromagnetic theory, one can write the integral relation (13) on-shell between the energy-momentum flow T^λ and the gravitational superpotential $U^{\mu\lambda}$. The examples of fermion

fields and Proca fields [9] show that the superpotential $U^{\mu\lambda}$ does not depend on matter fields. Then, by analogy with electromagnetic and gauge theories, one can think of (13) as being the relation between the energy-momentum flow T^λ in a compact 3-dimensional manifold $N \subset X$ and the flux $U^{\mu\lambda}$ of a gravitational field generated by this flow through the boundary ∂N .

The peculiarity of gravitation theory as like as any non-Abelian gauge theory lies in the fact that the integral relation (22) depends on a gauge parameter. In the case of general covariant transformations, this gauge parameter is a world vector field τ whose canonical prolongation on a natural bundle is the generator of general covariant transformations. As a consequence, the energy-momentum conservation law in gravitation theories as like as in mechanics [20] depends on a reference frame, but it is form-invariant under general covariant transformations. In particular, with respect to the local gauge $\tau = \text{const.}$, the energy-momentum flow in the conservation law (62) is the canonical energy-momentum tensor.

A direct computation also shows that, for a number of gravitation models, the generalized Komar superpotential (4) appears to be an energy-momentum superpotential associated with gauge invariance under general covariant transformations.

References

- [1] J.Goldberg, in *General Relativity and Gravitation*, edited by A.Held (Plenum Press, N.Y., 1980); D.Bak, D.Cangemi and R.Jackiw, Phys. Rev. D **49**, 5173 (1994); B. van der Heuvel, J. Math. Phys. **35**, 1668 (1994).
- [2] N.Konopleva and V.Popov, *Gauge Fields* (Harwood Academic Publishers, London, 1981).
- [3] F.Hehl, J.McCrea, E.Mielke and Y.Ne'eman, Phys. Rep. **258**, 1 (1995).
- [4] G.Giachetta and L.Mangiarotti, Int. J. Theor. Phys. **36**, 125 (1997).
- [5] G.Sardanashvily, J. Math. Phys. **38**, 847 (1997).
- [6] G.Sardanashvily, Class. Quant. Grav. **14**, 1371 (1997).
- [7] M.Mayer, in *Lecture Notes in Physics* **67** (Springer-Verlag, Berlin, 1977).
- [8] M.Gotay and J.Marsden, Contemp. Math. **132**, 367 (1992).
- [9] G.Giachetta, L.Mangiarotti and G.Sardanashvily, *New Lagrangian and Hamiltonian Methods in Field Theory* (World Scientific, Singapore, 1997).

- [10] G.Giachetta and L.Mangiarotti, Int. J. Theor. Phys. **29**, 789 (1990).
- [11] F.Hehl, P. von der Heyde, G.Kerlick and J.Nester, Rev. Mod. Phys. **48**, 393 (1976).
- [12] D. Ivanenko and G. Sardanashvily, Phys. Rep. **94**, 1 (1983).
- [13] J.Novotný, in *Geometrical Methods in Physics, Proceeding of the Conference on Differential Geometry and its Applications (Czechoslovakia 1983)*, edited by D.Krupka (University of J.E.Purkyně, Brno, 1984).
- [14] A.Komar, *Phys. Rev.* **113**, 934 (1959).
- [15] G.Murphy, Int. J. Theor. Phys. **29** 1003 (1990); R.Dick, Int. J. Theor. Phys. **32** 109, (1993) 109; J.Novotný, Int. J. Theor. Phys. **32** 1033 (1993).
- [16] G.Giachetta and G.Sardanashvily, Class. Quant. Grav. **13**, L67 (1996).
- [17] A.Borowiec, M.Ferraris, M.Francaviglia and I.Volovich, GRG **26**, 637 (1994).
- [18] A.Borowiec, M.Ferraris, M.Francaviglia and I.Volovich, Class. Quant. Grav **15**, 43 (1998).
- [19] G.Sardanashvily, Int. J. Theor. Phys **37**, 1257 (1998); J. Math. Phys. **39** N9 (1998).
- [20] A.Echeverría Enríquez, M.Muñoz Lecanda and N.Román Roy, J. Phys.A **28**, 5553 (1995); G.Sardanashvily, J. Math. Phys. **39**, 2714 (1998); L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).