

A Stationary Asymptotically Einstein Static Universe Solution

Mihaela Time

7th September 1998

Abstract

Here I present a stationary cylindrically symmetric asymptotically Einstein static universe solution with the matter consisting of a cosmological and rotating dust term which admits predicted black hole event horizon.

1 Introduction

I will consider here the strong gravitational fields in some form of gravitational collapse, when the gravitational forces cannot be compensated by other forces. As a special case I introduce an asymptotically Einstein static universe space-time. This space-time is far from the strong gravitational field and the metric has the form of the Einstein static universe at infinity.

The spatially closed, static Einstein universe,

$$(1) \quad ds_E^2 = d\eta^2 + \sin^2 \eta (d\theta^2 + \sin^2 \theta d\varphi^2) - c^2 d\psi^2$$

$$\varphi \in [0, 2\pi], \quad \eta \in [0, \pi], \quad \theta \in [0, \pi], \quad \psi \in \mathbb{R}$$

is the simplest cosmological dust model with constant curvature $K = \text{const.}$; the energy-momentum tensor T_{ab} has the form (15), where $\Lambda = \frac{1}{K^2}$ and $\mu = \frac{2}{K^2} = 2\Lambda$.

The space-time of Special Relativity is described mathematically by the Minkowski space (M, η) . The flat metric η , in coordinates ¹

$$(2) \quad ds^2 = dr^2 + r^2(d\theta^2 + \sin^2 \theta d\varphi^2) - c^2 dt^2$$

$$r \in [0, \infty), \quad \theta \in [0, \pi], \quad \varphi \in [0, 2\pi], \quad t \in \mathbb{R},$$

¹The metric is apparently singular for $r = 0$ and $\sin \theta = 0$.

is wholly conformal to a finite region of the Einstein static universe (1).

We have:

$$(3) \quad ds^2 = \Theta^{-2} ds_E^2,$$

where $\Theta = 2 \cos \frac{\psi+\eta}{2} \cos \frac{\psi-\eta}{2}$ is a smooth strictly positive function, under the conformal transformation:

$$t + r = \operatorname{tg} \frac{\psi + \eta}{2}, \quad t - r = \operatorname{tg} \frac{\psi - \eta}{2}, \quad -\frac{\pi}{2} \leq \psi - \eta \leq \psi + \eta \leq \frac{\pi}{2}$$

(the boundaries $\psi \pm \eta = \pm \frac{\pi}{2}$ are the null surfaces \mathcal{I}^+ and \mathcal{I}^-).

The de Sitter space-times are also conformal to a (finite) part of ds_E^2 and generally, all the closed Robertson-Walker metrics (Minkowski space, the Sitter space are included as special cases) are conformally equivalent to the Einstein static universe. The Robertson-Walker metric,

$$(4) \quad ds_{R-W}^2 = K^2(ct)[d\eta^2 + \sin^2 \eta(d\theta^2 + \sin^2 \theta d\varphi^2)] - c^2 dT^2$$

under the transformation

$$(5) \quad dT = \frac{1}{K(ct)} d\psi$$

gives (1) .

In this paper I will construct a solution of the Einstein dust equation (M, g) , which asymptotically approaches the Einstein static universe. This asymptotic limit requires the energy-momentum tensor to have the form of dust and $\Lambda \neq 0$ to be positive.

Here I shall adopt in the definition of this kind of asymptotically Einstein static universe space-times that the boundary ∂M is consisted by two disjoint (closed) spacelike ² surfaces \mathcal{I}^+ and \mathcal{I}^- (for the Robertson-Walker metrics, \mathcal{I}^+ and \mathcal{I}^- are T_b , respectively T_e , the beginning of world and the end of universe).

2 A stationary asymptotically Einstein static universe solution

Stationary gravitational fields are characterized by the existence of a timelike Killing vector field ξ . So, in a stationary space-time (M, g) we can have a global causal structure, i.e. we can introduce an adapted coordinate system $(x^a) = (x^\alpha, t)$, $\xi = \frac{\partial}{\partial t}$ so that the metric g_{ab} is independent of t ,

$$(6) \quad ds^2 = h_{\alpha\beta} dx^\alpha dx^\beta + F(dt + A_\alpha dx^\alpha)^2, \quad F \equiv \xi_a \xi^a < 0.$$

²In asymptotically flat spaces, \mathcal{I}^+ and \mathcal{I}^- are null surfaces each of which of $\mathbb{R}^1 \times S^2$ topology.

The timelike unitary vector field $h^0 \equiv (-F)^{-\frac{1}{2}}\xi$, which is globally defined on M , indicates not only the time-orientation in every point $p \in M$, but also gives a global time coordinate t on M .

Stationarity (i.e. time translation symmetry) means that there exists a 1-dimensional group G_1 of isometries ϕ_t whose orbits are timelike curves parametrized by t .

Using the 3-projection formalisme (Geroch (1971)) developed on the 3-dimensional differentiable factor manifold \mathcal{S}_3 , the Einstein's field equations

$$(7) \quad R_{ab} - \frac{1}{2}Rg_{ab} + \Lambda g_{ab} = \kappa T_{ab},$$

for stationary fields take the following simplified form:

$$(8) \quad \left\{ \begin{array}{l} R_{ab}^{(3)} = \frac{1}{2}F^{-2}(\frac{\partial F}{\partial x^a}\frac{\partial F}{\partial x^b} + \omega_a\omega_b) + \kappa(h_a^c h_b^d - F^{-2}\tilde{h}_{ab}\xi^a\xi^b)(T_{cd} - \frac{1}{2}Tg_{cd}); \\ F_{,a}^{\parallel a} = F^{-1}\tilde{h}_{ab}(\frac{\partial F}{\partial x^a}\frac{\partial F}{\partial x^b} - \omega_a\omega_b) - 2\kappa F^{-1}\xi^a\xi^b(T_{ab} - \frac{1}{2}Tg_{ab}); \\ \omega_a^{\parallel a} = 2F^{-1}\tilde{h}_{ab}\frac{\partial F}{\partial x^a}\omega_b \\ F\epsilon^{abc}\omega_{c,b} = 2\kappa h_b^a T_c^b \xi^c \end{array} \right.$$

Here, “ \parallel ” denotes the covariant derivative associated with the conformal metric tensor $\tilde{h}_{ab} = -Fh_{ab}$ on \mathcal{S}_3 ($h_{ab} = g_{ab} + h_a^0 h_b^0$ is the projection tensor) and $\omega^a = \frac{1}{2}\epsilon^{abcd}\xi_b\xi_{c;d} \neq 0$ ³ is the rotational vector ($\omega^a\xi_a = 0$, $\mathcal{L}_\xi\omega = 0$).

This Einstein static universe asymptotic limit suggests there are more than one symmetry on M . I suppose the metric g_{ab} admits also an Abelian group of isometries G_2 generated by the two spacelike Killing vector fields η and ζ , $\mathcal{L}_\eta g_{ab} = \mathcal{L}_\zeta g_{ab} = 0$, $\eta_a\eta^a > 0$, $\zeta_a\zeta^a > 0$ and the integral curves of η are closed (spatial) curves.

The circularity theorem (due to Kundt) states that an (axisymmetric) metric can be written in the (2+2)-split if and only if the conditions

$$(9) \quad (\eta^{[a}\xi^b\xi^{c;d]})_{;e} = 0 = (\xi^{[a}\eta^b\eta^{c;d]})_{;e}$$

are satisfied.

The existence of the orthogonal 2-surfaces is possible for dust solutions, provided that the 4-velocity of dust satisfies the condition:

$$(10) \quad u_{[a}\xi_b\eta_{c]} = 0, \quad u^a = (-H)^{-\frac{1}{2}}(\xi^a + \Omega\eta^a) = (-H)^{-\frac{1}{2}}l^i\xi_i^a, \quad \text{where} \\ l^i \equiv (1, \Omega), \quad H = \gamma_{ij}l^il^j, \quad \gamma_{ij} \equiv \xi_i^a\xi_{aj}, \quad i, j = 1, 2, \quad \xi_1 = \xi; \quad \xi_2 = \eta$$

³Here I use the convention: round brackets denote symmetrization and square brackets antisymmetrization and Ω is the angular velocity.

) i.e., the trajectories of the dust lie on the transitivity surfaces of the group generated by the Killing vectors ξ , η .

Using an adapted coordinate system, the metric (6) can be written in the following form:

$$(11) \quad ds^2 = e^{-2U} [e^{2V} (dr^2 + dz^2) + W^2 d\varphi^2] - e^{2U} (dt + Ad\varphi)^2$$

where the functions ⁴ U , V , W and A depend only on the coordinates (r, z) ; these coordinates are also conformal flat coordinates on the 2-surface S_2 orthogonal to 2-surface T_2 of the commuting Killing vectors $\xi = \partial_t$ and $\eta = \partial_\varphi$.

If we identify the 4-velocity of the dust u^a with timelike Killing vector $\xi^a = \partial_t = (0, 0, 0, 1)$ then (11) represents a co-moving system ($x^1 = r$, $x^2 = z$, $x^3 = \varphi$, $x^0 = t$) with dust, $u_a = \xi_a = (0, 0, -e^{2U}A, -e^{2U})$ and

$$(12) \quad \begin{cases} g_{11} = g_{22} = e^{-2U+2V} = h_{11} = h_{22}, \\ g_{33} = e^{-2U}W^2 - e^{2V}A^2 = h_{33}, \quad g_{00} = \xi_0 = -e^{2U} = F, \\ g_{03} = \xi_3 = -e^{2U}A, \quad g_{13} = g_{23} = g_{10} = g_{20} = 0 \end{cases}$$

We can use the complex coordinates (q, \bar{q}) on the 2-surface S_2 :

$$(13) \quad q = \frac{1}{\sqrt{2}}(r + iz)$$

The surface element on T_2 is $f_{ab} = 2\xi_{[a}\eta_{b]}$, $f_{ab}f^{ab} < 0$ and the surface element on S_2 is \tilde{f}_{ab} , the dual tensor of f_{ab} , $\tilde{f}_{ab} = \frac{1}{2}\epsilon_{abcd}f^{cd}$

Thus the Einstein's dust equations with constant $\Lambda > 0$ (8) for the metric (12) will take the following form:

$$(14) \quad \left\{ \begin{aligned} \frac{\partial^2 W}{\partial q \partial \bar{q}} &= -\Lambda W e^{2V-2U} \\ \frac{\partial^2 U}{\partial q \partial \bar{q}} + \frac{1}{2W} \left(\frac{\partial U}{\partial q} \frac{\partial W}{\partial \bar{q}} + \frac{\partial U}{\partial \bar{q}} \frac{\partial W}{\partial q} \right) + \frac{1}{2W^2} e^{4U} \frac{\partial A}{\partial q} \frac{\partial A}{\partial \bar{q}} &= (\mu - 2\Lambda) \frac{e^{2V-2U}}{4} \\ \frac{\partial^2 A}{\partial q \partial \bar{q}} - \frac{1}{2W} \left(\frac{\partial A}{\partial q} \frac{\partial W}{\partial \bar{q}} + \frac{\partial A}{\partial \bar{q}} \frac{\partial W}{\partial q} \right) + 2 \left(\frac{\partial A}{\partial q} \frac{\partial U}{\partial \bar{q}} + \frac{\partial A}{\partial \bar{q}} \frac{\partial U}{\partial q} \right) &= 0 \\ \frac{\partial^2 W}{\partial q \partial \bar{q}} - 2 \frac{\partial W}{\partial q} \frac{\partial V}{\partial \bar{q}} + 2W \left(\frac{\partial U}{\partial q} \right)^2 - \frac{1}{2W} e^{4U} \left(\frac{\partial A}{\partial q} \right)^2 &= 0 \\ \frac{\partial^2 V}{\partial q \partial \bar{q}} + \frac{\partial U}{\partial q} \frac{\partial U}{\partial \bar{q}} + \frac{1}{(2W)^2} e^{4U} \frac{\partial A}{\partial q} \frac{\partial A}{\partial \bar{q}} &= -\Lambda \frac{e^{2V-2U}}{2} \end{aligned} \right.$$

⁴The function W is defined invariantly as $W^2 \equiv -2\xi_{[a}\eta_{b]}\xi^a\eta^b$.

Here $\Delta = \frac{\partial^2}{\partial r^2} + \frac{\partial^2}{\partial z^2} = 2\frac{\partial^2}{\partial q\partial\bar{q}}$ is the Laplace operator and the energy-momentum tensor T_{ab} has the form as:

$$(15) \quad \kappa T_{ab} = -\Lambda g_{ab} + \mu u_a u_b, \quad \mu > 0, \Lambda = \text{const.} > 0.$$

The conservation law, $T_{;b}^{ab} = 0$ implies $U_{;a} = 0$ and is a consequence of the field equations, being used in place of one of the equations (14).

Using the simplification $U = 0$ in (12) then the matter current is geodesic ($\dot{u}_a = u_{a;b}u^b = 0$) and without expansion ($\theta = u^a_{;a} = 0$), but has a non-rigidly rotation, ($\omega = \sqrt{\frac{1}{2}\omega_{ab}\omega^{ab}} \neq 0$) and $\sigma \neq 0$; the field equations (14) will take the following simplified form:

$$(16) \quad \left\{ \begin{array}{l} \frac{\partial^2 W}{\partial q \partial \bar{q}} = -\Lambda W e^{2V} \\ \frac{1}{2W^2} \frac{\partial A}{\partial q} \frac{\partial A}{\partial \bar{q}} = (\mu - 2\Lambda) \frac{e^{2V}}{4} \\ \frac{\partial^2 A}{\partial q \partial \bar{q}} - \frac{1}{2W} \left(\frac{\partial A}{\partial q} \frac{\partial W}{\partial \bar{q}} + \frac{\partial A}{\partial \bar{q}} \frac{\partial W}{\partial q} \right) = 0 \\ \frac{\partial^2 W}{\partial q \partial \bar{q}} - 2 \frac{\partial W}{\partial q} \frac{\partial V}{\partial \bar{q}} - \frac{1}{2W} \left(\frac{\partial A}{\partial q} \right)^2 = 0 \\ \frac{\partial^2 V}{\partial q \partial \bar{q}} + \frac{1}{(2W)^2} \frac{\partial A}{\partial q} \frac{\partial A}{\partial \bar{q}} = -\Lambda \frac{e^{2V}}{2} \end{array} \right.$$

Finally, if we take into account the third symmetry ($\zeta = \partial_z$ the spacelike Killing vector field) as a special case of the stationary axisymmetric (with ξ and η) solution (16) and if we match the parameters by a particular choice such that the exterior field is conformal with Einstein static universe⁵, then we obtain a stationary asymptotically Einstein static universe solution:

$$ds^2 = e^{2V(r)}(dr^2 + dz^2) + W^2(r)d\varphi - (dt + A(r)d\varphi)^2$$

where $V(r)$, $W(r)$, $A(r)$ and $\mu(r)$ depend only on r .

⁵For the case when $\Lambda = 0$ the extern field becomes static ($\omega^a = \frac{1}{2}\epsilon^{abcd}\xi_b\xi_{c;d} = 0$) even the dust was in rotation with $\Omega = \text{const.}$

Indeed denoting $\frac{\partial}{\partial r} = '$ the field equations (16) take the form:

$$(17) \quad \begin{cases} W'' = -2\Lambda W e^{2V} \\ A'^2 = (\mu - 2\Lambda) W^2 e^{2V} \\ \frac{A''}{A'} = \frac{W'}{W} \\ 2W'' - 4W'V' - \frac{1}{W} A'^2 = 0 \\ V'' + \frac{1}{4W^2} A'^2 = -\Lambda e^{2V} \end{cases}$$

The third equation of the previous system is integrable: $A' = \alpha W$, where $\alpha = \text{const.} \neq 0$. Taking $\alpha = 2$ and inserting in the rest of the system we get:

$$(18) \quad \begin{cases} \mu = 2\Lambda + 4e^{-2V} \\ \frac{W'}{W} = -\frac{1 + \Lambda e^{2V}}{V'} \\ V'' + \Lambda e^{2V} + 1 = 0 \end{cases}$$

Because the cosmological constant Λ is very small (less than 10^{-57}cm^{-2}) we can approximate $\Lambda e^{2V} \approx \Lambda(1 + 2V)$. In this approximation the system gets the form:

$$(19) \quad \begin{cases} \mu = 2\Lambda + \frac{4}{1 + 2V} \\ \frac{W'}{W} = -\frac{1 + \Lambda(1 + 2V)}{V'} \\ V'' + \Lambda(1 + 2V) + 1 = 0 \end{cases}$$

Integrating first the last linear equation ($V'' + 2\Lambda V + \Lambda + 1 = 0$) we obtain:

$$(20) \quad \begin{aligned} V(r) &= C_1 \sin(\sqrt{2\Lambda}r + \alpha) - \frac{1 + \Lambda}{2\Lambda} \\ \mu(r) &= 2\Lambda + 4 \frac{1}{1 + 2C_1 \sin(\sqrt{2\Lambda}r + \alpha) - \frac{1 + \Lambda}{\Lambda}} \\ W(r) &= \frac{1}{2} C_3 C_1 \sqrt{2\Lambda} \cos(\sqrt{2\Lambda}r + \alpha) \\ A(r) &= C_3 \left(C_1 \sin(\sqrt{2\Lambda}r + \alpha) - \frac{1 + \Lambda}{2\Lambda} \right) + C_4 \end{aligned}$$

where α, C_1, C_2, C_3 are parameters of integration of the system which must be chosen so that for $r \rightarrow \infty$ we have $\mu(r) \rightarrow 2\Lambda$ and $A(r) \rightarrow 0$, so the solution found is asymptotically Einstein static universe solution...

3 Some discussion

Another type of event horizon, caused by the presence of the repulsive Λ term, is defined as the boundary of the points of spacetime from which light can never reach the observer. This boundary is called *the cosmological event horizon* of the observer (see [1]).

On such a space-time ⁶ which admits predicted black hole event horizons and it is conformal with Einstein static universe at infinity, there are points from where light rays (null geodesics) that escape to null infinity \mathcal{I}^+ would return to \mathcal{I}^- on a proper finite time interval (as in general closed universes) and would also return to their source. Such a region of points $p \in M, p \in J^+(\mathcal{I}^-) \cap J^-(\mathcal{I}^+)$ contains causal closed curves (null or temporal piecewise geodesics). Indeed, this is the case since the behaviour near infinity could be determined by the light ray structure of space-time and that light rays remain light rays (and causal structure has not changed either) of the Einstein static universe under a conformal transformation ($ds^2 = 0$ implies $ds_E^2 = \Theta^2 ds^2 = 0$) and light rays of the Einstein universe have the property that they all return to their source at a time interval to cover $\Delta T = \frac{2\pi}{c}$.

References

- [1] **Gibbons, G.W., Hawking S.W.**, *Cosmological event horizons, thermodynamics, and particle creation*, Physical Review D, (1976)
- [2] **Hawking, S.W.**, *Black Holes*, eds. Dewitt & Dewitt (Gordon and Breach), (1973)
- [3] **Huesler, M.**, *Stationary Black Holes: uniqueness and beyond*, Living Reviews, (1998)
- [4] **Iftime, M.D.**, *Spacetime distribution*, Geometry & Topology, (1998)
- [5] **Kramer, D., Stephani, H., Herlt, E., MacCallum, M., Schmutzer, E.**, *Exact solutions of Einstein's field equations*, Cambridge Univ. Press, (1980)

⁶No strong causality condition but only a stable causal condition is holded on space-time.