

An Expansion Term In Hamilton's Equations.

Mark D. Roberts,

Department of Mathematics and Applied Mathematics,
University of Cape Town, South Africa

roberts@gmunu.mth.uct.ac.za

November 20, 2018

Published: Europhysics Letters **45**(1991)26-31

<http://www.edpsciences.com/docinfos/EURO/OnlineEURO.html>

Epreprint: <http://xxx.lanl.gov/abs/gr-qc/9810090>

Comments: Final Version, no diagrams, LaTeX2e, small changes from previous eprint and from the published version.

1999 PACS Physics and Astronomy Classification Scheme:

<http://publish.aps.org/eprint/gateway/pacslist>

11.10.Ef, 04.60+Ds, 52.60+h, 03.70+k

1991 Mathematical Reviews Classification Scheme:

<http://www.ams.org/msc>

81S99, 83C45, 83C43

Keywords:

Hamilton's Equations, Preferred Vector Field,

Expansion, Symplectic Geometry.

Abstract

For any given spacetime the choice of time coordinate is undetermined. A particular choice is the absolute time associated with a preferred vector field. Using the absolute time Hamilton's equations are

$$\begin{aligned} -\frac{\delta H_c}{\delta q} &= \dot{\pi} + \Theta\pi, \\ +\frac{\delta H_c}{\delta \pi} &= \dot{q}, \end{aligned}$$

where $\Theta = V_{;a}^a$ is the expansion of the vector field. Thus there is a hitherto unnoticed term in the expansion of the preferred vector field. Hamilton's equations can be used to describe fluid motion. In this case the absolute time is the time associated with the fluid's co-moving vector. As measured by this absolute time the expansion term is present. Similarly in cosmology, each observer has a co-moving vector and Hamilton's equations again have an expansion term. It is necessary to include the expansion term to quantize systems such as the above by the canonical method of replacing Dirac brackets by commutators. Hamilton's equations in this form do not have a corresponding symplectic form. Replacing the expansion by a particle number $N \equiv \exp(-\int \Theta d\tau)$ and introducing the particle numbers conjugate momentum π^N the standard symplectic form can be recovered with two extra fields N and π^N . Briefly the possibility of a non-standard symplectic form and the further possibility of there being a non-zero Finsler curvature corresponding to this are looked at.

1 Introduction

One of the problems with the standard treatment of Hamilton's equations is that it assumes that a preferred time can be "taken out". This is not a restriction in special relativity or in static general relativity; however in a non-static spacetime one wants to define time with respect to a given preferred vector field. Below it is shown that defining time by this specific preferred vector field, the expansion of the vector field has to be included in Hamilton's equations. The expansion term also occurs in the Poisson bracket and the equation for ordinary time evolution of an arbitrary variable. The type of systems where the expansion term is important are perfect fluids, where the preferred vector field is that of a co-moving element of the fluid; and in cosmology where the preferred vector field is that of an observer co-moving with the expansion. If one wants to quantize such systems one has to replace the Poisson (or for constrained systems the Dirac) bracket by a quantum commutator. After this is done the term in the expansion remains. For a perfect fluid quantization has been done with expansion present Roberts (1998) [10]. Quantization of cosmology, Tipler (1986) [12] is usually approached via the ADM (Arnowitt, Deser, and Misner (1962)) formalism [3]. In this time is taken out so that the normal to the surface has no expansion, vorticity, or acceleration. It is possible that this can be generalized using the parametric manifold approach of Boersma and Dray (1995) [4]. The inclusion of a normal vector with expansion would require use of Hamilton's equations with the expansion term presented here. Expansion is the main qualitative property in cosmology so that the addition of this term is of importance. The additional expansion term was first found for the Hamiltonian description of particular fluids and then generalized. The expansion term might be directly derivable by different methods using Peierls (1952) [8] brackets. The manner in which time is usually taken out is simply to label one coordinate as t . If there is a preferred vector field V^a present it is associated with a preferred time via the absolute derivative

$$\dot{X}_{abc\dots} = \frac{D}{d\tau} X_{abc\dots} = V^e \nabla_e X_{abc\dots} = V^e X_{abc\dots;e}. \quad (1)$$

The usual approach is recovered by letting $V^a = (1, 0, 0, 0)$. In general V^a has a more complex structure than this and can be decomposed into the shear, vorticity, etc. . . , see for example Hawking and Ellis (1973) [7] p.83. Additions to Hamilton's equations have also been used to overcome conceptual problems in quantum mechanics Ghirardi, Rimini, and Weber

(1986) [5].

2 Hamilton's equations

The canonical Hamiltonian H_c for fields is

$$H_c = \int \sqrt{-g} d^4x H_d = \int \sqrt{-g} d^4x (\pi^i \dot{q}_i - L), \quad (2)$$

where H_d is the Hamiltonian density. The corresponding canonical phase space Lagrange action is

$$I_{ph} = \int \sqrt{-g} d^4x (\pi^i \dot{q}_i - H_d). \quad (3)$$

Varying I_{ph}

$$\begin{aligned} \delta I_{ph} &= \int \sqrt{-g} d^4x \left[\pi^i \delta \left(\frac{d\dot{q}_i}{d\tau} \right) + \frac{d\dot{q}_i}{d\tau} \delta \pi^i - \left(\frac{\partial H_d}{\partial q^i} \right) \delta q_i \right. \\ &\quad \left. - \left(\frac{\partial H_d}{\partial q_a^i} \right) \delta q_a^i - \left(\frac{\partial H_d}{\partial \pi^i} \right) \delta \pi^i - \left(\frac{\partial H_d}{\partial \pi_a^i} \right) \delta \pi_a^i \right], \end{aligned} \quad (4)$$

replacing $\frac{D}{d\tau}$ by $V^a \nabla_a$, as in equation 1 taking $\delta V^a = 0$, interchanging infinitesimals thus

$$\begin{aligned} \pi_i \left(\frac{\delta d\dot{q}^i}{d\tau} \right) &= \pi_i \delta (V^a q_{;a}^i) = \pi_i V^a \delta (q_{;a}^i) \\ &= \pi V^a \nabla_a (\delta q^i) = \nabla_a (V^a \pi_i \delta q^i) - (\delta q^i) \nabla_a (\pi_i V^a), \end{aligned} \quad (5)$$

and disarding the surface term $\nabla_a (V^a \pi_i \delta q^i)$ leaves

$$\begin{aligned} \delta I_{ph} &= \int \sqrt{-g} d^4x \left[\delta q^i \left(-\nabla_a (\pi_i V^a) - \frac{\partial H_d}{\partial q^i} + \nabla_a \frac{\partial H_d}{\partial q_{;a}^i} \right) \right. \\ &\quad \left. + \delta \pi^i \left(V^a \nabla_a q_i - \frac{\partial H_d}{\partial \pi^i} + \nabla_a \frac{\partial H_d}{\partial \pi_{;a}^i} \right) \right]. \end{aligned} \quad (6)$$

For the δq^i term to vanish

$$-\frac{\delta H_c}{\delta q^i} = -\frac{\partial H_d}{\partial q^i} + \nabla \left(\frac{\partial H_d}{\partial q_{;a}^i} \right) = \nabla_a (\pi^i V^a) = \dot{\pi}^i + \Theta \pi^i \quad (7)$$

For the $\delta\pi^i$ term to vanish

$$+ \frac{\delta H_c}{\delta\pi^i} = + \frac{\partial H_d}{\partial\pi^i} - \nabla \left(\frac{\partial H_d}{\partial\pi^i_{;a}} \right) = V^a \nabla_a q^i = \dot{q}^i \quad (8)$$

Equations 7 and 8 generalize Hamilton's equations. The equations are unaffected if π^i and q^i have spacetime indices $\pi^i \rightarrow \pi^i_{ab\dots}$, $q^i \rightarrow q^i_{ab\dots}$ and so on. The standard Poisson bracket is defined as

$$\{X, Y\} \equiv \frac{\delta X}{\delta q_i} \frac{\delta Y}{\delta\pi^i} - \frac{\delta X}{\delta\pi_i} \frac{\delta Y}{\delta q^i}. \quad (9)$$

Equations 7 and 8 suggest alternative Poisson bracket to the standard bracket,

$$\begin{aligned} \{A, B\}_A &= \left(\Theta\pi_i + \frac{\delta A}{\delta q_i} \right) \frac{\delta B}{\delta\pi^i} - \frac{\delta A}{\delta\pi^i} \left(\Theta\pi_i + \frac{\delta B}{\delta q_i} \right) \\ &= \{A, B\} + \Theta\pi_i \left(\frac{\delta B}{\delta\pi^i} - \frac{\delta A}{\delta\pi^i} \right), \end{aligned} \quad (10)$$

containing the expansion Θ . The alternative bracket does not give $\{\pi_i, q^i\} = 1$, also for particular constrained systems the correct Lagrange multipliers are not recovered. Thus the standard bracket is preferred. Perhaps Peierls' brackets [8] can show that the alternative bracket was not suitable by a general method. The time evolution of an arbitrary variable $X(\tau, q^i, \pi^i)$ is given by

$$\frac{dX}{d\tau} = \frac{\partial X}{\partial\tau} + \dot{q}^i \frac{\delta X}{\delta q^i} + \dot{\pi}^i \frac{\delta X}{\delta\pi^i}, \quad (11)$$

substituting q^i and π^i using equations 7 and 8 gives the time evolution

$$\begin{aligned} \frac{dX}{d\tau} &= \frac{\partial X}{\partial\tau} + \{X, H_c\} - \Theta\pi^i \frac{\delta X}{\delta\pi^i} \\ &= \frac{\partial X}{\partial\tau} + \{X, H_c\}_A - \Theta\pi^i \frac{\delta H_c}{\delta\pi^i}. \end{aligned} \quad (12)$$

Note in particular that the time evolution of the canonical Hamiltonian H_c is no longer solely given by its partial derivative with respect to time and is

$$\frac{dH_c}{d\tau} = \frac{\partial H_c}{\partial\tau} - \Theta\pi^i \frac{\delta H_c}{\delta\pi^i}. \quad (13)$$

3 Symplectic Structure

Following the standard, see for example Wald (1994), [13] p.11 procedure for producing the symplectic form let

$$y^\mu = (q^1, \dots, q^n; \pi^1, \dots, \pi^n) \quad (14)$$

to produce the equations

$$\frac{dy^\mu}{d\tau} = \sum_{\nu=1}^{2n} \Omega^{\mu\nu} \frac{\delta H_c}{\delta y^\nu}, \quad (15)$$

when $\nu = \mu + n$

$$\begin{aligned} \Omega^{\mu\nu} &= 1, \\ \Omega^{\nu\mu} &= -\frac{\dot{\pi}^\mu}{\dot{\pi}^\mu + \Theta \pi^\mu}, \end{aligned} \quad (16)$$

otherwise $\Omega = 0$. This is not a symplectic matrix as (unless $\Theta = 0$) it is not antisymmetric. Another problem is that the expansion Θ is not explicitly defined in term of the fields q^i and the momenta π^i . Also recall that for symplectic spaces the Hamiltonian is usually conserved, Arnold 1978 [2] p.207, but from equation 13, $\frac{dH_c}{d\tau}$ is not necessarily zero here.

In order to produce a symplectic structure first remove the expansion Θ so that there are only fields and momenta present. By analogy with fluids, Roberts(1997) [9], let $\Theta = -\frac{\dot{N}}{N}$ where N is a particle number. The existing 2i equations are

$$\begin{aligned} -\frac{\delta H_c}{\delta q^i} &= \dot{\pi}^i + \frac{\dot{N}}{N} \pi^i = N \dot{\pi}'^i, & \dot{\pi}'^i &\equiv \frac{\pi^i}{N}, \\ +\frac{\delta H_c}{\delta \pi^i} &= \dot{q}^i. \end{aligned} \quad (17)$$

N is a new field so it also has to be added to the Hamiltonian density to give the new Hamiltonian density

$$H_a = H_d + \pi_N \dot{N} = \pi^{i+1} \dot{q}_{i+1} - L, \quad (18)$$

and new canonical Hamiltonian H_A . In addition to the 2i equations described by 17 there are two new equations

$$\begin{aligned} -\frac{\delta H_A}{\delta N} &= \dot{\pi}^N - \frac{\dot{N}}{N} \pi^N = N \dot{\pi}'^N, & \pi'^N &\equiv \frac{\pi^N}{N} \\ +\frac{\delta H_A}{\delta \pi^N} &= \dot{N}, \end{aligned} \quad (19)$$

From equations 18 and 19 we can now define

$$q'^i \equiv \int d\tau \left(\frac{\dot{q}^i}{N} \right), \quad (20)$$

and

$$z^\mu \equiv (q'^1, \dots, q'^n, N; \pi'^1, \dots, \pi'^n, \pi'^N), \quad (21)$$

to produce equations of the same form as 15 with z^μ replacing y^μ and

$$\Omega^{\mu\nu} = N, \quad \Omega^{\nu\mu} = -N, \quad (22)$$

otherwise $\Omega = 0$. This symplectic matrix describes the system and is two dimensions higher than the original dimension, also it is N times the standard form. To produce a symplectic matrix of the standard form rather than N times it, leave q^i in its original form and define

$$\pi'^i \equiv \int d\tau \left(\dot{\pi}^i + \frac{\dot{N}}{N} \pi^i \right). \quad (23)$$

A possibility is that the extra structure present because of the expansion can lead to a non standard form of the symplectic matrix and that this in turn would give non-vanishing Finsler curvature, Rund (1959) [11], of some geometric object. In order to attempt investigation of this **first** try adding a term $\gamma(q, \dot{q}, \pi, \dot{\pi})$ to the Hamiltonian density H_a to give a new Hamiltonian density

$$H_\gamma = \pi^{i+1} \dot{q}_{i+1} - L + \gamma(q, \dot{q}, \pi, \dot{\pi}), \quad (24)$$

and a new canonical Hamiltonian H_Γ . Hamilton's equations become

$$\begin{aligned} -\frac{\delta H_\Gamma}{\delta q^i} &= \dot{\pi}^i - \frac{\dot{N}}{N} \pi^i + \frac{\partial \gamma}{\partial q^i} - \left(\frac{\partial \gamma}{\partial \dot{q}} \right)' + \frac{\dot{N}}{N} \frac{\partial \gamma}{\partial \dot{q}^i}, \\ +\frac{\delta H_\Gamma}{\delta \pi^i} &= \dot{q}^i + \frac{\partial \gamma}{\partial \pi^i} - \left(\frac{\partial \gamma}{\partial \dot{\pi}^i} \right)' + \frac{\dot{N}}{N} \frac{\partial \gamma}{\partial \dot{\pi}^i}. \end{aligned} \quad (25)$$

Taking an ansatz for γ of the form

$$\begin{aligned} \gamma &= k_1 + k_2 q + k_3 \pi + k_4 q \pi \\ &+ k_5 \dot{q} + k_6 q \dot{q} + k_7 \pi \dot{q} + k_8 q \pi \dot{q} \\ &+ k_9 \dot{\pi} + k_{10} q \dot{\pi} + k_{11} \pi \dot{\pi} + k_{12} q \pi \dot{\pi}, \end{aligned} \quad (26)$$

it is immediately apparent that $k_1 = k_2 = k_3 = k_4 = k_5 = k_7 = k_9 = 0$ and can be shown that nothing of this form will produce an antisymmetric symplectic matrix. **Secondly** recall, Hargreaves (1908) [6] that in fluid mechanics variation of the Lagrangian and Hamiltonian is achieved via the infinitesimal first law of thermodynamics, typically

$$\delta L = \delta p = N\delta h - NT\delta s + \sum X_\mu \delta x^\mu, \quad (27)$$

where p is the pressure, N the particle number, h the enthalpy, T the temperature, and s the entropy. X_μ are any additional intensive variables, x^μ are any additional extensive variables. For definitions of intensive and extensive variables see Adkins (1983) [1]; briefly if an intensive variable is doubled the mass remains the same, but if an extensive variable is doubled the mass doubles. In particular term $\mu\delta N$ can be added, where μ is the chemical potential. By analogy with the first conservation law 27 choose new Hamiltonian density H_e which can be expressed infinitesimally in the form

$$\delta H_e = \delta H_n - \alpha \times (\delta\beta). \quad (28)$$

It immediately follows that the choice $\alpha = \frac{\dot{\pi}^N}{\pi^N} q^i, \beta = \pi_i$ gives the equations

$$\begin{aligned} -\frac{\delta H_E}{\delta q^i} &= \dot{\pi}^i - \frac{\dot{N}}{N} \pi^i, \\ +\frac{\delta H_E}{\delta \pi^i} &= \dot{q}^i - \frac{\dot{\pi}^N}{\pi^N} q^i, \end{aligned} \quad (29)$$

where i ranges from 1 to $n+1$ so as to include the particle number N and its associated momentum π^N . Using equations 14 and 15 with n replaced with $n+1$ to try and remove the time derivatives and produce a symplectic matrix one instead arrives at

$$\begin{aligned} \frac{dq^i}{d\tau} &= \frac{\delta H_E}{\delta \pi_i} + x q^i \left(-\frac{1}{\pi^N} \frac{\delta H_E}{\delta N} + \frac{1}{N} \frac{\delta H_E}{\delta \pi^N} \right), \\ \frac{d\pi^i}{d\tau} &= -\frac{\delta H_E}{\delta q_i} - y \pi^i \left(-\frac{1}{N} \frac{\delta H_E}{\delta \pi^N} + \frac{1}{\pi^N} \frac{\delta H_E}{\delta N} \right) \end{aligned} \quad (30)$$

where x, y are the number of iterations used to remove \dot{N} and $\dot{\pi}^N$. Taking x and y to be finite,

$$y^\mu = (q^i, \pi^i, N, \pi^N), \quad (31)$$

and equation 15 gives in place of $\Omega^{\mu\nu}$ the matrix

$$M^{\mu\nu} = \begin{pmatrix} 0 & 1 & -\frac{xq^i}{\pi^N} & \frac{xq^i}{N} \\ -1 & 0 & -\frac{y\pi^i}{\pi^N} & \frac{y\pi^i}{N} \\ 0 & 0 & -x\frac{N}{\pi^N} & 1+x \\ 0 & 0 & -1-x & y\frac{\pi^N}{N} \end{pmatrix}, \quad (32)$$

which has trace $y\frac{\pi^N}{N} - x\frac{N}{\pi^N}$ and determinant $1+x+y$, however it is not anti-symmetric and so is not symplectic.

4 Conclusion

It was shown that when “time” used is that of a preferred vector field that Hamilton’s equations gain a term in the expansion of the vector field. The implications of this for symplectic structure were discussed.

5 Acknowledgements

This work has been supported in part by the South African Foundation for Research and Development (FRD).

References

- [1] Adkins,C.J.(1983) Equilibrium Thermodynamics, Third Edition, Cambridge University Press.
- [2] Arnold,V.I.(1978) Mathematical Methods of Classical Mechanics. Translated by K.Vogtmann and A.Weinstein, Math.Rev.57#5460 Springer-Verlag, Berlin.
- [3] Arnowitt,R.,Deser,S. and Misner,C.W.(1962) The Dynamics of General Relativity, in An Introduction to Current Research, Edited L.Witten, Math.Rev. 26#1182
- [4] Boersma,S. and Dray,T.(1995) Parametric Manifolds. Math.Rev. 95m:53093 *J.Math.Phys.* **36**,1378.
- [5] Ghirardi,G.C., Rimini,A., and Weber,T.(1986) Unified dynamics fo microscopic and macroscopic systems. Math.Rev.87e:81014 *Phys.Rev.D*,**34**,470.

- [6] Hargreaves,R.(1908) A Pressure-integral as Kinetic Potential. *Phil.Mag.***16**,436.
- [7] Hawking,S.W. and Ellis,G.F.R.(1973) The Large Scale Structure of Space-Time. Cambridge University Press.
- [8] Peierls,R.E.(1952) The Commutation Laws of Relativistic Field Theory. *Math.Rev.*14#520. *Proc.Roy.Soc.***A214**,143.
- [9] Roberts,M.D.(1997) Fluid Symmetry Breaking II: Velocity Potential Method. *Hadronic J.***20**,73.
- [10] Roberts,M.D. The Quantum Commutator Algebra of a Perfect Fluid. <http://xxx.lanl.gov/abs/gr-qc/9810089>
- [11] Rund,H.(1959) The Differential Geometry of Finsler Spaces. *Math.Rev.*21#4462 Springer-Verlag, Berlin.
- [12] Tipler,F.(1986) Interpreting the Wave Function of the Universe. *Math.Rev.*87g:83161 *Phys.Rep.***C137**,231.
- [13] Wald,R.M.(1994) Quantum Field Theory in Curved Spacetimes and Black Hole Thermodynamics. *Math.Rev.*95i:81176 University of Chicago Press.