

Geodesics on non-Riemannian Geometric theory of Planar Defects

L.C. Garcia de Andrade¹

Abstract

The method of Hamilton-Jacobi is used to obtain geodesics around non- Riemannian planar torsional defects. It is shown that by perturbation expansion in the Cartan torsion the geodesics obtained are parabolic curves along the plane x-z when the wall is located at the plane x-y. In the absence of defects the geodesics reduce to straight lines. The family of parabolas depend on the torsion parameter and describe a gravitationally repulsive domain wall. Torsion here plays the role of the Burgers vector in solid state physics.

¹Departamento de Física Teórica - Instituto de Física - UERJ
Rua São Fco. Xavier 524, Rio de Janeiro, RJ
Maracanã, CEP:20550-003 , Brasil.
E-Mail.: GARCIA@SYMBCOMP.UERJ.BR

The investigation of topological defects [1] in cosmology follows closed analogous models in solid state physics [2]. Riemannian [3, 4, 5] and non-Riemannian [6, 7] models have been used to investigate these distributional defects such as disclinations [8] and dislocations [9] and domain walls analogous to ferromagnetic domains in crystallography. Lower dimensional models like disclinations in liquid crystals presented helical torsion structures as demonstrated by E. Dubois-Violette [10] and James Sethna [11]. More recently F. Moraes [8] has been used the same method of Hamilton-Jacobi (H-J) equations to investigate geodesics around torsional defects like electrons around dislocated metals. In this Letter I follow this same pattern to obtain geodesics around domain walls in 3-Dimensional gravity. Although isolated domain walls are nowadays of no interest in physics they still may play a role in mixed system in the condensed matter systems which serve as a laboratory for Cosmology. In this letter we find a simple example of a two dimensional gravity model from a domain wall solution of Einstein-Cartan gravity obtaining by reducing the dimensions of the spacetime metric

$$ds^2 = e^{8\pi G\sigma z} dt^2 - e^{-J^1 z} (dx^2 + dy^2) - dz^2 \quad (1)$$

where J^1 is the constant torsion at the planar wall, σ is the planar wall distribution of the wall energy density and G is the Newtonian gravitational constant. Due to translational symmetry of defects we shall consider the following metric

$$ds^2 = -e^{-J^1 z} dx^2 - dz^2 \quad (2)$$

at the $y = \text{constant}$ section. To obtain the geodesics I shall solve the H-J equation

$$\frac{\partial W}{\partial t} + H\left(\frac{\partial W}{\partial x^i}\right) = 0 \quad (3)$$

where

$$W[\gamma] = \frac{1}{2} \int_{t_0}^{t_1} g_{ij} \dot{x}^i \dot{x}^j dt \quad (4)$$

where γ is a parametrized curve where

$$s[\gamma] = \int_{t_0}^{t_1} (g_{ij} \dot{x}^i \dot{x}^j)^{\frac{1}{2}} dt \quad (5)$$

Here $H = L = \frac{1}{2} g_{ij} \dot{x}^i \dot{x}^j$ is the Hamiltonian and L is the Lagrangian. We must stress that although in general Riemann-Cartan spaces test particles

follow autoparallels and not geodesics ,off the domain wall there is vacuum and there GR remains valid. Nevertheless geodesics suffer indirectly the influence of torsion since they are also present in the metric considered. In the case of linearized EC gravity we have the following metric

$$ds^2 = -(1 - J^1 z)dx^2 - dz^2 \quad (6)$$

The H-J equation applied to metric (6) reduces to

$$\frac{\partial W}{\partial t} + (1 + J^1 z)\left(\frac{\partial W}{\partial x}\right)^2 + \left(\frac{\partial W}{\partial z}\right)^2 = 0 \quad (7)$$

where we have considered H in terms of the momenta $p_i = \frac{\partial W}{\partial x^i} = \frac{\partial L}{\partial \dot{x}^i}$. Follows from (7) that

$$W = c_1 t + W'(x, z) \quad (8)$$

and this in turn implies

$$c_1 + \left(\frac{\partial W'}{\partial x}\right)^2 + \left(\frac{\partial W'}{\partial z}\right)^2 = 0 \quad (9)$$

Writing the perturbation series for W' as

$$W' = W'_0 + J^1 W'_1 + (J^1)^2 W'_2 + \dots \quad (10)$$

and substituting (10) into (9) one obtains

$$\left(\frac{\partial W'_0}{\partial x}\right)^2 + \left(\frac{\partial W'_0}{\partial z}\right)^2 = -c_1 \quad (11)$$

and

$$-\frac{\partial W'_0}{\partial x} z + 2\frac{\partial W'_1}{\partial z}\frac{\partial W'_0}{\partial z} + 2\frac{\partial W'_1}{\partial x}\frac{\partial W'_0}{\partial x} = 0 \quad (12)$$

Equation (11) is solved at once by the ansatz

$$W'_0 = c_2 x + c_3 z \quad (13)$$

where the constants satisfy

$$c_1 = -\frac{1}{2}(c_2^2 + c_3^2) \quad (14)$$

Now the equation (12) can be written as

$$c_2 \frac{\partial W'_1}{\partial x} + c_3 \frac{\partial W'_1}{\partial y} = -\frac{1}{2} c_2^2 z \quad (15)$$

Equation (15) is solved by the method of characteristics where we first differentiate W'_1 with respect to the auxiliary variable u such as

$$\frac{dW'_1}{du} = \frac{\partial x}{\partial u} \frac{\partial W'_1}{\partial x} + \frac{\partial y}{\partial u} \frac{\partial W'_1}{\partial y} \quad (16)$$

Comparison of (15) with (16) yields

$$\frac{dW'_1}{du} = \frac{1}{2} c_2^2 z \quad (17)$$

and

$$\frac{\partial x}{\partial u} = c_2 \quad (18)$$

and

$$\frac{\partial y}{\partial u} = c_3 \quad (19)$$

Thus

$$x = c_2 u + t_1 \quad (20)$$

and

$$y = c_3 u \quad (21)$$

where t_1 is another parameter. Now by integration of the equation (17) we obtain

$$W'_1 = -\frac{1}{2} c_2^2 z u^2 + d \quad (22)$$

Where d is an integration constant. Thus by substituting (22) into (10) one has

$$W = -\frac{1}{2} (c_2^2 + c_3^2) t + c_2 x + c_3 y + J^1 d + \frac{1}{2} c_2^4 J^1 u^2 \quad (23)$$

The solution of the motion equations is obtained by differentiation with respect to c_2 and c_3 of W . Thus

$$0 = \frac{\partial W}{\partial c_2} = -c_2 t + x - 2 c_2^3 J^1 u^2 \quad (24)$$

and

$$0 = \frac{\partial W}{\partial c_2} = -c_2 t + x \quad (25)$$

Equations (24) and (25) yield the following geodesic equations

$$x = c_2 t \quad (26)$$

and

$$z = -\frac{c_3 x}{c_2} + 4J^1 c_3 z^2 \quad (27)$$

which yields the desired trajectory. From (27) one observes that torsion contributes to the parabolic curve. In the absence of the torsional defect the geodesics are reduced to the straight lines. Our result describes a gravitationally repulsive domain wall is algebraically simpler than the one obtained by F. Moraes due to the simpler geometry of the torsional defect we are dealing with. Future prospects in this work include the investigation of autoparallels in propagating torsion theories. After I finished this paper I came across with an interesting paper by Katanaev and Volovich [12] where they considered elastic waves on parallel wedge dislocations in the geometric theory of defects also dealing with the Hamilton-Jacobi methods. Nevertheless they did not handle domain walls as we did here.

Acknowledgement

I would like to express my gratitude to Prof. H. Kleinert for kindly suggesting this work to me and for his kind hospitality at the Institut für theoretische Physik at Berlin Freie Universität. Thanks are also due Prof. P. S. Letelier for helpful discussions on the subject of this Letter. Special thanks go to Prof. M. O. Katanaev and Prof. James Sethna for sending me their papers. Financial support from CNPq (Brazil) and DAAD (Bonn) is gratefully acknowledged.

References

- [1] A. Vilenkin and P.S. Shellard, Cosmic String and Other Topological Defects, (1993), Cambridge Univ. Press.
- [2] E.Kroner, Continuum theory of Defects, course 3, in Physics of Defects, Les Houches Session XXXV, North-Holland, Amsterdam, (1981).
- [3] P.S.Letelier, J.Math.Phys.(1995).
- [4] P.S.Letelier, Class. and Quant. Grav., (1995), 12, 2221.
- [5] L.C.Garcia de Andrade, J.Math.Phys.(1998), Jan.issue.
- [6] L.C.Garcia de Andrade, Mod.Phys.Lett.A, 12, 27 (1997).
- [7] L.C.Garcia de Andrade, On non-Riemannian domain walls, Gen. Rel. and Grav.(1998), in press.
- [8] F.Moraes, Phys. Lett. A 214, (1996) 189.
- [9] L.C.Garcia de Andrade, Non-Riemannian cosmic walls as boundaries of spinning polarized matter, Mod.Phys.Lett.A, in press.
- [10] E.Dubois-Violette and B.Pansu, Geometry of Condensed Matter Physics, Directions in condensed matter physics vol.9, (1990), World Scientific.
- [11] J.P.Sethna, Phys.Rev.B31, (1985), 10, 6278.
- [12] M.O.Katanaev and I.V.Volovich, Scattering on dislocations and cosmic string in the geometric theory of defects, Los Alamos preprint archives gr-qc/9801081.