

Causal Dissipative Cosmology With Variable G and Λ

Arbab I.Arbab¹ and A.Beesham²

Abstract

A cosmological model with variable G and Λ is considered in the framework of Israel-Stewart-Hiscock(ISH) causal theory. Power law as well as inflationary solutions are obtained. The gravitational constant is found to increase with time.

Keywords: Cosmology, Israel-Stewart-Hiscock Theory, Variable G and Λ

¹ICTP, P.O. Box 586, 34100-Trieste, ITALY. E-mail: arbab@ictp.trieste.it / arbab64@hotmail.com

²Department of Applied Mathematics, University of Zululand, Private Bag X1001, Kwa-Dlangezwa 3886, South Africa

The role of dissipative effects on the evolution of the universe has been investigated by many works [6,8,11,15]. Weinberg has stressed that the entropy of the early universe may have emerged from bulk viscosity dissipation [12]. The bulk viscosity may arise from the conversion of massive string modes to massless modes as well as particle creation effects around the grand unification (GUT) era [6]. It has been shown by Murphy and Barrow[8,11] that bulk viscosity can lead to inflation. In a recent paper [10], we have shown that this solution also arises in the presence of variable gravitational and cosmological constant. In the same view Singh *et al* [13] obtained a similar solution with different energy conservation. Most of these investigation are based on Eckart theory. However, it is shown that this theory is non-causal and all of it equilibrium states are unstable. The presently available theory considering dissipative effects in the universe is the full Israel-Stewart-Hiscock casual thermodynamics [2,3]. In this paper we would like to investigate the evolution of bulk viscous stress in this casual theory where both G and Λ vary with time. Our investigation shows that the variation of the gravitational constant is also consistent with the full casual theory. For age values consistent with present data the theory allows only increasing G for the present phase. We further show that the functional dependence of the coefficient of bulk viscosity assumed by many workers is an additional ad hoc assumption. We have found power law as well as inflationary solutions in the full casual theory.

The Field Equations

In a Robertson Walker(RW) flat metric, the Einstein field equations with a perfect fluid energy-momentum tensor and variable G and Λ yield

$$3\frac{\dot{R}^2}{R^2} = 8\pi G\rho + \Lambda \quad , \quad (1)$$

$$3\frac{\ddot{R}}{R} = -4\pi G(3p + \rho) + \Lambda \quad (2)$$

and

$$3(p + \rho)\dot{R} = -\left(\frac{\dot{G}}{G}\rho + \dot{\rho} + \frac{\dot{\Lambda}}{8\pi G}\right)R \quad . \quad (3)$$

The usual energy-momentum conservation, $T_{;\nu}^{\mu\nu} = 0$, leads to

$$\dot{\rho} + 3\frac{\dot{R}}{R}(p + \rho) = 0 \quad . \quad (4)$$

In the presence of bulk viscous stress (Π), the pressure term becomes $p \rightarrow p + \Pi$.

Section A

Equation (3) and (4) yields

$$\dot{\rho} + 3H(\rho + p) = -3\Pi H \quad (5)$$

and

$$8\pi\dot{G}\rho + \dot{\Lambda} = 0 \quad . \quad (6)$$

where a dot denotes a derivative with respect to time t . The density ρ and the pressure p are related by the equation of the state

$$p = (\gamma - 1)\rho , \quad (7)$$

where $\gamma=\text{constant}$. Following [10,13] we take

$$\Lambda = 3\beta H^2 . \quad (8)$$

Using eq.(6), eqs.(1) & (8) yield

$$G = BH^{-2\beta/(1-\beta)}, \quad B = \text{const.} , \quad (9)$$

$$\rho = DH^{2/(1-\beta)}, \quad D = \text{const.} \quad (10)$$

while eq.(2) gives

$$\dot{H} + \frac{3\gamma(1-\beta)}{2}H^2 = -4\pi G\Pi, \quad \beta \neq 1 . \quad (11)$$

The causal evolution equation for the bulk viscous pressure is given by [2,3]

$$\Pi + \tau\dot{\Pi} = -3\zeta H - \frac{\epsilon}{2}\tau\Pi(3H + \frac{\dot{\tau}}{\tau} - \frac{\dot{\zeta}}{\zeta} - \frac{\dot{T}}{T}), \quad (12)$$

where T is the temperature, ζ the bulk viscosity coefficient and τ the relaxation time. Setting $\epsilon = 0, 1$ gives the truncated and the full-ISH causal theory, respectively. Now consider the parameterization

$$\zeta = \tau\rho, \quad \text{and} \quad T = T_0\rho^r , \quad (13)$$

where $r = \frac{\gamma-1}{\gamma}$, T_0 are constants. Upon using eq.(9), eq.(11) yields

$$\dot{H} + \frac{3\gamma(1-\beta)}{2}H^2 = -4\pi BH^{-2\beta/(1-\beta)}\Pi, \quad \beta \neq 1 . \quad (14)$$

We will consider here two cases:

Case (1):

(a) $H = H_0 = \text{const.}$ and $\beta \neq 1$, so that $R \propto \exp(H_0 t)$, i.e., an inflationary solution. It follows from eqs.(8)-(12) that

$$\Lambda = \text{const.}, \quad G = \text{const.}, \quad \rho = \text{const.}, \quad \Pi = \text{const.}, \quad (15)$$

and

$$\zeta = \text{const.}, \quad \text{and} \quad \tau = \text{const.} \quad (16)$$

We see that the bulk viscosity remains constant during the inflationary phase. This solution is obtained by Wolf for the non-causal theory and by Banerjee and Beesham for the casual theory.

(b) $H = \text{const.}$ and $\beta = 1$: We see that $\Lambda = 3H^2, R \propto \exp(\sqrt{\frac{\Lambda}{3}}t), G = \text{const.}$ and $\rho = 0$.

From eqs.(5) and (13) it follows that $\Pi = 0$, $\zeta = 0$ and $T = 0$, so we get the perfect fluid solution. Therefore, during this de Sitter solution the bulk viscosity vanishes. Also, if $\alpha = \frac{3\gamma(1-\beta)}{2}$, $\Pi = 0$ and we recover the model in [9].

Case(2):

A power law form of R , i.e., $R = At^\alpha$, $A, \alpha = \text{const.}$. Applying this in eq.(15) one obtains,

$$\Pi = -Nt^{-2/(1-\beta)}, \beta \neq 1, \quad (17)$$

where $N = \frac{[-2+3\gamma\alpha(1-\beta)]\alpha^{(1+\beta)/(1-\beta)}}{8\pi A}$. Equations (9) and (10) give

$$\rho = B't^{-2/(1-\beta)}, G = Ct^{2\beta/(1-\beta)}, C, B' \text{ const.}, \beta \neq 1. \quad (18)$$

From the above equation we see that $\Pi = -a\rho$, $a = \text{const.}$ Using eqs.(12) and (17), eq.(13) gives

$$\tau = Bt, B = \text{const.} \quad (19)$$

and

$$\zeta = \zeta_0 t^{-(1+\beta)/(1-\beta)}, \zeta_0 = \text{const}, \beta \neq 1. \quad (20)$$

It is clear that the viscosity is decreasing with time. This solution is obtained by Desikan [5] for Brans-Dicke theory with constant deceleration parameter under the framework of ISH casual theory. This is manifested in the replacement of $[\alpha/(1+b)]$ by $-2\beta/(1-\beta)$. We see that these relations are valid irrespective of the functional dependence form of ζ on ρ . We would like to mention that Banerjee and Beesham followed a lengthy way to arrive at this result. They, however, claimed that this solution is possible only for particular behavior i.e., $\zeta \propto \rho^{1/2}$. In fact this solution is valid for all values of $[q]$, see eqs.(2.8) and (2.14)]

Section B

Another way of looking at eq.(3) is by assuming that

$$\dot{\rho} + 3\gamma H\rho = 0 \quad (21)$$

and

$$\frac{\dot{G}}{G}\rho + \frac{\dot{\Lambda}}{8\pi G} = -3\Pi H. \quad (22)$$

Using eqs.(1) and (8) this can be written as

$$\left(\frac{2}{1-\beta}\frac{\dot{H}}{H} + 3\gamma H\right)\rho = -3\Pi H. \quad (23)$$

We now assume a power law for R , i.e., $R = At^\alpha$, $A, \alpha = \text{const.}$. Equation (21) and (23) now yield

$$\Pi = -N\rho, \text{ and } \rho = Dt^{-3\alpha\gamma}, D = \text{const.}, \beta \neq 1, \quad (24)$$

where $N = \frac{-2+3\alpha\gamma(1-\beta)}{3\alpha(1-\beta)}$. Using eqs.(13) and (24), eq.(12) gives

$$\tau = Bt \quad (25)$$

where $B^{-1} = 3\alpha[\gamma(1 - \frac{\epsilon}{2}(1 + r)) + (\frac{1}{N} - \frac{\epsilon}{2})]$.

In section A, we obtained a similar relation for the relaxation time and upon using eq.(13) we obtain

$$\zeta = \tau\rho = \eta_0 t^{1-3\alpha\gamma}, \eta_0 = \text{const.} \quad (26)$$

From eqs.(1), (8) and (24), the gravitational parameter is given by

$$G \propto t^{-2+3\alpha\gamma} \quad (27)$$

Therefore, G increases or decreases with time according to whether $3\alpha\gamma > 2$ or < 2 . However, G remains constant in the pure radiation and matter epochs. We therefore, see that the casual bulk effects perturb the usual FRW solution for the radiation and matter phases. For the present era, the gravitational parameter is increasing with time(to get a reasonable age parameter for the universe).

Inflationary Solution

This is obtained when $H = H_0 = \text{const}$, i.e., $R = R_0 \exp(H_0 t)$, $R_0 = \text{const}$ In this case eq.(20) gives

$$\rho = D \exp(-3\gamma H_0 t), D = \text{const} \quad (28)$$

so that

$$G = F \exp(3\gamma H_0 t), F = \text{const.} \quad (29)$$

and from eq.(23)

$$\Pi = -\gamma\rho. \quad (30)$$

Applying this to eq.(12) we get

$$\tau = \text{const} \quad \text{and} \quad \zeta \propto \exp(-3\gamma H_0 t). \quad (31)$$

Note that during inflation both ζ and ρ decreases with time exponentially. We also note that this solution is applicable irrespective of the value of ϵ , i.e., for $\epsilon = 1$, or 0 .

Acknowledgement

AIA wishes to thank the FRD (South Africa) for support and the University of Zululand for hospitality.

References

1. N. Banerjee and A.Beesham, Pramana 46(1996)213
2. W.A.Hiscock and L.Lindblom, Ann. Phys.(NY) 100, (1976)310
3. W.Israel, J.Stewart, Physics Letter A58, (1976)213
4. P.C. Paul, S.Mukhrejee and A.Beesham,(University of Zululand Preprint-1997)
5. K.Desikan, Pramana, 45(1995)511
6. C.Wolf, S.-Afr. Tydskr, 14(1991)68
7. R.Maartens, Class. Quantum Gravit. 12(1995)1455
8. D.J. Barrow, Phy. Lett.B180(1986)335
9. A.I.Arbab, (University of Zululand Preprint-1997)
10. A.I.Arbab, Gen. Rel. Gravit.29(1997)61
11. G.Murphy, Phys. Rev.D8 (1973)4321
12. S.Weinberg, Gravitation and Cosmology (J.Wiely, NY, 1972)
13. T.Singh, A.Beesham and W.S.Mbokazi, Gen. Relat. Gravit.30(1998)573
14. C.Ekart, Phys. Rev.58(1940)919
15. O.Gron, Astr. Phys. Space Science.173(1990)191