

RELATIVISTIC CHARGED SPHERES II: REGULARITY AND STABILITY *

Fernando de Felice †, Liu Siming ‡ and Yu Yunqiang §

†Department of Physics *G. Galilei*, University of Padova
via Marzolo 8, I-35131 Padova Italy and INFN, Sezione di Padova

‡§Department of Physics, Peking University, Beijing 100871, China
(October 15, 2018)

Abstract

We present new results concerning the existence of static, electrically charged, perfect fluid spheres that have a regular interior and are arbitrarily close to a maximally charged black-hole state. These configurations are described by exact solutions of Einstein's field equations. A family of these solutions had already been found (de Felice et al., 1995) but here we generalize that result to cases with different charge distribution within the spheres and show, in an appropriate parameter space, that the set of such physically reasonable solutions has a non zero measure. We also perform a perturbation analysis and identify the solutions which are stable against adiabatic radial perturbations. We then suggest that the stable configurations can be considered as classic models of charged particles. Finally our results are used to show that a conjecture of Kristiansson et al. (1998) is incorrect.

Key words: relativity - exact solutions - black-hole physics

I. INTRODUCTION

It is well known that static, spherically symmetric, uncharged perfect fluids cannot be held in equilibrium below a certain radius without developing singularities inside. This radius is $9M/4$ for an incompressible fluid sphere of total mass energy M ; it can be even larger than that for more realistic equations of state (Buchdahl, 1959). The possibility of holding a non-singular object in stable equilibrium but compact enough to be close (in fact arbitrarily close) to a black-hole state, is of great interest not only in order to judge the state of matter in this quasi-critical condition, that is being about to turn into a black-hole, but also to yield a classic model of charged massive particles which might have astrophysical and cosmological implications. Although one can reach this goal with non-perfect fluids, a

*This work was partially supported by the Gruppo Nazionale per la Fisica Matematica del CNR and by the Ministero della Ricerca Scientifica e Tecnologica (MURST) of Italy. †: defelice@pd.infn.it, ‡: liusiming@263.net, §: yqyu@hotmail.com.

perfect fluid solution of the type mentioned was recently found (de Felice et al., 1995) but with the presence of an electric charge. Electric charges inhibit the growth of space-time curvature and therefore they are an efficient means of avoiding singularities inside matter and enhance stability even for quasi-critical configurations.

In this paper we extend the above mentioned analysis (de Felice et al., 1995) to a more general case and, in Section 2, show the existence of regular solutions very close to a black-hole state as points in a parameter space (figures 1-3). In Section 3 we study the stability of these solutions against adiabatic radial perturbations and show the domain of stable solutions in the same parameter space (figures 1, 2). Having solutions which describe very compact sources, we are able to verify the correctness of a conjecture (Kristiansson et al., 1998) according to which the limit of regular embedding in the Euclidean space of conformally reduced space-like sections of the Reissner-Nordström metric, coincides with the limit of regularity of the internal solutions. In section 4 we show that this is not true.

Throughout the paper we use geometrized units ($c = 1 = G$) and metric signature $+2$.

II. EXISTENCE OF REGULAR QUASI-CRITICAL SPHERES

In Schwarzschild coordinates the line element of a spherically symmetric space-time reads:

$$ds^2 = -e^\eta dt^2 + e^\lambda dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2$$

where $\eta = \eta(r, t)$ and $\lambda = \lambda(r, t)$. The energy-momentum tensor of a perfect, electrically charged fluid takes the form (de Felice et al., 1990):

$$T^{\mu\nu} = (\rho + p)u^\mu u^\nu + pg^{\mu\nu} + \frac{1}{4\pi}(F^{\mu\alpha}F^\nu{}_\alpha - \frac{1}{4}g^{\mu\nu}F_{\alpha\beta}F^{\alpha\beta})$$

where $F^{\alpha\beta}$ satisfies Maxwell's equations:

$$F_{[\alpha\beta,\gamma]} = 0, \quad F^{\mu\nu}{}_{;\nu} = 4\pi j^\mu.$$

Here j^μ is the current four-vector, ρ and p are the energy density and the isotropic pressure of matter measured in its rest frame respectively; ρ and p are expressed in (geometrized) units of $length^{-2}$.

In the spherically symmetric case the only non vanishing component of Maxwell's tensor is (Bekenstein, 1971):

$$F^{tr} = e^{-\frac{\lambda+\eta}{2}} \frac{Q(r, t)}{r^2}$$

where:

$$Q(r, t) = \int_0^r e^{\frac{\lambda+\eta}{2}} 4\pi r^2 j^t dr \quad (1)$$

is the total electric charge within a sphere of radius r at time t ; it then follows that:

$$F^{\alpha\beta}F_{\alpha\beta} = -\frac{2Q^2}{r^4}$$

where charge is expressed in units of *length*. If we require that the configuration be static, then $u^\mu = e^{-\eta/2} \delta_t^\mu$, hence Einstein's equations which one needs to solve for the interior metric, take the form:

$$e^{-\lambda} \left(\frac{\lambda'}{r} - \frac{1}{r^2} \right) + \frac{1}{r^2} = \frac{Q^2}{r^4} + 8\pi\rho, \quad (2)$$

$$-e^{-\lambda} \left(\frac{\eta'}{r} + \frac{1}{r^2} \right) + \frac{1}{r^2} = \frac{Q^2}{r^4} - 8\pi p \quad (3)$$

where $(')$ means derivative with respect to r .

Let the sphere be incompressible with a constant matter energy density ρ_m and the charge distribution within the sphere be given by:

$$Q(r) = Q_0 \left(\frac{r}{R} \right)^n \quad (4)$$

where n is a constant parameter, R is the coordinate radius of the boundary and Q_0 is the total charge. Integrating equation (2) with respect to coordinate radius r , one obtains

$$e^{-\lambda} = 1 - \frac{8\pi}{3} \rho_m r^2 - \frac{Q_0^2 r^{2n-2}}{(2n-1)R^{2n}}. \quad (5)$$

Combining this result with (1) and (4), we see that the charge density is given by:

$$e^{\frac{n}{2}j^t} = \frac{Q_0 n r^{n-3}}{4\pi R^n e^{\frac{\lambda}{2}}} = \frac{Q_0 n r^{n-3} \sqrt{1 - \frac{8\pi}{3} \rho_m r^2 - \frac{Q_0^2 r^{2n-2}}{(2n-1)R^{2n}}}}{4\pi R^n},$$

then we have $n \geq 3$ which prevents the divergence of charge density at the center.

Introducing scale parameters such as:

$$p_0 = \frac{1}{3} \rho_m, \quad r_0 = \left(\frac{3}{4\pi \rho_m} \right)^{1/2} \quad (6)$$

and setting:

$$Y \equiv \frac{p}{p_0}, \quad \bar{\xi} \equiv \frac{r}{r_0} \quad (7)$$

the Oppenheimer-Wolkoff equation and equation (5) for such a sphere read respectively (de Felice et al., 1995):

$$\frac{dY}{d\bar{\xi}} = n\alpha \bar{\xi}^{2n-5} - (Y+3) \frac{(Y+1)\bar{\xi} - [(n-1)/(2n-1)]\alpha \bar{\xi}^{2n-3}}{1 - 2\bar{\xi}^2 - [\alpha/(2n-1)]\bar{\xi}^{2(n-1)}}, \quad (8)$$

$$e^{-\lambda} = 1 - 2\bar{\xi}^2 - [\alpha/(2n-1)]\bar{\xi}^{2(n-1)} \quad (9)$$

where

$$\alpha = \left(\frac{Q_0 r_0^{n-1}}{R^n} \right)^2.$$

An obvious condition is $\alpha > 0$. Now we see that the set of equations (3), (8) and (9) is complete.

FIGURES




FIG. 1. Configurations in the parameter space for $n=5$

The boundary of the physical configuration is identified by the conditions:

$$Y(\bar{\xi}_b) = 0, \quad e^\nu(\bar{\xi}_b) = e^{-\lambda}(\bar{\xi}_b) = 1 - 2\bar{\xi}_b^2 - \frac{\alpha}{2n-1}\bar{\xi}_b^{2(n-1)}. \quad (10)$$

where $\bar{\xi}_b = R/r_0$. Then, for a given set of $(\alpha, \bar{\xi}_b, n)$, one can solve equations (3), (8) and (9) and obtain the structure of the corresponding configuration (Note: When one integrates equation (8) numerically from its boundary inward, pressure Y may diverge before $\bar{\xi}$ reaches zero for some values of $(\alpha, \bar{\xi}_b, n)$. Since there is a singularity in such a configuration, it is physically meaningless. But we still use the parameter $(\alpha, \bar{\xi}_b, n)$ to represent such solutions). But not all of the solutions are acceptable, two basic requirements of a regular static configuration are:

- 1: That there be no singularity inside the sphere,
- 2: That the radius of the boundary be larger than the external horizon size of the black hole with the same mass and charge.




FIG. 2. Configurations in the parameter space for $n=3$




FIG. 3. Enlargement of figure 2 around the meeting point

For a given n , each point in the $(\bar{\xi}_b^2, \alpha)$ -plane represents a solution. In figures 1-3 we plot the functions which allow one to single out the points which correspond to physical configurations.

Let M be the total mass energy of the solution, then the relations between $\bar{\xi}_b$ and the other parameters are:

$$\frac{R}{M} = \frac{1}{\bar{\xi}_b^2 \left[1 + \bar{\xi}_b^{2(n-2)} \frac{n\alpha}{2n-1} \right]}, \quad (11)$$

$$\frac{Q_0}{M} = \frac{\sqrt{\alpha}}{\bar{\xi}_b^{3-n} \left[1 + \bar{\xi}_b^{2(n-2)} \frac{n\alpha}{2n-1} \right]}. \quad (12)$$

One can show that R/M changes monotonically along the curve $Q_0/M = \text{constant}$. Let $Q_0 = M$, then (12) gives the curve b where $Q_0 = M$. Inside (figure 1, for $n > 3$) or below (figure 2, for $n = 3$) this curve, $Q_0 > M$.

The horizon sizes for the charged black hole are defined as the solutions of equation $e^\lambda(R) = 0$ and the results are $R_\pm = M \pm \sqrt{M^2 - Q_0^2}$ for a given M and Q_0 . Let $e^\lambda(\bar{\xi}_b) = 0$, (9) gives curve a where $R = R_+$. Above this curve, we have $R < R_+$, so the solutions are unacceptable. We notice this curve is tangential to the curve $Q_0 = M$ at the point $\bar{\xi}_b^2 = (n-1)/(2n-1)$, $\alpha = [(n-1)/(2n-1)]^{1-n}$. On the left hand side of this point, curve a represents configurations with their boundary at the external horizon. Consequently physical configurations must lie below this curve.

For given α and $\bar{\xi}_b^2$, one can solve equation (8) numerically to get $Y(\bar{\xi})$. We find that the central pressure $Y(0)$ will diverge as $\bar{\xi}_b^2$ increases to some value for fixed $\alpha < [(n-1)/(2n-1)]^{1-n}$, then we get the curve c $Y(0) = \text{inf}$ which approaches the tangential point of curves a and b . Above this curve, there are pressure singularities inside the configurations. So the regular static configurations must lie below this curve. (Curves d and e are for later use).

We see from the figures that, for these special models, no regular solution exists which describes a compact charged sphere having $Q_0 < M$ and radius R arbitrarily close to R_+ since any such solution would lie above curve c where a pressure singularity emerges inside the sphere. A more general result is assured by the following:

Theorem: if the total electric charge of a static perfect fluid ball is smaller than its total mass, then there is no regular static configuration having a radius arbitrarily close to the size of the external horizon (Yu et al., 1999).

As stated in (de Felice et al., 1995), the curves in figures 1-3 show that a finite region in the parameter space exists, which is the region between curves b and c , where regular solutions describing static charged balls of matter may be found with a compactness arbitrarily close to the critical one for transition to a black-hole state, namely with $R \rightarrow R_+$ and $Q_0 \rightarrow M$.




FIG. 4. The pressure of the configurations for $n=3$

It is of interest to study the behavior of the pressure inside some of these configurations as a function of parameter α . For a fixed n , this parameter gives a measure of the total charge within the body, so we show in figure 4 plots of the internal pressure as function of radius $\bar{\xi}$, for $n = 3$, $\bar{\xi}_b^2 = 0.36$ and increasing values of α (the corresponding configurations are indicated by '+' in figure 2). It is interesting to see how charge contributes to a decrease of the central pressure, a property which justifies the very existence of regular solutions describing compact balls of matter.

Regular solutions also exist inside (for $n > 3$) or below (for $n = 3$) curve b in figures 1 and 2, where $Q_0 > M$. However, as appears from figure 4, the central pressure is negative in these solutions.

In what follows we shall discuss the stability of these solutions against adiabatic radial perturbations.

III. STABILITY

We will consider small radial perturbations for the charged, perfect fluid balls, then any fluid element which was at r in the unperturbed configuration is displaced to $r + \xi(r, t)$ in the perturbed one where $\xi \ll r$. In what follows we shall use subscript " i " to denote the unperturbed configuration, then we have (Misner et al., 1973):

$$\begin{aligned}\eta(r, t) &= \eta_i(r) + \delta\eta(r, t) \\ \lambda(r, t) &= \lambda_i(r) + \delta\lambda(r, t) \\ p(r, t) &= p_i(r) + \delta p(r, t)\end{aligned}$$

$$\begin{aligned}\rho(r, t) &= \rho_i(r) + \delta\rho(r, t) \\ \bar{n}(r, t) &= \bar{n}_i(r) + \delta\bar{n}(r, t)\end{aligned}\tag{13}$$

where \bar{n} is the baryon number density; the variations are small and such that $\delta a/a_i \sim \xi/r$, a being any of the above parameters. Furthermore we assume that the charge distribution in the vibrating configuration remains unchanged, namely $Q(r + \xi(r, t)) = Q_i(r)$, meaning that there are no electric currents for the comoving observer. To first order in the variations, the Lagrangian perturbations read:

$$\begin{aligned}\Delta p(r, t) &= p(r + \xi, t) - p_i(r) \\ &= \delta p(r, t) + p_i' \xi \\ \Delta \rho(r, t) &= \delta \rho(r, t) + \rho_i' \xi \\ \Delta \bar{n}(r, t) &= \delta \bar{n}(r, t) + \bar{n}_i' \xi \\ \Delta Q(r, t) &= 0.\end{aligned}\tag{14}$$

From baryon conservation we have:

$$\frac{d\Delta\bar{n}}{d\tau} = -\bar{n}u^\mu{}_{;\mu}.\tag{15}$$

Since $u^r/u^t = \dot{\xi}$, the dot meaning differentiation with respect to the coordinate time t and $u^\mu u_\mu = -1$, in the perturbations we have to first order:

$$\begin{aligned}u^t &= e^{-\eta_i/2} \left(1 - \frac{\delta\eta}{2}\right) \\ u^r &= e^{-\eta_i/2} \dot{\xi}.\end{aligned}\tag{16}$$

Using the relation

$$u^\mu{}_{;\mu} = (-g)^{-1/2} [(-g)^{1/2} u^\mu]_{,\mu}$$

and integrating equation (15) with respect to coordinate time (first-order analysis), we deduce:

$$\Delta\bar{n} = -\bar{n}_i \left[r^{-2} e^{-\lambda_i/2} \left(r^2 e^{\lambda_i/2} \xi \right)' + \frac{\delta\lambda}{2} \right].\tag{17}$$

For adiabatic variations we have:

$$\frac{\Delta p}{\Delta\bar{n}} = \frac{p_i}{\bar{n}_i} \gamma$$

where γ is the adiabatic index, then from (17) and (14) we get:

$$\delta p = -\gamma p_i \left[r^{-2} e^{-\lambda_i/2} \left(r^2 e^{\lambda_i/2} \xi \right)' + \frac{\delta\lambda}{2} \right] - \xi p_i'.\tag{18}$$

From Einstein's equations, we obtain:

$$-e^{-\lambda_i} \frac{r\eta_i' + 1}{r^2} \delta\lambda + e^{-\lambda_i} \frac{(\delta\eta)'}{r} = \frac{2Q_i Q_i'}{r^4} \xi + 8\pi \delta p\tag{19}$$

$$e^{-\lambda_i} \frac{(\dot{\delta\lambda})}{r} = -8\pi(\rho_i + p_i) \dot{\xi} = -\frac{e^{-\lambda_i}}{r} (\lambda_i' + \eta_i') \dot{\xi}\tag{20}$$

thus, integrating (20) with respect to coordinate time and choosing the integration constant so that $\delta\lambda = 0$ when $\xi = 0$, we get:

$$\delta\lambda = -e^{\lambda_i} 8\pi r(\rho_i + p_i)\xi = -(\lambda_i' + \eta_i')\xi. \quad (21)$$

Using (21), (18) and (3) to eliminate respectively $\delta\lambda$, δp and η_i' in (19), we have:

$$\begin{aligned} (\delta\eta)' &= e^{\lambda_i} r \left(\frac{2Q_i Q_i'}{r^4} \xi + 8\pi \delta p \right) + \frac{r\eta_i' + 1}{r} \delta\lambda \\ &= -8\pi \gamma p_i \frac{e^{\lambda_i + \eta_i/2}}{2} \left(r^2 e^{-\eta_i/2} \xi \right)' + 8\pi [p_i' r - (p_i + \rho_i)] e^{\lambda_i} \xi - \frac{2Q_i Q_i'}{r^3} \xi e^{\lambda_i}. \end{aligned} \quad (22)$$

If we project the identity:

$$T^\nu_{\mu;\nu} = 0 \quad (23)$$

parallelly to the matter flow, namely:

$$u^\mu T^\nu_{\mu;\nu} = 0,$$

we obtain, from the chosen forms of u^μ and $T^{\mu\nu}$:

$$-\frac{d\rho}{d\tau} + \frac{(\rho + p)}{\bar{n}} \frac{d\bar{n}}{d\tau} - \frac{Q}{4\pi r^4} \frac{dQ}{d\tau} = 0. \quad (24)$$

Integrating (24) with respect to time and using (14) and (17) to remove ΔQ , $\Delta\rho$ and $\Delta\bar{n}$ we deduce after some algebra:

$$\delta\rho = -(\rho_i + p_i) \left[r^{-2} e^{-\lambda_i/2} \left(r^2 e^{\lambda_i/2} \xi \right)' + \frac{\delta\lambda}{2} \right] - \xi \rho_i'. \quad (25)$$

If we now project (23) transversely to the matter flow, namely:

$$h^\mu_\nu T^\sigma_{\mu;\sigma} = 0,$$

where $h^\mu_\nu = \delta^\mu_\nu + u^\mu u_\nu$, we obtain:

$$(\rho + p) u^\mu u_{\nu;\mu} = -p_{,\nu} - (T^\mu_{\nu;\mu})^{em} - u_\nu \frac{Q}{4\pi r^4} \frac{dQ}{d\tau}. \quad (26)$$

The r -component of (26) yields:

$$(\rho_i + p_i) e^{\lambda_i - \eta_i} \ddot{\xi} = -(\delta p)' - (\delta\rho + \delta p) \frac{\eta_i'}{2} - (\rho_i + p_i) \frac{(\delta\eta)'}{2} - \frac{Q_i Q_i'}{4\pi r^4} \xi' - \frac{Q_i^2}{4\pi r^4} \xi, \quad (27)$$

while the t -component is trivial and gives:

$$p_i' = \frac{Q_i Q_i'}{4\pi r^4} - (\rho_i + p_i) \frac{\eta_i'}{2}.$$

By using the initial value equations (18), (25) and (22) to reexpress δp , $\delta\rho$ and $(\delta\eta)'$ in terms of ξ in equation (27), we obtain:

$$\begin{aligned}
(\rho_i + p_i)e^{\frac{3}{2}\lambda_i}\ddot{\xi} &= \left\{ \left[\gamma p_i r^{-2} e^{\eta_i/2} \zeta' + \xi p_i' \right]' + \left[(\rho_i + p_i(1 + \gamma)) r^{-2} e^{\eta_i/2} \zeta' + \xi(\rho_i' + p_i') \right] \frac{\eta_i'}{2} \right. \\
&\quad + (\rho_i + p_i) \left[4\pi \gamma p_i r^{-1} e^{\lambda_i + \eta_i/2} \zeta' + \frac{Q_i Q_i'}{r^3} \xi e^{\lambda_i} + 4\pi(\rho_i + p_i) e^{\lambda_i} \xi - 4\pi p_i' r e^{\lambda_i} \xi \right] \\
&\quad \left. - \frac{Q_i Q_i'}{4\pi r^4} \zeta' - \frac{Q_i'^2}{4\pi r^4} \xi \right\} e^{\eta_i + \lambda_i/2} \\
&= \left[\gamma p_i r^{-2} e^{\lambda_i/2 + 3\eta_i/2} \zeta' \right]' + e^{\lambda_i/2 + 3\eta_i/2} \zeta \left[\frac{(p_i' - (Q_i Q_i')/(4\pi r^4))^2}{r^2(\rho_i + p_i)} \right. \\
&\quad + \frac{4}{r^3} \left(\frac{Q_i Q_i'}{4\pi r^4} - p_i' \right) - 8\pi p_i(\rho_i + p_i) \frac{e^{\lambda_i}}{r^2} - \frac{e^{\lambda_i}}{r^6} Q_i^2(\rho_i + p_i) \\
&\quad \left. + \frac{Q_i Q_i''}{4\pi r^6} - \frac{Q_i Q_i'}{\pi r^7} \right]
\end{aligned} \tag{28}$$

where $\zeta = r^2 e^{-\eta_i/2} \xi$ is a *renormalized displacement function*. From the above, the physically acceptable solutions of the dynamic equation must satisfy the following boundary conditions:

$$\frac{\xi}{r} \quad \text{finite or zero as} \quad r \rightarrow 0 \tag{29}$$

$$\Delta p = -\gamma p_i r^{-2} e^{\eta_i/2} \left(r^2 e^{-\eta_i/2} \xi \right)' \rightarrow 0 \quad \text{as} \quad r \rightarrow R \tag{30}$$

where R is the radius of the ball.

Assume that $\zeta(r, t)$ has a sinusoidal time dependence:

$$\zeta(r, t) = \zeta(r) e^{-i\omega t} \tag{31}$$

hence the dynamic equation and the boundary conditions reduce to an eigenvalue problem for the angular frequency ω and amplitude $\zeta(r)$:

$$(F\zeta')' + (H + \omega^2 W)\zeta = 0 \tag{32}$$

so that ζ/r^3 is finite or zero as $r \rightarrow 0$ and $\gamma p_i r^{-2} e^{\eta_i/2} \zeta'(r) \rightarrow 0$ as $r \rightarrow R$, where:

$$F = \gamma p_i r^{-2} e^{\lambda_i/2 + 3\eta_i/2} \tag{33}$$

$$\begin{aligned}
H &= e^{\lambda_i/2 + 3\eta_i/2} \left[\frac{(p_i' - (Q_i Q_i')/(4\pi r^4))^2}{r^2(\rho_i + p_i)} + \frac{4}{r^3} \left(\frac{Q_i Q_i'}{4\pi r^4} - p_i' \right) - 8\pi p_i(\rho_i + p_i) \frac{e^{\lambda_i}}{r^2} \right. \\
&\quad \left. - \frac{e^{\lambda_i}}{r^6} Q_i^2(\rho_i + p_i) + \frac{Q_i Q_i''}{4\pi r^6} - \frac{Q_i Q_i'}{\pi r^7} \right]
\end{aligned} \tag{34}$$

$$W = (\rho_i + p_i) r^{-2} e^{3\lambda_i/2 + \eta_i/2} \tag{35}$$

By using the variational principle (Mathews J. and Walker R.L., 1970), one gets:

$$\omega^2 = \text{extreme of} \quad \frac{\int_0^R (F\zeta'^2 - H\zeta^2) dr}{\int_0^R W\zeta^2 dr} \tag{36}$$

so, the configuration is stable against adiabatic radial perturbations only if:

$$\int_0^R (F\zeta'^2 - H\zeta^2)dr > 0 \quad (37)$$

for any ζ which satisfies the boundary conditions (29), (30).

Using the simplest *trial function* $\zeta \propto r^3$ (Chandrasekhar, 1964) and assuming that the adiabatic index γ was constant throughout the spheres, we calculated the stability of such configurations. The domain of stable solutions in the parameter space is shown in figures 1 and 2:

curve *d* for $\gamma=4$, the configurations are unstable above this curve,

curve *e* for $\gamma=5/3$, the configurations are unstable above this curve,

and the stability curves approach curve *c* as the adiabatic index approaches infinity. Evidently this limit corresponds to perfect rigidity.

With the inclusion of these curves, figures 1 and 2 allow us to identify a region on the parameter space which corresponds to solutions which are physically significant, being regular and stable against adiabatic radial perturbations. Since these very compact configurations are electrically charged, they are probably not describing astrophysical objects but, perhaps, classic models of massive elementary particles. In this case it would be desirable to extend our analysis to a more detailed modelling of the interior, since the adiabatic index depends critically on the equation of state (Akmal et al., 1998).

IV. REGULAR SOLUTIONS AND REGULAR EMBEDDINGS

Given a Riemannian manifold M with Lorentzian metric g , it may be heuristically helpful to find embedding diagrams of any subset of M into Euclidean space. More specifically, one has to find a surface in a three-Euclidean space which is isometric to a given two-surface of M (Bini et al., 1999). This problem does not always have a solution. It has been argued (Kristiansson et al., 1998) that the limiting condition for regular embedding of space-like sections of the vacuum Schwarzschild and Reissner-Nordström solutions, conformally reduced by the factor $\Omega = g_{00}$, does coincide with the limiting condition for their regular, static, spherically symmetric internal solutions. We shall show here that this is not true.

Consider the Reissner-Nordström space-time solution:

$$ds^2 = - \left(1 - \frac{2m}{r} + \frac{Q_0^2}{r^2}\right) dt^2 + \left(1 - \frac{2m}{r} + \frac{Q_0^2}{r^2}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (38)$$

then consider the conformal $t = \text{constant}$ space-like subspace:

$$d\Sigma^2 = \left(1 - \frac{2m}{r} + \frac{Q_0^2}{r^2}\right)^{-2} dr^2 + \left(1 - \frac{2m}{r} + \frac{Q_0^2}{r^2}\right)^{-1} r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (39)$$

The two-dimensional surface $\theta = \pi/2$, namely:

$$d\sigma^2 = \left(1 - \frac{2m}{r} + \frac{Q_0^2}{r^2}\right)^{-2} dr^2 + \left(1 - \frac{2m}{r} + \frac{Q_0^2}{r^2}\right)^{-1} r^2 d\phi^2 \quad (40)$$

is matched to a surface $z = z(R)$ in the three-dimensional Euclidean space:

$$\begin{aligned}
d\sigma_e^2 &= dR^2 + R^2 d\phi^2 + dz^2 \\
&= \left[1 + \left(\frac{dz}{dR} \right)^2 \right] dR^2 + R^2 d\phi^2
\end{aligned} \tag{41}$$

where R, ϕ, z are cylindrical coordinates. Comparing (41) with (40) we obtain the condition for regular embedding as:

$$\left(\frac{dz}{dR} \right)^2 = \left(1 - \frac{2m}{r} + \frac{Q_0^2}{r^2} \right) \left(1 - \frac{3m}{r} + \frac{2Q_0^2}{r^2} \right)^{-2} - 1 \geq 0 \tag{42}$$

From this we deduce, for each M and Q_0 , the condition for regular embedding as $r \geq \tilde{r}$ where \tilde{r} is the solution of the equation:

$$1 - \frac{2m}{\tilde{r}} + \frac{Q_0^2}{\tilde{r}^2} = \left(1 - \frac{3m}{\tilde{r}} + \frac{2Q_0^2}{\tilde{r}^2} \right)^2 \tag{43}$$




FIG. 5. Configurations in the parameter space for different n and enlargement

If $Q_0 = 0$, we easily deduce $\tilde{r} = 9M/4$, a well known limit for regular internally static, electrically neutral, spherically symmetric and perfect fluid solutions (Buchdahl, 1959). If $Q_0 \neq 0$, there are several internal solutions which match the external Reissner-Nordström one. Here, of the various solutions we shall consider those with $n = 3$ and $n = \infty$. The plots of the limit of regularity for these solutions, namely where the central pressure diverges, are compared in the $(r - Q_0)$ -plane and shown in Figure 5 where r is the coordinate radius of the sphere's boundary. Although the two curves coincide at $r = 9M/4$ when $Q_0 = 0$ and $r = M$ when $Q_0 = M$, as expected, they differ from each other everywhere else, curve $n = \infty$ being at larger radii than curve $n = 3$ for each value of Q_0 . Now it happens that the condition for infinite central pressure in the $n = \infty$ case (a perfect conductor) can be expressed analytically through solving the Oppenheimer-Wolkoff equation for conductive incompressible perfect fluids and reads:

$$\sqrt{1 - \frac{2m}{\tilde{r}} + \frac{Q_0^2}{\tilde{r}^2}} = \frac{\tilde{r}m - Q_0^2}{3\tilde{r}m - 2Q_0^2}. \tag{44}$$

This coincides with condition (43) for regular embedding. It is then clear that regular solutions for $n = 3$ exist which have radii smaller than the corresponding limit of regular embedding; this is sufficient to falsify the conjecture.

V. CONCLUSIONS

The use of plots in Figures 1 to 5 highlights the result that regular solutions of Einstein's equations exist describing charged spheres arbitrarily close to a black-hole. These solutions fill a region in a parameter space which is thin but finite so it is not a set of measure zero. However not all these solutions are stable against adiabatic radial perturbations. We have made a perturbation analysis and have established a stability criterion given by relation (37). We have deduced the stability curves for configurations with adiabatic indices $\gamma = 5/3$ and $\gamma = 4$ (respectively curves e and d in figures (1) and (2)), assuming that γ was constant throughout the sphere. We see that, for moderate values of γ , the limits of stability prevent stable compact charges from being arbitrarily close to a black-hole state but also from having a total charge Q_0 arbitrarily close to their mass. These situations would be possible if the adiabatic index diverged; in this case, in fact, the stability curve approaches curve c which, we recall, is the limit of regularity for our solutions. Clearly, a diverging γ corresponds to a complete rigidity while, in a less extreme case, γ would critically depend on the equation of state (Akmal et al., 1998).

Although our solutions can be thought of as being classic (non quantum) models of charged particles, it would be interesting to find solutions of Einstein's equations which describe charged spheres with a more detailed physical description of their interior.

VI. ACKNOWLEDGMENTS

One of us (F.de F.) would like to thank the Director of the Physics Department of Peking University for hospitality. We thank Prof. Mary Evans Properi for correcting the English text.

REFERENCES

- [1] Akmal A, Pandharipande V R and Ravenhall D G 1998 Phys. Rev. C **58** (1998) 1804
- [2] Bekenstein J D 1971 Phys. Rev. D **4** 2185
- [3] Bini Donato and Jantzen Robert T *Circular Orbits in Kerr Space Times: Embedding Diagrams*: To appear
- [4] Buchdahl H A 1959 Phys. Rev. **116** 1027
- [5] Chandrasekhar S 1964 Phys. Rev. Letters **12** 114
- [6] de Felice F and Clarke C J S 1990 *Relativity on Curved Manifolds* Cambridge Univ. Press
- [7] de Felice F, Yu Yunqiang and Fang Jing 1995 Mon. Not. R. astron. Soc. **277** L17
- [8] Kristiansson S, Sonogo S and Abramowicz M 1998 Gen. Rel. & Grav. **30** 275
- [9] Mathews J and Walker R L 1970 *Mathematical Methods of Physics* The Benjamin/Cummings Publishing Company
- [10] Misner C W, Thorne K S and Wheeler J A 1973 *Gravitation* W. H. Freeman and Co.
- [11] Yunqiang Yu and Siming Liu 1999 *Relativistic Charged Balls* : To appear.