

de Broglie-Bohm Interpretation for Analytic Solutions of the Wheeler-DeWitt Equation in Spherically Symmetric Space-time

Masakatsu Kenmoku *

Department of Physics, Nara Women's University, Nara 630-8506, Japan

Hiroto Kubotani †

Faculty of Engineering, Kanagawa University, Yokohama 221-8686, Japan

Eiichi Takasugi ‡

Department of Physics, Osaka University, Osaka 560-0043, Japan

Yuki Yamazaki §

Graduate School of Human Culture, Nara Women's University, Nara 630-8506, Japan

(August 19, 2019)

We discuss the implications of a wave function for quantum gravity, which involves nothing but 3-dimensional geometries as arguments and is invariant under general coordinate transformations. We derive an analytic wave function from the Wheeler-DeWitt equation for spherically symmetric space-time with the coordinate system arbitrary. The de Broglie-Bohm interpretation of quantum mechanics is applied to the wave function. In this interpretation, deterministic dynamics can be yielded from a wave function in fully quantum regions as well as in semiclassical ones. By introducing a coordinate system additionally, we obtain a cosmological black hole picture in compensation for the loss of general covariance. Our analysis shows that the de Broglie-Bohm interpretation gives quantum gravity an appropriate prescription to introduce coordinate systems naturally and extract information from a wave function as a result of breaking general covariance.

03.65.Bz, 04.60.-m, 04.70.Dy, 97.60.Lf

I. INTRODUCTION

Quantum theory of gravity is one of the most attractive subjects in particle physics and cosmology. The canonical formalism of gravity formulated by Arnowitt, Deser, and Misner (ADM) [1] and by Dirac [2] describes the dynamics of gravity as a totally constraint system. For quantization, the constraints are used to construct possible quantum states. The equation which corresponds to the Hamiltonian constraint is called the Wheeler-DeWitt equation [3,4]. A great deal of effort has been made to analyze this equation [5]. Especially the dynamics of spherically symmetric geometries has been studied extensively [6–8]. In this case, although gravitational and electromagnetic waves cannot be realized, we can treat heuristic geometrical structures of space-time: black holes as small scale structures and expanding universe as large scale one.

In solving the Wheeler-DeWitt equation, there arise the problem of operator ordering ambiguity and the regularization problem [4,9,10]. It has been argued that although the constraint algebra does close in certain operator ordering, it does not necessarily close in another ordering. In our previous paper [11], we considered spherically symmetric geometry and fixed operator ordering in the Hamiltonian, momentum, and mass constraints so that the algebra among these constraints is closed. For the consistent ordering, we also found an analytic wave function of the quantized spherically symmetric space-time as a simultaneous solution of the constraint equations.

To proceed further, in this paper, we try to extract quantum properties of the space-time from the analytic wave function [11,12]. For this aim, we use the de Broglie-Bohm (dBB) interpretation (quantum potential interpretation or pilot wave approach) of quantum mechanics [13–15]. In this interpretation, a deterministic rigid trajectory in the

*e-mail address: kenmoku@phys.nara-wu.ac.jp

†e-mail address: kubotani@yukawa.kyoto-u.ac.jp

‡e-mail address: takasugi@het.phys.sci.osaka-u.ac.jp

§e-mail address: yamazaki@phys.nara-wu.ac.jp

configuration space, what we call *a de Broglie-Bohm trajectory*, is defined well. The interpretation merely assigns the gradient of the phase of a wave function to the momentum of a quantum particle as

$$m \frac{dx}{dt} = p \equiv \frac{\partial \Theta}{\partial x}, \quad (1.1)$$

where the wave function is expressed as $\Psi = |\Psi| \exp(i\Theta)$. A dBB trajectory is obtained by integrating Eq. (1.1) with respect to t . There can exist many trajectories specified by the configuration on which the quantum system is set at an initial time. The amplitude $|\Psi|^2$ is interpreted as the probability density in a statistical ensemble of the trajectories. For the potential problem of quantum particles, the equation to determine the phase of a wave function is derived from the Schrödinger equation. It is described as a form of the Hamilton-Jacobi type equation, which distinctively includes *a quantum potential*

$$V_Q = -\frac{\hbar^2}{2|\Psi|} \frac{\partial^2}{\partial x^2} |\Psi|. \quad (1.2)$$

A similar equation is derived also for gravity from the Wheeler-DeWitt equation.

The dBB interpretation of quantum mechanics is favorable especially for quantum theory of gravity because this interpretation is able to resolve the conceptual problems of quantum gravity: the disappearance of the dynamics (the loss of time) [16] and the problem of the observation [17]. The dBB interpretation introduces the time parameter t through Eq. (1.1), which surmounts the problem of time [18–20]. Further, a rigid trajectory can be traced without any observation even in fully quantum regions and connects the regions smoothly to semi-classical regions, where the quantum potential is merely negligible and the trajectory behaves like a classical path.

This paper is organized as follows. In Sec. 2, we prepare for the main purpose of this paper. First we perform canonical quantization of the Einstein-Maxwell theory with a cosmological constant in spherically symmetric space-time. Then we derive an analytic solution of the Wheeler-DeWitt equation. For the space-time with no electromagnetic field, we obtained one in our previous paper [11]. Sec. 3 is devoted to introducing the dBB interpretation to the analytic wave function. In Sec. 4, we present the explicit expressions of the dBB trajectories. Here we have to give coordinate conditions. Summary of this paper is given in Sec. 5.

II. CANONICAL QUANTIZATION OF THE EINSTEIN-MAXWELL THEORY IN SPHERICALLY SYMMETRIC SPACE-TIME

In this section we consider canonical quantization of the 4-dimensional Einstein-Maxwell theory with a cosmological term. Space-time is assumed to be spherically symmetric. We quantize the space-time following our previous work [11], where we fixed operator ordering in the Hamiltonian, momentum, and mass constraints and showed that they form a closed algebra. We also derive an analytic wave function which satisfies these constraints. Here we extend our previous model [11] to include the electromagnetic field. The inclusion of the electromagnetic field will be profitable to examine extreme black holes [21] and cosmological black holes [22]. We use natural geometrical units $c = \hbar = G = 1$ and adopt the conventions in the Kuchař's work [7] and ours [11].

A. Canonical quantization

We start to consider a general spherically symmetric metric in the ADM decomposition,

$$ds^2 = -N^2 dt^2 + \Lambda^2 (dr + N^r dt)^2 + R^2 d\Omega^2, \quad (2.1)$$

where $d\Omega$ is a line element on the unit sphere, and the metric components N , N^r , Λ , and R are functions of the time coordinate t and the radial one r . Hereafter we refer to N and N^r as the lapse and shift functions, respectively, which express the degrees of freedom in the choice of coordinate systems. The action of the Einstein-Maxwell theory has a form of

$$I = \frac{1}{16\pi} \int d^4x \sqrt{-^{(4)}g} (^{(4)}R - 2\lambda - F_{\mu\nu} F^{\mu\nu}), \quad (2.2)$$

where λ denotes a cosmological term, and $^{(4)}R$ and $^{(4)}g$ are the scalar curvature and the determinant of the metric tensor, respectively. Here the electromagnetic field strength is denoted by $F_{\mu\nu}$. However, spherical symmetry reduces

the number of non-vanishing components of $F_{\mu\nu}$. The nontrivial part is described as $F_{01} = -F_{10} = \dot{A}_1 - A_0'$, where A_0 and A_1 are the components of the electromagnetic field. Hereafter a dot and a prime denote the derivatives with respect to t and r , respectively. Substituting the metric (2.1) into Eq. (2.2), we transform the action into a form of the ADM decomposition

$$I = \int dt \int dr [-N^{-1} (R(-\dot{\Lambda} + (\Lambda N^r)')(-\dot{R} + R'N^r) + \frac{1}{2}\Lambda(-\dot{R} + R'N^r)^2) + N(-\Lambda^{-1}RR'' - \frac{1}{2}\Lambda^{-1}R'^2 + \Lambda^{-2}\Lambda'RR' + \frac{1}{2}\Lambda(1 - \lambda R^2)) + \frac{1}{2}N^{-1}\Lambda^{-1}R^2(\dot{A}_1 - A_0')^2] \quad (2.3)$$

$$= \int dt \int dr [P_\Lambda \dot{\Lambda} + P_R \dot{R} + P_A \dot{A}_1 - N(\frac{1}{2}\Lambda R^{-2}P_\Lambda^2 - R^{-1}P_R P_\Lambda + \frac{1}{2}R'^2\Lambda^{-1} + RR''\Lambda^{-1} - RR'\Lambda^{-2}\Lambda' - \frac{\Lambda}{2}(1 - R^{-2}P_A^2 - \lambda R^2)) - N^r(R'P_R - \Lambda(P_\Lambda)') - A_0\hat{P}_A'], \quad (2.4)$$

where P_Λ , P_R , and P_A are the canonical momenta conjugated to Λ , R , and A_1 , respectively.

We quantize the dynamical variables Λ , R , and A_1 . In the Schrödinger picture, their canonical momenta are represented by functional differential operators:

$$\begin{aligned} \hat{P}_\Lambda(r) &= -i \frac{\delta}{\delta \Lambda(r)}, \\ \hat{P}_R(r) &= -i \frac{\delta}{\delta R(r)}, \\ \hat{P}_A(r) &= -i \frac{\delta}{\delta A_1(r)}. \end{aligned} \quad (2.5)$$

Here and in the rest of this section, we use the notation “hat” for differential operators and do not express the argument t explicitly, because we always treat products of simultaneous operators only.

B. Constraint equations

The action (2.4) includes the non-dynamical variables N , N^r , and A_1 . Variation with respect to them induces the classical constraint equations: the Hamiltonian and momentum constraints and the Gaussian law. The constraints are adapted to restrict a wave function of quantum gravity, Ψ in a form of the equations

$$\begin{aligned} \hat{H}\Psi &= 0, \\ \hat{H}_r\Psi &= 0, \\ \hat{H}_A\Psi &= 0, \end{aligned} \quad (2.6)$$

where the concrete expressions for the operators \hat{H} , \hat{H}_r , and \hat{H}_A are given later. The first equation is called the Wheeler-DeWitt equation, and the third one corresponds to the conservation of electric charge. For spherically symmetric space-time with no matter, we can introduce the mass of a black hole, M as a dynamical variable [6,7,23], which is a constant of motion. In quantum theory, we can construct mass eigenstates through

$$\hat{M}\Psi = m\Psi, \quad (2.7)$$

where \hat{M} and m are the quantized mass operator and a mass eigenvalue, respectively.

In quantum theory of gravity, it is troublesome to fix operator ordering. First we introduce the Hamiltonian, momentum, electric charge, and mass operators with ordering factors as

$$\begin{aligned} \hat{H} &= \frac{1}{2}\Lambda R^{-2}\hat{P}_\Lambda^{(C)}\hat{P}_\Lambda - R^{-1}\hat{P}_R\Lambda\hat{P}_\Lambda^{(B)}\Lambda^{-1} + \frac{1}{2}R'^2\Lambda^{-1} \\ &\quad + RR''\Lambda^{-1} - RR'\Lambda^{-2}\Lambda' - \frac{\Lambda}{2}\left(1 - \hat{P}_A^2R^{-2} - \lambda R^2\right), \end{aligned} \quad (2.8)$$

$$\hat{H}_r = R' \hat{P}_R - \Lambda(\hat{P}_\Lambda)', \quad (2.9)$$

$$\hat{H}_A = -(\hat{P}_A)', \quad (2.10)$$

$$\hat{M} - m = \frac{1}{2} R^{-1} \hat{P}_\Lambda^{(A)} \hat{P}_\Lambda - \frac{1}{2} R(\chi - \hat{F}), \quad (2.11)$$

where

$$\chi \equiv R'^2 \Lambda^{-2}, \quad (2.12)$$

$$\hat{F} \equiv 1 - 2mR^{-1} + \hat{P}_A^2 R^{-2} - \frac{\lambda}{3} R^2. \quad (2.13)$$

In Eqs. (2.8) and (2.11), the momentum \hat{P}_Λ is accompanied with an ordering function A as

$$\begin{aligned} \hat{P}_\Lambda^{(A)} &\equiv A \hat{P}_\Lambda A^{-1}, \\ \hat{P}_\Lambda^{(B)} &\equiv \frac{1}{2} \left(\hat{P}_\Lambda + \hat{P}_\Lambda^{(A)} \right) = A^{1/2} \hat{P}_\Lambda A^{-1/2}, \\ \hat{P}_\Lambda^{(C)} &\equiv \hat{P}_\Lambda^{(A)} - i R R'^{-1} \left(A^{-1} \frac{\delta A}{\delta \Lambda} \right)' = C \hat{P}_\Lambda C^{-1}, \end{aligned} \quad (2.14)$$

where

$$C = A \exp \left[- \int^r dr R R'^{-1} \int^\Lambda d\Lambda \left(A^{-1} \frac{\delta A}{\delta \Lambda} \right)' \right]. \quad (2.15)$$

In this ordering, \hat{M} has a favorable relation

$$\hat{M}' = -\Lambda^{-1} R' \hat{H} - R^{-1} \hat{P}_\Lambda^{(B)} \Lambda^{-1} \hat{H}_r - R^{-1} \hat{P}_A \hat{H}_A, \quad (2.16)$$

which guarantees that \hat{M} is spatially conserved as well as in classical theory. It shows that we have provided the quantized constraints (2.8) - (2.11) with consistent ordering.

Next we fix the ordering function A so that \hat{H} , \hat{H}_r , and \hat{M} forms a closed algebra. According to [11], the commutators between \hat{H} and \hat{H} or \hat{H}_r are evaluated from the commutators between \hat{M} and \hat{M} or \hat{H}_r , and A is shown to take a form

$$A = A_Z(Z) \bar{A}(R, \chi), \quad (2.17)$$

where A_Z and \bar{A} are arbitrary functions, and the argument Z is defined as

$$Z \equiv \int dr \Lambda f(R, \chi) = \int dr \int^\Lambda d\Lambda \bar{f}(R, \chi) \quad (2.18)$$

by an arbitrary function, f or \bar{f} . Here f and \bar{f} have a relation

$$f(R, \chi) = -\frac{\chi^{1/2}}{2} \int^\chi dx x^{-3/2} \bar{f}(R, x). \quad (2.19)$$

In the proof of the closure of the constraint algebra, we make the most of a special property that

$$[Z, H_r(r)] = i \left(R'(r) \frac{\delta Z}{\delta R(r)} - \Lambda(r) \left(\frac{\delta Z}{\delta \Lambda(r)} \right)' \right) = 0. \quad (2.20)$$

Tsamis and Woodard [10] pointed out that the closure of the constraints in quantum gravity is ill-defined in the sense that formal evaluation of the commutator yields the multiple product of delta functions, and that the theory must be regulated. Especially, the formal adoption of calculation rules for the delta function to the multiple product of delta functions with the same argument leads even to contradictory results. In our evaluation of the commutator [11], on the other hand, we only have to utilize the commutators of \hat{M} and thus need not treat the multiple product of delta functions with the same argument. Only multiple products of delta functions in our calculation comprise $\delta(0)$ that comes from the commutation of coincident operators. Further, all the noncancelable ones are multiplied by the constraints in the canonical gravity. Thus, we hopefully proceed further through the formal manipulation of the unregulated theory.

C. Solutions of constraint equations

The wave function Ψ is a functional of the electromagnetic field A_1 and the geometrical fields Λ and R . The wave function is assumed to be in a separable form

$$\Psi = \Psi_{EM}[A_1]\Psi_G[\Lambda, R] . \quad (2.21)$$

Each of $\Psi_{EM}[A_1]$ and $\Psi_G[\Lambda, R]$ is constructed as follows. The quantized Gaussian law $\hat{H}_A\Psi_{EM}[A_1] = 0$ is trivially satisfied by the equation

$$\hat{P}_A\Psi_{EM} = Q\Psi_{EM}, \quad (2.22)$$

whose solution is

$$\Psi_{EM}[A_1] = \exp(i \int dr Q A_1(r)). \quad (2.23)$$

Here Q is an eigenvalue of the conserved charge. Considering Eq. (2.20), a solution of the momentum constraint $\hat{H}_r\Psi_G[\Lambda, R] = 0$ is

$$\Psi_G = \Psi_G(Z), \quad (2.24)$$

where $\Psi_G(Z)$ on the right hand side is an arbitrary function of Z .

From Eq. (2.16), we see that if $\Psi_G(Z)$ satisfies the mass constraint (2.7), it simultaneously satisfies the Wheeler-DeWitt equation. Thus we only have to consider the mass constraint instead of the Wheeler-DeWitt equation. As to the ordering function \bar{A} in Eq. (2.17), we assume that

$$\frac{\delta Z}{\delta \Lambda} (\equiv \bar{f}) = \bar{A} \quad \text{and} \quad \bar{A}^2 = R^2(\chi - F(R)), \quad (2.25)$$

where $F \equiv \hat{F}(\hat{P}_A \rightarrow Q)$. This assumption reduces the differential functional equation (2.7) with Eq. (2.11) to an ordinary differential equation on Z ,

$$\frac{d^2\Psi_G}{dZ^2} - \frac{1}{A_Z} \frac{dA_Z}{dZ} \frac{d\Psi_G}{dZ} + \Psi_G = 0 . \quad (2.26)$$

The variable Z in Eq. (2.18) is also determined as

$$\begin{aligned} Z &= \int dr \int^\Lambda d\Lambda R \sqrt{\chi - F(R)} \\ &= \int dr R \Lambda \left(\sqrt{\chi - F(R)} + \frac{\sqrt{\chi}}{2} \ln \left| \frac{\sqrt{\chi - F(R)} - \sqrt{\chi}}{\sqrt{\chi - F(R)} + \sqrt{\chi}} \right| \right) . \end{aligned} \quad (2.27)$$

Finally we choose the rest of the ordering factors, A_Z such that the solution of Eq. (2.26) becomes a special function.

(i) Bessel type solutions

As the simplest case, we take

$$A_Z = Z^{2\nu-1}. \quad (2.28)$$

Then the solution of Eq. (2.26) is given by the Hankel (or Bessel) functions as

$$\Psi_G^{(\nu)}(Z) = Z^\nu (a_1 H_\nu^{(1)}(Z) + a_2 H_\nu^{(2)}(Z)) , \quad (2.29)$$

where a_1 and a_2 are integration constants.

(ii) Hypergeometric type solutions

If we choose

$$A_Z = Z^\sigma (Z - 1)^\delta, \quad (2.30)$$

the solution of Eq. (2.26) is given by the hypergeometric functions as

$$\Psi_G^{(\sigma, \delta)}(Z) = Z^{\sigma+1}(Z-1)^{\delta+1} \left\{ b_1 F(\alpha, \beta, \gamma; Z) + b_2 Z^{1-\gamma} F(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; Z) \right\}, \quad (2.31)$$

where b_1 and b_2 are integration constants, and α , β , and γ are constants under the constraints

$$\alpha\beta = \sigma + \delta + 2, \quad \alpha + \beta = \sigma + \delta + 3, \quad \gamma = \sigma + 2. \quad (2.32)$$

Each function of Eqs. (2.29) and (2.31) shows an eigenstate of the mass m . Eigenstates with different mass eigenvalues can be superposed. We also mention generality of the wave functions (2.29) and (2.31). For the Hamilton-Jacobi equation, a complete solution contains the same number of arbitrary constants as of the dynamical variables. In our model, there are two dynamical variables R and Λ at each point of r . However, the Wheeler-DeWitt equation works as if it is a kind of Hamilton-Jacobi equation with zero energy. Therefore, the zero energy condition is forced at each point of r . Further, we set the mass constraint equation (2.7) at each point. As a result, our solutions contain only the universal mass eigenvalue as an arbitrary constant unlike a complete solution of the Hamilton-Jacobi equation.

III. DE BROGLIE-BOHM INTERPRETATION

In the previous section, we obtained the wave function for spherically symmetric space-time. It should be noted that, although some assumptions are imposed in order to get the analytic formula, any coordinate condition on N and N^r has not been undertaken. For classical relativity, we need to fix a coordinate system to get an explicit expression of a space-time geometry. By using the dBB interpretation of quantum mechanics, we try to extract physical meanings from the wave function of quantum gravity. For ordinary quantum systems, as briefly seen in Sec. 1, the dBB interpretation gives us deterministic rigid trajectories with no ambiguity instead of wave functions. In this section, we show how the dBB interpretation induces a rigid space-time picture in quantum theory of gravity.

We express a wave function of quantum gravity in a polar coordinate as

$$\Psi(Z) = |\Psi(Z)| \exp(i\Theta(Z)), \quad (3.1)$$

where the phase $\Theta(Z)$ depends only on Z as well as the amplitude $|\Psi(Z)|$. In a similar way to Eq. (1.1), the derivatives of the phase Θ are identified with the canonical momenta P_Λ , P_R , and P_A which are conjugate respectively to Λ , R , and A_1 . Estimating the derivatives as

$$\begin{aligned} \frac{\delta\Theta}{\delta\Lambda} &= \frac{\delta Z}{\delta\Lambda} \frac{d\Theta}{dZ} = \bar{f} \frac{d\Theta}{dZ}, \\ \frac{\delta\Theta}{\delta R} &= \frac{\delta Z}{\delta R} \frac{d\Theta}{dZ} = \frac{\Lambda}{R'} \left(\frac{\delta Z}{\delta\Lambda} \right)', \quad \frac{d\Theta}{dZ} = \frac{\Lambda}{R'} \bar{f}' \frac{d\Theta}{dZ}, \\ \frac{\delta\Theta}{\delta A_1} &= Q, \end{aligned} \quad (3.2)$$

we obtain the equations to determine the dBB trajectory of the space-time geometry and the electromagnetic field as

$$\dot{R} - R' N^r = -\frac{N}{R} \bar{f} \frac{d\Theta}{dZ}, \quad (3.3)$$

$$R(\dot{\Lambda} - (\Lambda N^r)') + \Lambda(\dot{R} - R' N^r) = -\frac{N\Lambda}{R'} \bar{f}' \frac{d\Theta}{dZ}, \quad (3.4)$$

$$\dot{A}_1 - A'_0 = Q N \Lambda R^{-2}, \quad (3.5)$$

where $\bar{f} = R\sqrt{\chi - F(R)}$ from the assumption (2.25) and $\chi \equiv R'^2 \Lambda^{-2}$ by the definition (2.12). In the estimation (3.2), we have used Eq. (2.20). Equations (3.3)-(3.5) form simultaneous differential equations with respect to the time and radial coordinates. By comparing a solution of the equations with the classical one which is derived from the classical equation of motion, we can know quantum gravity effects on the space-time geometry qualitatively. When the difference between them is negligible, we only have to assert that the quantum space-time is reduced to the classical one spontaneously, or the classical state is realized without any observer.

Before integrating Eqs. (3.3)-(3.5), we observe them for a while. At sight we find that the equation for the electromagnetic part, (3.5) is the same as the classical one. By taking the ratio of Eq. (3.4) to Eq. (3.3), we obtain

$$\frac{\dot{\Lambda}}{\Lambda} + \frac{\dot{R}}{R} - \frac{\dot{R}}{R'} \frac{\bar{f}'}{\bar{f}} = N^{r'} + N^r (\ln \frac{\Lambda R}{\bar{f}})' . \quad (3.6)$$

We note that Eq. (3.6) doesn't include the phase Θ and the solution of Eq. (3.6) will not depend on an explicit form of the wave function. For the gravitational part, thus the correlation between the dynamical variables Λ and R corresponds to the classical path in the configuration space. By contrast to Eq. (3.6), Eq. (3.3) includes a functional form of Θ and the time slicing function N . Combined with Eq. (3.6), Eq. (3.3) is used to determine finally the dBB trajectory of the space-time geometry which is parametrized by the time t .

In order to proceed to get dBB trajectories concretely, we specify a wave function for our analysis. We adopt the Bessel type solution (2.29) with $a_1 = 0$,

$$\Psi_G^{(\nu)}(Z) = a_2 Z^\nu H_\nu^{(2)}(Z), \quad (3.7)$$

since its asymptotic behavior is simple and thus it is hopeful to derive physical meanings of the wave function. For the wave function (3.7), we can estimate $d\Theta/dZ$ analytically by using the relations $H_\nu^{(1)}(Z)^* = H_\nu^{(2)}(Z)$ on the real Z and $H_\nu^{(2)}(dH_\nu^{(1)}/dZ) - H_\nu^{(1)}(dH_\nu^{(2)}/dZ) = 4i/(\pi Z)$. The result is

$$n(Z)^{-1} \equiv -\frac{d\Theta}{dZ} = \frac{2}{\pi Z |H_\nu^{(2)}(Z)|^2}. \quad (3.8)$$

Thus $n(Z)$ is asymptotically

$$n(Z) \rightarrow 1 \quad \text{for } Z \rightarrow \infty. \quad (3.9)$$

Although $\Psi_G^{(\nu)}(Z)$ is an analytic function on the complex Z , the phase $\Theta(Z)$ is not. Therefore, we confine $n(Z)$ on the real Z and refrain from extending $n(Z)$ analytically to the complex plane. We note that the wave function (3.7) satisfies the Vilenkin's boundary condition [24], since the time derivative of R has a positive sign as seen in Eq. (3.3). It may be worthwhile to mention other choices of the wave function. If $b_1 = b_2$ or $b_1 = -b_2$, the wave function remains real or purely imaginary, and the dBB interpretation gives no dynamics. For the case $b_2 = 0$, the sign of the phase in the wave function is opposite to that in the wave function (3.7) and thus the direction of the time parameter t is reversed.

IV. DE BROGLIE-BOHM TRAJECTORY

In this section, we solve the differential equations (3.3) and (3.4) and obtain the dBB trajectories hidden in the wave function (3.7). For the equations to be simplified and integrated analytically, we assume that χ defined in Eq. (2.12) depends on r only through R :

$$\chi \equiv R'^2/\Lambda^2 = \bar{\chi}(R), \quad (4.1)$$

where $\bar{\chi}(R)$ is a function of R . The assumption requires that \bar{f} defined in Eq. (2.25) is also a function only of R . If we take a coordinate condition $N^r = 0$, Eq. (3.6) is reduced to a simple form

$$\frac{\dot{\Lambda}}{\Lambda} + \frac{\dot{R}}{R} - \frac{\dot{\bar{f}}}{\bar{f}} = 0. \quad (4.2)$$

By integrating Eq. (4.2), we obtain a relation

$$\bar{f} = c_0(r) R \Lambda, \quad (4.3)$$

where c_0 is an arbitrary function of r as an integration constant with respect to t . From the definition (2.25), we can determine Λ through R :

$$\Lambda = \sqrt{\bar{\chi}(R) - F(R)/c_0(r)}. \quad (4.4)$$

From Eq. (4.1), on the other hand, R' is expressed as

$$R' = \sqrt{\bar{\chi}(R)}\Lambda = \sqrt{\bar{\chi}(R)(\bar{\chi}(R) - F(R))}/c_0(r) . \quad (4.5)$$

If the right hand side of Eq. (4.5) is nonzero, we obtain a solution

$$G(R) \equiv \int \frac{dR}{\sqrt{\bar{\chi}(R)(\bar{\chi}(R) - F(R))}} = \int \frac{dr}{c_0(r)} + \phi(t), \quad (4.6)$$

which determines implicitly the dependence of R on r . Here ϕ is an arbitrary function of t as an integration constant with respect to r . The remaining equation of motion (3.3) is used to determine the time dependence of $\phi(t)$:

$$\dot{\phi} = \frac{dG(R)}{dR} \dot{R} = \frac{N}{n(Z)} \frac{1}{\sqrt{\bar{\chi}(R)}}. \quad (4.7)$$

It is noted that, if only we take $N^r = 0$ as a coordinate condition, the ansatz (4.1) is compatible with the canonical evolution and, resultantly, corresponds to a condition to pick up a special solution.

In the following, we evaluate dBB trajectories by giving explicit functional forms of $\bar{\chi}(R)$. The notations ds_{dBB} and Z_{dBB} are used for the line element of the rigid geometry described by the dBB picture and Z which is evaluated on the rigid geometry from Eq. (2.27), respectively.

Case A: $\bar{\chi}(R) \equiv F(R) \geq 0$

This is a special case where $\bar{f} = 0$ and, therefore, Eqs. (4.4) and (4.6) do not apply. Directly from Eqs. (3.3) and (3.4) with $N^r = 0$, we find

$$\dot{\Lambda} = \dot{R} = 0, \quad (4.8)$$

which means that there is no dynamical evolution. This conclusion is confirmed by observing that the integrand of Z in Eq. (2.18) or Eq. (2.27) vanishes and thus the wave function becomes constant. To keep the ansatz $\bar{\chi}(R) \equiv F(R)$, or $R'^2 = \Lambda^2 F(R)$ during the time evolution we require a condition

$$\frac{\partial}{\partial t} R' = 0. \quad (4.9)$$

If we take $R' = R$, that is, $R = \exp r$, which satisfies the consistency condition (4.9), then we obtain $\Lambda^2 = R^2 F(R)^{-1}$ and

$$ds_{dBB}^2 = -N^2 dt^2 + F(R)^{-1} dR^2 + R^2 d\Omega^2. \quad (4.10)$$

Here the lapse function N remains an arbitrary function. The Reissner-Norström-de Sitter metric corresponds to the case of the choice $N = \sqrt{F(R)}$.

Case B: $\bar{\chi}(R) \equiv 0$

First we take the integration constant in Eq. (4.3) as $c_0 = 1$. From Eq. (4.4), when $F(R) \leq 0$, Λ is given by

$$\Lambda^2 = -F(R). \quad (4.11)$$

From Eq. (4.5), we have $R' = 0$. Therefore,

$$R = \phi(t), \quad (4.12)$$

where $\phi(t)$ is an arbitrary function of t as an integration constant with the respect to r . Equations (4.11) and (4.12) show an initial configuration to pick up a dBB trajectory.

If the lapse function N is taken as

$$N = \frac{1}{\sqrt{-F(R)}} n(Z), \quad (4.13)$$

then Eq. (3.3) is reduced to $\dot{\phi} = 1$, and the dBB trajectory is

$$R(r, t) = t. \quad (4.14)$$

As a result, the line element ds_{dBB} is

$$ds_{dBB}^2 = \frac{1}{F(t)} n(Z_{dBB})^2 dt^2 - F(t) dr^2 + t^2 d\Omega^2, \quad (4.15)$$

where

$$Z_{dBB} = - \int dr R F(R) = -t F(t) R_0, \quad (4.16)$$

and R_0 is the world size. For large t , the metric (4.15) approaches to the inside geometry of the Reissner-Nordström-de Sitter black hole, which is discussed in Ref. [12], and, when $Q = \lambda = 0$, to that of the Schwarzschild black hole, which is discussed in Refs. [20,25]. These classical solutions are generalizations of the Kantowski-Sachs metric [26].

Case C: $\bar{\chi}(R) \equiv 1 - 2m/R + Q^2/R^2 \geq 0$

The ansatz reduces Eq. (2.27) into

$$Z = \int dr R^2 \left(\sqrt{\frac{\lambda}{3}} R + \frac{\sqrt{\bar{\chi}(R)}}{2} \ln \left| \frac{\sqrt{\lambda/3} R - \sqrt{\bar{\chi}(R)}}{\sqrt{\lambda/3} R + \sqrt{\bar{\chi}(R)}} \right| \right). \quad (4.17)$$

If we set $c_0(r) = \sqrt{\lambda/3}$ in Eq. (4.3), then Λ is given as

$$\Lambda = R \quad (4.18)$$

from Eq. (4.4), and Eq. (4.5) becomes

$$R' = R \sqrt{1 - \frac{2m}{R} + \frac{Q^2}{R^2}}. \quad (4.19)$$

Integrating Eq. (4.19), we obtain

$$R = x \left(1 + \frac{m}{x} + \frac{m^2 - Q^2}{4x^2} \right), \quad (4.20)$$

where $x \equiv \exp(r + \sqrt{\frac{\lambda}{3}} \phi(t))$, and $\phi(t)$ is an arbitrary function of t as an integration constant with respect to r . Equations (4.18) and (4.20) describe an initial configuration.

Next we choose a time coordinate condition

$$N = \sqrt{1 - \frac{2m}{R} + \frac{Q^2}{R^2}} n(Z) = \left(1 - \frac{m^2 - Q^2}{x^2} \right) \left(1 + \frac{2m}{x} + \frac{m^2 - Q^2}{x^2} \right)^{-1} n(Z). \quad (4.21)$$

Then Eq. (4.7) is reduced to $\dot{\phi} = 1$. Therefore, the parameter x is determined to be

$$x = a_0 \exp \left(\sqrt{\frac{\lambda}{3}} t + r \right) \equiv a(t) \rho(r), \quad (4.22)$$

where a_0 is a constant, and $a(t)$ corresponds to the scale factor of an expanding de Sitter universe with a cosmological term λ . As a result, the line element ds_{dBB} is

$$\begin{aligned} ds_{dBB}^2 &= - \left(1 - \frac{2m}{R} + \frac{Q^2}{R^2} \right) n(Z_{dBB})^2 dt^2 + R^2 (dr^2 + d\Omega^2) \\ &= - \left(\frac{1 - \frac{m^2 - Q^2}{4x^2}}{1 + \frac{m}{x} + \frac{m^2 - Q^2}{4x^2}} \right)^2 n(Z_{dBB})^2 dt^2 + a(t)^2 \left(1 + \frac{m}{x} + \frac{m^2 - Q^2}{4x^2} \right)^2 (d\rho^2 + \rho^2 d\Omega^2). \end{aligned} \quad (4.23)$$

Here we study some limiting cases. First we consider the special case $m = Q = 0$. Equation (4.23) is reduced to

$$ds_{dBB}^2 = -n(Z_{dBB})^2 dt^2 + a(t)^2(d\rho^2 + \rho^2 d\Omega^2), \quad (4.24)$$

whose asymptotic form behaves like a classical de Sitter universe. Horiguchi [27] showed that quantization of the homogeneous universe with a cosmological term gives a Bessel type wave function of order $\nu = 1/3$ (the Airy function) with the argument

$$Z = \int d\rho \rho^2 a(t)^3 = a(t)^3 V_0, \quad (4.25)$$

where V_0 is the world volume. The argument (4.25) is different from our Z , Eq. (4.17) in the absence of the second term. This is due to the difference in the order of the two procedures: quantization and reduction of the degrees of freedom. In our case, the cosmological isotropic symmetry is taken after quantization, while it is taken before quantization in Ref. [27]. The property of quantum fluctuations depends on how to construct minisuperspaces. Next we refer to the special case $m = Q \neq 0$. The asymptotic line element is

$$ds_{dBB}^2 \rightarrow -(1 + \frac{m}{x})^{-2} dt^2 + a(t)^2 (1 + \frac{m}{x})^2 (d\rho^2 + \rho^2 d\Omega^2), \quad (4.26)$$

where $x = a(t)\rho$. This is called an extreme black hole. In classical theory, extension of the Majumdar-Papetrou geometry to the cosmological black hole was discussed by Ref. [22].

Case D: $\bar{\chi}(R) \equiv \frac{1}{2}(F(R) + \sqrt{F(R)^2 + 4R^2})$

The ansatz yields

$$\Lambda^2 = \frac{1}{2}(-F(R) + \sqrt{F(R)^2 + 4R^2}) \quad (4.27)$$

and $R' = R$ from Eqs. (4.4) and (4.5), respectively. Here $c_0 = 1$ has been chosen. Thus the initial R takes a simple form

$$R = \exp(r + \phi(t)), \quad (4.28)$$

where $\phi(t)$ is an arbitrary function of t as an integration constant with respect to r .

If we take the lapse function N as

$$N^2 = \frac{\lambda}{6}(F(R) + \sqrt{F(R)^2 + 4R^2})n(Z)^2, \quad (4.29)$$

Eq. (3.3) is reduced to $\dot{\phi} = \sqrt{\frac{\lambda}{3}}$. Therefore the form of R is determined as

$$R = a_0 \exp(\sqrt{\frac{\lambda}{3}}t + r) \equiv a(t)\rho(r), \quad (4.30)$$

where a_0 is a constant. Resultantly the line element ds_{dBB} is given by

$$\begin{aligned} ds_{dBB}^2 = & -\frac{\lambda}{6}(\sqrt{F(R)^2 + 4R^2} + F(R))n(Z_{dBB})^2 dt^2 \\ & + a^2(t)\left(\frac{2d\rho^2}{\sqrt{F(R)^2 + 4R^2} + F(R)} + \rho^2 d\Omega^2\right). \end{aligned} \quad (4.31)$$

The classical limit of Eq. (4.31) shows a cosmological black hole geometry in the standard form, which contrasts with the isotropic form in the case C.

In the asymptotic region where $n_{dBB} \rightarrow 1$, or the classical limit $\hbar \rightarrow 0$, Eqs. (4.15), (4.23), and (4.31) show the same classical Reissner-Nordström-de Sitter space-time. They are different from each other merely in the coordinate system with respect to which they are described and the region which they cover in the fully extended space-time. In Eq. (4.15) of the case B, the coordinate t plays the role of the Schwarzschild radial coordinate. The associated condition $F(R) \leq 0$ shows that the corresponding classical geometry covers only the dynamical region, or one patch bounded by the horizons. On the other hand, the asymptotic form of Eq. (4.23) is described with respect to the cosmological isotropic coordinate and covers the region from the inner of the event horizon of the black hole to the

outer of the event horizon of the de Sitter universe. The transformation between the cosmological isotropic coordinate and the static one is given by

$$a\Omega\rho = \bar{r}, \quad t = \bar{t} - \sqrt{\frac{\lambda}{3}} \int \frac{1}{F(\bar{r})} \frac{\bar{r}^2}{\sqrt{\bar{r}^2 - 2m\bar{r} + Q^2}} d\bar{r}, \quad (4.32)$$

where \bar{t} and \bar{r} are the time and radial coordinates in the static Reissner-Nordström-de Sitter space-time. We find that the coordinate t plays the role of an advanced null coordinate. Alternatively, the coordinate transformation

$$a\rho = \bar{r}, \quad t = \bar{t} - \int \frac{1}{F(\bar{r})} \frac{\sqrt{F^2(\bar{r})^2 + \bar{r}^2} - F(\bar{r})}{\bar{r}} d\bar{r} \quad (4.33)$$

relates the asymptotic form of Eq. (4.31) with that of Eq. (4.15) to extend the dynamical region to the static region.

Returning our attention to quantum theory, we find that the rigid space-time picture is not covariant under general coordinate transformations. For example, under the coordinate transformation (4.33), Eq. (4.15) is not equivalent to Eq. (4.31) due to the presence of the quantum gravity factor $n(Z_{dBB})$. In other word, the quantum effect appears differently depending on the choice of coordinate systems. It may appear strange. Indeed, a quantum state on the superspace is constructed to be invariant. However, the deterministic picture labels quantum fluctuations on the superspace by introducing the new time coordinate. In this way, the dynamics is essential to quantum mechanics.

By giving the initial and coordinate conditions concretely, in this section, we obtained the various space-times which are related with each other by the coordinate transformations. For other conditions, we can also perform analytic evaluations. In Appendix, some calculations are given. As a nontrivial case, we here present a space-time with the positive constant curvature. For $m = Q = 0$, the configuration variables $(R, \Lambda) = (a(t)r, a(t)/\sqrt{1 - Kr^2})$ satisfies Eq. (3.6) when we set the shift function as $N^r = 0$. Here $K(> 0)$ is a sign of the constant curvature. The variable $a(t)$ is determined by Eq. (3.3) as

$$\dot{a} = \frac{N}{n(Z)} \sqrt{\frac{\lambda}{3} a^2 - K}. \quad (4.34)$$

When we fix the lapse function as $N = n(Z)$, the solution is

$$a(t) = \sqrt{\frac{3K}{\lambda}} \cosh \sqrt{\frac{\lambda}{3}} t. \quad (4.35)$$

In the classical limit $\hbar \rightarrow 0$, this solution corresponds to the closed de Sitter universe whose scale factor is Eq. (4.35). Now the corresponding classical geometry cannot be related with the cases A-D by any coordinate transformation.

V. SUMMARY

We have studied canonical quantization of the Einstein-Maxwell theory with a cosmological term in spherically symmetric space-time from the viewpoint of the de Broglie-Bohm interpretation. First we constructed the canonical formalism of the spherically symmetric geometry and quantized it. To resolve the operator ordering ambiguity, we followed the procedure proposed by our previous work [11]. We obtained an analytic wave function, which is the simultaneous eigenstate of the mass and electric charge operators. It is noted that, at this stage, any coordinate fixing had not been taken and the lapse and shift functions N and N^r were still arbitrary. Next the dBB interpretation of quantum mechanics was applied to the wave function, as if the Wheeler-DeWitt equation were the Schrödinger equation with zero energy. The equations of motion for the quantized space-time geometry were given as deterministic partial differential equations. Giving the ansatz to specify an initial geometry and choosing a coordinate system, we integrated the equations analytically. In the asymptotic limit, the obtained rigid metrics correspond to the classical solutions of the Einstein-Maxwell theory, or the various representations of the Reissner-Norström-de Sitter space-time and the closed de Sitter universe.

As discussed in Sec. 4 in detail, comparison between the rigid geometries as the dBB trajectories shows that the dBB trajectory picture is not covariant under coordinate transformations. When we translate a quantum state from the wave function picture to the dBB trajectory representation, we need additionally the coordinate system which describes the canonical evolution. If once we give a coordinate condition and obtain the rigid space-time picture, the metric of the space-time cannot be transformed to the others induced by the different choice of coordinate systems under any coordinate transformation. In other words, realization of the quantum world breaks the covariance under

general coordinate transformations, while a quantum state as bundle of possible rigid geometries has invariance. It may remind us the effective action at the spontaneous symmetry breaking.

In our analysis, a coordinate system is chosen a priori. However, if it has the important meaning on the quantum gravity, it must be treated as physical process fundamentally. In order to make our analysis more plausible, we have to throw probes into quantum-mechanically fluctuating space-times rather than choose a coordinate condition as a special one. The probe will play the role of the coordinate system in the sense that it set up how to measure a space-time structure. The dBB trajectory picture constructed by the information from the probe will teach us what quantum gravitational state is realized. We may think the probe and the observable information as a quantum field and the Hawking radiation, for example. The analysis on this line is needed for the conceptual problem.

There is another interest in our work from the angle of model analysis. Using the model discussed here, we can construct the quantum mechanics of extreme black holes [21], especially of cosmological extreme black holes [22], and analyze the gravitational fluctuation near the event horizon at $N^r \neq 0$ gauge analogous to the Vaidya metric, which gives us a different viewpoint of the Hawking radiation [28].

APPENDIX

Following the analysis in Sec. 4, we present other calculations of the dBB trajectory.

$$\mathbf{1. \text{ Ansatz: }} \bar{\chi}(R) \equiv -\frac{\lambda}{3}R^2$$

We give the ansatz $\bar{\chi}(R) \equiv -\frac{\lambda}{3}R^2$ and treat only the region $\bar{\chi}(R) - F(R) < 0$. Taking $c_0(r) = 1$ in Eq. (4.3), we find

$$\Lambda^2 = -\left(1 - \frac{2m}{R} + \frac{Q^2}{R^2}\right). \quad (\text{A1})$$

Here we allow Λ to become purely imaginary only if the reality of Z is reserved. From Eq. (4.6), we obtain

$$R = y\left(1 + \frac{m}{y} + \frac{m^2 - Q^2}{4y^2}\right), \quad (\text{A2})$$

where $y \equiv \exp(\sqrt{\frac{\lambda}{3}}r + \phi(t))$, and $\phi(t)$ is an arbitrary function of t as an integration constant. If we adopt a coordinate condition

$$N^2 = -y^2\left(1 + \frac{m}{y} + \frac{m^2 - Q^2}{4y^2}\right)^2 n(Z)^2, \quad (\text{A3})$$

where the evolution of the variables is traced on the purely imaginary time t , Eq. (3.3) is reduced to $\dot{\phi} = 1$, and thus $y = a_0 \exp(\sqrt{\frac{\lambda}{3}}r + t)$, where a_0 is a constant. As a result, the line element ds_{dBB} is

$$ds_{dBB}^2 = -\left(\frac{1 - \frac{m^2 - Q^2}{4y^2}}{1 + \frac{m}{y} + \frac{m^2 - Q^2}{4y^2}}\right)^2 dr^2 + y^2\left(1 + \frac{m}{y} + \frac{m^2 - Q^2}{4y^2}\right)^2 (n(Z_{dBB})^2 dt^2 + d\Omega^2). \quad (\text{A4})$$

For $N^r = 0$, we define a space-time dual transformation as

$$r \rightarrow t, \quad t \rightarrow r, \quad \Lambda \rightarrow iN \quad \text{and} \quad N \rightarrow -i\Lambda. \quad (\text{A5})$$

Under this transformation, the action (2.4) is invariant and the classical solutions (asymptotic forms) derived in Sec. 4 are related with each other:

$$\text{case A} \leftrightarrow \text{case B} \quad \text{and} \quad \text{case C} \leftrightarrow \text{Appendix}. \quad (\text{A6})$$

However the dBB trajectory representation is not covariant under the space-time dual transformation.

2. Ansatz: $\bar{\chi}(R) \equiv 0$ ($N^r \neq 0$)

In Sec. 4, we use the shift function $N^r = 0$ as a coordinate condition. Here we present an example of the nonzero shift function. From the ansatz $\bar{\chi}(R) \equiv 0$, we find $R' = 0$ and that Eq. (2.27) is reduced to

$$Z = \int dr \Lambda R \sqrt{-F(R)}. \quad (\text{A7})$$

By estimating the derivative of Z with respect to R , we have

$$\frac{\delta \Theta}{\delta R} = \Lambda (\sqrt{-F(R)} + R \frac{d}{dR} \sqrt{-F(R)}) \frac{d\Theta}{dZ}. \quad (\text{A8})$$

The equations of motion for the dBB trajectory are reduced to

$$\dot{R} = \frac{N}{n(Z)} \sqrt{-F(R)} \quad \text{and} \quad \dot{\Lambda} - (N^r \Lambda)' = \frac{N}{n(Z)} \Lambda \frac{d}{dR} \sqrt{-F(R)}. \quad (\text{A9})$$

If we set $N = n(Z)/\sqrt{-F(R)}$ and $(N^r)' = 0$, then we find $R = t$ and

$$\Lambda = G(r + \int N^r dt) \sqrt{-F(t)}, \quad (\text{A10})$$

where G is an arbitrary function. Thus the line element ds_{dBB} is

$$ds_{dBB}^2 = \frac{n(Z_{dBB})}{F(t)} dt^2 - F(t) G(r + \int N^r dt)^2 (dr + N^r dt)^2 + t^2 d\Omega^2, \quad (\text{A11})$$

where

$$Z_{dBB} = t \sqrt{-F(t)} \int dr G(r + \int N^r dt). \quad (\text{A12})$$

Equation (A11) can be related with Eq. (4.15) by the coordinate transformation $\bar{t} = t$ and $\bar{r} = \int G(\alpha) d\alpha$, where $d\alpha = dr + N^r dt$, and \bar{t} and \bar{r} denote the coordinates in the static Reissner-Norström-de Sitter metric.

-
- [1] R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation: An Introduction to Current Research*, edited by L. Witten (Wiley, New York, 1962).
 - [2] P. A. M. Dirac, *Lectures on Quantum Mechanics* (Yeshiva University, New York, 1964).
 - [3] J. A. Wheeler, in *Batelle Rencontres: 1967 Lectures in Mathematics and Physics*, edited by C. DeWitt and J. A. Wheeler (Benjamin, New York, 1968).
 - [4] B. S. DeWitt, Phys. Rev. **160**, 1113 (1967).
 - [5] K. V. Kuchař, “Canonical quantum gravity”, e-print gr-qc/9304012, 1993.
 - [6] W. Fischler, D. Morgan, and J. Polchinski, Phys. Rev. D **42**, 4042 (1990).
 - [7] K. V. Kuchař, Phys. Rev. D **50**, 3961 (1994).
 - [8] B. K. Berger, D. M. Chitre, V. E. Moncrief, and Y. Nutku, Phys. Rev. D **5**, 2467 (1972); F. Lund, *ibid.* **8**, 3247 (1973); H. A. Kastrup and T. Thiemann, Nucl. Phys. **B425**, 665 (1994); T. Thiemann, *ibid.* **B436**, 681 (1995); J. Louko and S. N. Winters-Hilt, Phys. Rev. D **54**, 2647 (1996); J. Mäkelä and P. Repo, *ibid.* **57**, 4899 (1998).
 - [9] J. Schwinger, Phys. Rev. **132**, 1317 (1963); A. Komar, Phys. Rev. D **20**, 830 (1979).
 - [10] N. C. Tsamis and R. P. Woodard, Phys. Rev. D **36**, 3641 (1987); J. L. Friedman and I. Jack, *ibid.* **37**, 3495 (1988).
 - [11] M. Kenmoku, H. Kubotani, E. Takasugi, and Y. Yamazaki, Phys. Rev. D **59**, 124004 (1999).
 - [12] M. Kenmoku, H. Kubotani, E. Takasugi, and Y. Yamazaki, Int. J. Mod. Phys. A **15**, 2059 (2000).
 - [13] D. Bohm, Phys. Rev. **85**, 166 (1952); **85**, 180 (1952).
 - [14] J. S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, England, 1987).
 - [15] P. R. Holland, *The Quantum Theory Of Motion* (Cambridge University Press, Cambridge, England, 1993).
 - [16] C. J. Isham, “Canonical quantum gravity and the problem of time”, Report Imperial/TP/91-92/25, gr-qc/9210011, 1992.
 - [17] J. J. Halliwell, Phys. Rev. D **36**, 3626 (1987).
 - [18] S. P. de Alwis and D. A. MacIntire, Phys. Rev. D **50**, 5164 (1994).
 - [19] M. Kenmoku, K. Otsuki, K. Shigemoto, and K. Uehara, Class. Quantum Grav. **13**, 1751 (1996).
 - [20] M. Kenmoku, H. Kubotani, E. Takasugi, and Y. Yamazaki, Phys. Rev. D **57**, 4925 (1998).
 - [21] J. B. Hartle and S. W. Hawking, Commun. Math. Phys. **26**, 87 (1972).
 - [22] D. Kastor and J. Traschen, Phys. Rev. D **47**, 5370 (1993).
 - [23] Y. Nambu and M. Sasaki, Prog. Theor. Phys. **79**, 96 (1988).
 - [24] A. Vilenkin, Phys. Rev. D **37**, 888 (1988).
 - [25] K. Nakamura, S. Konno, Y. Oshiro, and A. Tomimatsu, Prog. Theor. Phys. **90**, 861 (1993).
 - [26] R. Kantowski and R. Sachs, J. Math. Phys. **7**, 443 (1966).
 - [27] T. Horiguchi, Mod. Phys. Let. A **9**, 1429 (1994).
 - [28] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).