

\\

Title: Dimensionless Constants of the Fundamental Physical Interactions  
viewed by the Model of Expansive Nondecelerative Universe  
Authors: Jozef Sima, Miroslav Sukenik and Magdalena Sukenikova  
Comments: 5 pages, AmsTex  
Report-no: SS-99-03

\\

Abstract

A profound relationship between the ENU (Expansive Nondecelerative Universe) and dimensionless constants of the fundamental physical interactions is presented. The contribution corrects the Dirac presumption on a time decrease of the gravitational constant  $G$  and using simple relations it precises the values of the vector bosons  $x$  and  $y$ .

\\

### **Dimensionless Constants of the Fundamental Physical Interactions viewed by the Model of Expansive Nondecelerative Universe**

Jozef Sima, Miroslav Sukenik and Magdalena Sukenikova  
Slovak Technical University, Radlinskeho 9, 812 37 Bratislava, Slovakia

Abstract. A profound relationship between the ENU (Expansive Nondecelerative Universe) and dimensionless constants of the fundamental physical interactions is presented. The contribution corrects the Dirac presumption on a time decrease of the gravitational constant  $G$  and using simple relations it precises the values of the vector bosons  $x$  and  $y$ .

There is no problem in defining dimensionless physical constants, namely that of strong interaction  $\alpha_s$

$$\alpha_s = 1 \quad (1)$$

and electromagnetic interaction  $\alpha_e$

$$\alpha_e = \frac{e^2}{4\pi\epsilon_o \cdot \hbar \cdot c} = 7.29 \times 10^{-3} \quad (2)$$

For the constant of weak interaction  $\alpha_w$  various modes of expression are used, the simplest of which is [1]

$$\alpha_w = \frac{g_F \cdot m_P^2 \cdot c}{\hbar^3} \approx 10^{-6} \quad (3)$$

where  $g_F$  is the Fermi constant ( $g_F \sim 10^{-62} \text{ J.m}^3$ ) and  $m_P$  is the proton mass. Ambiguousness appears in the gravitational constant  $\alpha_g$  expression, the most familiar one being

$$\alpha_g = \frac{G \cdot m_P^2}{\hbar \cdot c} \approx 10^{-39} \quad (4)$$

The value of  $\alpha_g$  depends on the mass option. As a matter of tradition, the mass of proton  $m_P$  is usually taken, there is, however, no justified reason for it. On the other hand, there should be a certain mass  $m_x$ , the introduction into  $\alpha_g$  of which will be rationalizable and justifiable. Gravitational forces are far-reaching, theoretically boundless, forces. Due to the operation of the hierarchical rotational gravitational systems (HRGS), the actual gravitational effects are considered as being finite. Gravitation can manifest itself only where

density of the gravitational energy of HRGS is higher than the critical density  $\varepsilon_{crit}$ . In the ENU model using the Vaidya metrics it was shown [2] that

$$\epsilon_g = -\frac{c^4 \cdot R}{8\pi \cdot G} = -\frac{3m \cdot c^2}{4\pi \cdot a \cdot r^2} \quad (5)$$

where  $\varepsilon_g$  is the density of gravitational energy created by a body with the mass  $m$  at the distance  $r$ ,  $R$  is the scalar curvature,  $a$  is the gauge factor that at present

$$a \approx 10^{26}m \quad (6)$$

In the ENU model [2, 3]  $\varepsilon_{crit}$  is expressed as

$$\epsilon_{crit} = \frac{3c^4}{8\pi \cdot G \cdot a^2} \quad (7)$$

In the cases when

$$|\epsilon_g| = \epsilon_{crit} \quad (8)$$

using equations (5) and (7) the following relation is obtained

$$r = r_{ef(m)} = (a \cdot R_{g(m)})^{1/2} \quad (9)$$

in which  $r_{ef(m)}$  represents the effective gravitational range of a body with the mass  $m$ ,  $R_{g(m)}$  is its gravitational radius. Since for the Compton wave  $\lambda$  of the particle with the mass  $m$

$$\lambda = \frac{\hbar}{m \cdot c} \quad (10)$$

a particle having the mass  $m_x$  for which  $\lambda = r_{ef(m)}$  must exist. It thus represents the lightest particle able to gravitationally influence its environment. Relating (9) and (10) leads to

$$m_x = \left( \frac{\hbar^2}{2G \cdot a} \right)^{1/3} \quad (11)$$

At present

$$m_x \approx 10^{-28}kg \quad (12)$$

This value can be introduced into  $\alpha_g$  without additional presumptions and then

$$\alpha_g = \frac{Gm_x^2}{\hbar \cdot c} = \frac{m_x^2}{m_{Pc}^2} \approx 10^{-40} \quad (13)$$

where  $m_{Pc}$  is the Planck mass

$$m_{Pc} \approx 10^{19}GeV \quad (14)$$

In the ENU model [2,3] it was proved that

$$a = c \cdot t = \frac{2G \cdot M_u}{c^2} \quad (15)$$

where  $t$  is the cosmological time,  $M_u$  is the mass of the Universe. It follows from (11), (13) and (15)

$$\alpha_g(t)^{-2/3} \quad (16)$$

It is worth pointing out at the importance of relation (16). It brings an evidence that  $\alpha_g$  is not a true constant but a time decreasing quantity. This fact leads to some interesting consequences.

The Compton dimension of particle  $m_x$  is about  $10^{-15}$  m and corresponding  $t_x \sim 10^{-23}$  s. Consequently, for the cosmologic time  $t$  expressed in  $t_x$  units using (11), (13) and (15), relation (17) can be obtained

$$\frac{t}{t_x} \approx \frac{1}{\alpha_g} \quad (17)$$

This fact was known to Dirac who, however,  $t_x$  put equal to the atomic time. It was, however, only a coincidence since at present the Compton wavelength

of the particle  $m_x$  approaches the dimension of atomic nucleus. Since Dirac introduced the constant mass of the proton when expressing  $\alpha_g$  and at the same time relied on the validity of relation (17), he formulated a false conclusion on time decrease of the gravitational constant  $G$ . In reality,  $G$  is a true constant and  $\alpha_g$  is a time decreasing quantity (see relations 11, 13 a 16).

In the ENU model the Universe is mass-space-time (in the language of special relativity) closed. In the framework of Friedmann model for such systems it is supposed [1] that

$$M_u \approx \frac{m_P}{\alpha_g^2} \quad (18)$$

The above relation cannot be unambiguously proved, using (11), (13), (15) it can be, however, derived that

$$M_u \approx \frac{m_x}{\alpha_g^2} \quad (19)$$

In addition, it holds in Friedmann model [1] that

$$k.T \approx \left( \frac{\hbar^3 \cdot c^5}{G \cdot t^2} \right)^{1/4} \quad (20)$$

where  $T$  is the temperature of the relict radiation. For closed systems it is supposed [1] that

$$k.T_{\min} \approx \alpha_g^{1/4} \cdot m_P \cdot c^2 \quad (21)$$

Relation (21) cannot be derived, however, it follows from (11), (13) and (20) that

$$k.T \approx \alpha_g^{1/4} \cdot m_x \cdot c^2 \quad (22)$$

If in a known empirical formula [1]

$$H \approx \frac{\alpha_g \cdot m_P \cdot c^2}{\hbar} \quad (23)$$

where  $H$  is the Hubble constant, the mass  $m_P$  is substituted for  $m_x$ , relation (24) emerges which can be derived directly using (11), (13) and (15)

$$H \approx \frac{\alpha_g \cdot m_x \cdot c^2}{\hbar} \quad (24)$$

It should be worth mentioning that relations (19), (22) and (24) are in fact closely connected to dependence (17) which can be directly derived from them. It is obvious that the introduction of  $m_x$  into the “constant”  $\alpha_g$  enables to formulate known empirical formulae as self-sustaining, rationalizable and derivable relations that are or should be valid in our Universe.

Due to the time decreasing of  $\alpha_g$  (16) there had to be the time in the past when

$$\alpha_g = \alpha_s \quad (25)$$

It follows of (1), (13) and (25) that at that time

$$m_{Pc} = m_x \approx 10^{19} GeV \quad (26)$$

It really happened at the beginning of the Universe expansion ( $t \sim 10^{-43}$  s).

At about  $t \sim 10^{-40}$  s following the beginning of the expansion the quantities  $\alpha_e$  and  $\alpha_g$  were of identical value

$$\alpha_e = \alpha_g \quad (27)$$

and, consequently, treatment of (2), (13) and (27) leads to relation

$$m_x = m_{Pc} \cdot \alpha_e^{1/2} \approx 10^{18} GeV \quad (28)$$

At about  $t \sim 10^{-34}$  s following the beginning of the expansion, it held

$$\alpha_g = \alpha_w \quad (29)$$

and treating (3), (13) and (29) we get in an independent way an expression for  $m_x$

$$m_x = m_{Pc} \cdot \alpha_w^{1/2} \cong 10^{16} GeV \quad (29)$$

We supposed the values  $m_x$  represent the mass of vector bosons  $x$  and  $y$ .

Conclusions

1. The present contribution rationalizes the “constant”  $\alpha_g$  and makes possible to prove on a exact base some relations describing our Universe that have been up to now formulated as empirical postulates and formulae.

2. The contribution corrects the Dirac presumption on a time decrease of the constant  $G$ .

3. Using simple relations the contribution determines in an independent way the mass of

vector bosons  $x$  and  $y$ .

References

- [1] I. L. Rozentahl, Adv. Math. Phys. Astr. 31 (1986) 241
- [2] J. Sima, M. Sukenik, General Relativity and Quantum Cosmology,  
<http://xxx.lanl.gov/abs/qg-qc/9903090> (1999)
- [3] V. Skalsky, M. Sukenik, Astrophys. Space Sci., 178 (1991) 169